

Ecological site R011XA010ID Churning Clay 8-12 PZ ARTRW8/PSSPS

Last updated: 4/06/2020
Accessed: 05/11/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

| | |
|---|--|
| Author(s)/participant(s) | Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676 |
| Contact for lead author | Brendan Brazee, State Rangeland Management Specialist USDA-NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709 |
| Date | 03/31/2008 |
| Approved by | Kendra Moseley |
| Approval date | |
| Composition (Indicators 10 and 12) based on | Annual Production |

Indicators

- Number and extent of rills:** rills rarely occur on this site. If rills are present they are most likely to occur on slopes greater than 10 percent. The churning action of the soil obliterates any signs of surface erosion each year.
- Presence of water flow patterns:** water-flow patterns do not occur on this site due to the relatively flat slopes. If water flow patterns do develop, they are broken up by the churning action of the soil.
- Number and height of erosional pedestals or terracettes:** both are rare on this site. Do not misinterpret frost heaving for pedestals.
- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** data not available. On sites in mid-seral status, bare ground may range from 40-60 percent.
- Number of gullies and erosion associated with gullies:** none.

-
6. **Extent of wind scoured, blowouts and/or depositional areas:** usually does not occur. Some wind erosion may occur immediately following a wildfire.
-
7. **Amount of litter movement (describe size and distance expected to travel):** fine litter in the interspaces may move less than 2 feet following a significant run-off event. Cracks in the soil surface that occur during the summer and fall, trap litter. Coarse litter generally does not move.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** values should range from 4 to 6 but needs to be tested.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** structure ranges from moderate fine granular to moderate medium granular. Soil organic matter (SOM) is 1 to 2 percent. The A or A1 horizon is typically 2 inches thick.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** bunchgrasses, especially deep-rooted perennials, slow run-off and increase infiltration. Shrubs accumulate snow in the interspaces.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** may be present in areas where livestock trailing occurs. It will be most prevalent if grazing occurs when the soils are wet. The churning action of the soil tends to breakdown the compaction layer where it occurs.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: season deep-rooted perennial bunchgrasses
- Sub-dominant: tall shrubs
- Other: perennial forbs
- Additional: shallow rooted grasses
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Wyoming big sagebrush becomes decadent in the absence of fire and ungulate grazing. Mortality of grasses and forbs occurs from the shrinking and swelling of the soil.
-
14. **Average percent litter cover (%) and depth (in):** annual litter cover in the interspaces will be 5-10 percent to a depth of <0.1. Under the mature shrubs litter is greater than 0.5 inches. Fine litter falls or blows into the surface cracks in the

soil.

-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** is 450 lbs. per acre in a year with normal precipitation and temperatures. Perennial grasses produce 45-55 percent of the total, forbs 15-25 percent, and shrubs 20-40 percent.
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** medusahead is the most troublesome invasive species on this site. Others that may be found on the site include cheatgrass, bulbous bluegrass, rush skeletonweed, scotch thistle, and spotted and diffuse knapweed.
-
17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in most years.
-