

Ecological site R013XY019ID Stony Loam 16-22 PZ ARTRV/PSSPS

Last updated: 2/13/2025
Accessed: 05/12/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

| | |
|---|--|
| Author(s)/participant(s) | Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676 |
| Contact for lead author | Brendan Brazee, State Rangeland Management Specialist USDA-NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709 |
| Date | 05/14/2008 |
| Approved by | Kirt Walstad |
| Approval date | |
| Composition (Indicators 10 and 12) based on | Annual Production |

Indicators

- Number and extent of rills:** rills can occur on this site. If rills are present they are likely to occur immediately following wildfire and on slopes greater than 15%. Surface stones reduce rills.
- Presence of water flow patterns:** water-flow patterns occur on this site. When they occur, they are short and disrupted by cool season grasses, tall shrubs and surface stones. They are not extensive.
- Number and height of erosional pedestals or terracettes:** both occur on this site but are not extensive. In areas where flow patterns and/or rills are present, a few pedestals may be expected. Terracettes also occur on the site uphill from surface stones, bases of tall shrubs and large bunchgrasses.
- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** may range from 10-20 percent but additional data is needed.
- Number of gullies and erosion associated with gullies:** does not occur on this site.

6. **Extent of wind scoured, blowouts and/or depositional areas:** usually not present.
-
7. **Amount of litter movement (describe size and distance expected to travel):** fine litter in the interspaces may move up to 3 feet following a significant run-off event. Coarse litter generally does not move.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** values should range from 4 to 6 but needs to be tested.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** structure ranges from weak very fine and fine, moderate very fine, fine, medium, and coarse, and strong very fine and fine granular to weak very fine platy to weak and moderate fine and medium subangular blocky. Soil organic matter (SOM) ranges from 1 to 5 percent. Surface color can be very dark brown to very dark grayish brown to dark brown. The A or A1 horizon is typically 2 to 11 inches thick.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** bunchgrasses, especially deep-rooted perennials, slow run-off and increase infiltration. Tall shrubs catch blowing snow in the interspaces.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** not present.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: cool season deep-rooted perennial bunchgrasses
- Sub-dominant: tall shrubs-perennial forbs
- Other: shallow rooted perennial bunchgrasses
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** mountain big sagebrush and antelope bitterbrush will become decadent in the absence of normal fire frequency and ungulate grazing. Grass and forb mortality will occur as tall shrubs increase.
-
14. **Average percent litter cover (%) and depth (in):** additional litter cover data is needed but is expected to be 15-30 percent to a depth of 0.1 inches. Under mature shrubs litter is >0.5 inches deep and is 90-100 percent ground cover.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** is 1100 pounds per acre (1232 kilograms per hectare) in a year with normal temperatures and precipitation.

Perennial grasses produce 55-65 percent of the total production, forbs 15-25 percent and shrubs 25-35 percent.

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: includes cheatgrass, Kentucky bluegrass, leafy spurge, and yellow salsify.
-
17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in most years.
-