

## Ecological site R024XY114OR SODIC LAKE TERRACE

Accessed: 05/11/2025

---

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Jeff Repp
Contact for lead author	State Rangeland Management Specialist for NRCS Oregon
Date	11/30/2016
Approved by	Bob Gillaspy
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

1. **Number and extent of rills:** None.
- 

2. **Presence of water flow patterns:** None, except following extremely high intensity storms and/or ponding in the spring when short (less than 1 meter) flow patterns may appear. Minimal evidence of past or current soil deposition or erosion.
- 

3. **Number and height of erosional pedestals or terracettes:** None.
- 

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 5-15%.
- 

5. **Number of gullies and erosion associated with gullies:** None.
- 

6. **Extent of wind scoured, blowouts and/or depositional areas:** None. Wind erosion hazard is slight to moderate.
- 

7. **Amount of litter movement (describe size and distance expected to travel):** Litter size is Small/Fine. Litter movement is limited, minimal, and short, associated with water flow patterns following extremely high intensity storms.

Litter also may be moved during intense wind storms.

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Site is Slightly to Moderately resistant to erosion. Stability class (Herrick et al. 2001) anticipated to be 3-6 at surface under perennial vegetation. Stability class at surface in the interspaces is anticipated to be less than or equal to that under perennial vegetation.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Surface layer structure is moderate coarse granular to weak medium platy. The A horizon has a dry color of 4 - 7 and is 2 - 5 inches thick. The Soil Organic Matter (SOM) content is low (0.5 to 3.0%).
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Plant foliar cover and basal cover with small gaps between plants should reduce raindrop impact and slow overland flow, providing increased time for infiltration to occur. High herbaceous vegetation on this site will retain more water from precipitation. Significant ground cover (60-80%) and gentle slopes (0-3%) effectively limit rainfall impact and overland flow.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Rhizomatous grass > Moderate to deep rooted bunchgrasses
- Sub-dominant: Shrubs
- Other: Other perennial grasses > perennial forbs
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Grasses will nearly always show some mortality and decadence. Normal decadence and mortality expected on other plants.
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Low 600 lbs/acre, Representative Value 800 lbs/acre, High 1000 lbs/acre
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize**

degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is **NOT** expected in the reference state for the ecological site: Annuals (Cheatgrass, Medusahead, Foxtail Barley, Mustards, and other forbs) invade sites that have lost deep rooted perennial grass functional groups. Perennial Pepperweed may also invade.

---

17. **Perennial plant reproductive capability:** All species should be capable of reproducing annually.
-