

## Ecological site R025XY414UT Mountain Mahogany Thicket (Curl-leaf Mountain Mahogany)

Accessed: 05/11/2025

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Brock Benson (NRCS), Shane A. Green (NRCS), Robert D. Stager (BLM), Mike Gates (BLM), Tyler Staggs (BLM), Alan Bass (BLM).
Contact for lead author	shane.green@ut.usda.gov
Date	03/30/2007
Approved by	Shane A. Green
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

- 1. Number and extent of rills:** Some rills present. Rill development may increase following large storm events, but should begin to heal during the following growing season. Frost heaving will accelerate recovery. Rill development may increase when run inflow enters site from other sites that produce large amounts of runoff (i.e. steeper sites, slickrock, rock outcrop).
- 2. Presence of water flow patterns:** Water flow patterns are common. Some are long (15-20'). They are generally very widely spaced (about 20-30' apart). Flow patterns occur in low places associated with microtopography commonly occurring on this site.
- 3. Number and height of erosional pedestals or terracettes:** Plants may have small pedestals (1-3") where they are adjacent to water flow patterns, but without exposed roots. Terracettes should be few and stable. Terracettes should be small (1-3") and show little sign of active erosion. Some plants may appear to have a pedestal but rather than be formed by erosion, they are the result of litter and soil accumulating at plant bases, forming the appearance of a pedestal.
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 15-30% bare ground (soil with no protection from raindrop impact). Very few if any bare spaces of greater than 1 square foot. In general, bare ground increases as production decreases. As species composition of shrubs relative to grasses increases, bare ground is likely to increase. Poorly developed biological soil crust that is susceptible to erosion from raindrop impact should be recorded as bare ground.

- 
5. **Number of gullies and erosion associated with gullies:** No gullies present.
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:** Very minor evidence of active wind-generated soil movement. Wind scoured (blowouts) and depositional areas are rarely present. If present they have muted features and are mostly stabilized with vegetation and/or biological crust.
- 
7. **Amount of litter movement (describe size and distance expected to travel):** Most litter resides in place but on steep slopes (>30%), at least half of the litter is likely to be transported downhill by wind or water short. Litter rarely moves more than 1-2' to next obstruction. Leaves, stems, and small twigs will accumulate at plant bases, against rocks, in soil depressions, or against larger woody litter. Woody litter is not likely to move.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** This site should have an erosion rating of 4 to 5 under plant canopies and a rating of 3 to 4 in the interspaces with an average rating of 4 using the soil stability kit test.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** A1--0 to 8 inches; brown (10YR 4/3) sandy loam, dark brown (10YR 3/3) moist; weak very fine granular structure; soft, very friable; 10 percent angular sandstone pebbles and flagstones; many roots; medium acid; clear wavy boundary. (5 to 15 inches thick)
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Bunchgrasses, rhizomatous grasses, and shrubs provide nearly continuous canopy cover, which effectively intercepts rainfall and protects soil from erosion. Extensive root systems provide ample organic matter to the soil, resulting in very high water-holding capacity. Infiltration rates vary with soil type. Runoff is expected to be minimal.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** A compaction layer is not expected.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Dominant: Sprouting shrubs (curl-leaf mountain mahogany, mountain snowberry) > Perennial cool-season bunchgrasses (Idaho fescue, bluebunch wheatgrass)
- Sub-dominant: Sub-dominant: Non-sprouting shrubs (mountain big sagebrush) = Tall perennial bunchgrass (basin wildrye)
- Other: Other: Perennial forbs > other perennial grasses > other shrubs
- Additional: Disturbance regime includes fire, drought, and insects. Assumed fire cycle of 50-80 years.

Dominance is based on average annual production, air dry weight: Sprouting shrubs > Perennial bunchgrasses > non-sprouting shrubs > native perennial and annual forbs.

Functional/structural groups may appropriately contain non-native species if their ecological function is the same as the native species in the reference state (e.g. Crested wheatgrass, intermediate wheatgrass, Siberian Wheatgrass etc.)

Following a recent disturbance such as drought or fire that remove the woody vegetation, forbs and perennial grasses (herbaceous species) may dominate the community. These conditions would reflect a functional community phase within the reference state.

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** During years with average to above average precipitation, there should be very little recent mortality or decadence apparent in either the shrubs or grasses. Some mortality of bunchgrass and other shrubs may occur during very severe (long-term) droughts. There may be partial mortality of individual bunchgrasses and shrubs during less severe drought and toward the end of the fire cycle. Long-lived species dominate the site. Open spaces from disturbance are quickly filled by new plants through seedlings and asexual reproduction (tillering).
- 

14. **Average percent litter cover (%) and depth ( in):** Litter cover includes litter under plants. Most litter will be fine (herbaceous) litter. Litter will be concentrated under plant canopies and sparser between plant canopies, with an average cover of 15-30% and an average depth of 0.5-1 inches. Litter cover may increase following years with favorable growing conditions. Excess litter may accumulate in absence of disturbance. Vegetative production may be reduced if litter cover exceeds 40%.
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 1100-1200 lbs/acre.  
Even the most stable communities exhibit a range of production values. Production will vary between communities and across the MRLA. Production will differ across the MLRA due to the naturally occurring variability in weather, soils, and aspect. The biological processes on this site are complex; therefore, representative values are presented in a land management context.
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Cheatgrass, Russian thistle, Utah juniper
- 

17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce sexually or asexually, except in drought years. Density of plants indicates that plants reproduce at level sufficient to fill available resource. Within capability of site there are no restrictions on seed or vegetative reproductive capacity.
-