

## Ecological site R027XY002NV MOIST FLOODPLAIN

Last updated: 6/03/2024  
Accessed: 05/12/2025

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	GK BRACKLEY
Contact for lead author	State Rangeland Management specialist
Date	06/20/2006
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

1. **Number and extent of rills:** None
- 

2. **Presence of water flow patterns:** None
- 

3. **Number and height of erosional pedestals or terracettes:** None
- 

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare Ground  $\pm$  20%; surface rock fragments minimal; shrub canopy less than 3% (willows, rose); foliar cover of perennial herbaceous plants  $\pm$  80%.
- 

5. **Number of gullies and erosion associated with gullies:** Gullies are rare to common depending on severity of associated stream channel entrenchment. Gullies and head cuts are healing or stable.
- 

6. **Extent of wind scoured, blowouts and/or depositional areas:** None
-

7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage of grasses and annual & perennial forbs) is only expected to move during periods of flooding by adjacent streams. Persistent litter (large woody material) will remain in place except during peak flooding periods.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values will range from 4 to 6. (To be field tested.)
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface structure is platy, subangular blocky or granular. Soil surface colors are very dark and have mollic epipedons. Organic carbon can range from 2 to 3.5 percent for much of the upper 20 inches. (OM values derived from lab characterization data.)
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Deep-rooted perennial herbaceous bunchgrasses slow runoff and increase infiltration. Tall stature and relatively coarse foliage of wildrye and associated litter break raindrop impact and provide opportunity for snow catch and snow accumulation on site.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None - Platy subsurface layers are not to be interpreted as compaction.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Reference Plant Community: Tall statured, deep-rooted, cool season, perennial bunchgrasses = rhizomatous, cool season, perennial bunchgrasses. (By above ground production)
- Sub-dominant: Deep-rooted, cool season, perennial forbs > shallow-rooted, cool season, perennial bunchgrasses and grass-like plants > fibrous, shallow-rooted, cool season, perennial forbs > tall shrubs (willow). (By above ground production)
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Herbaceous species show minimal mortality. Dead branches within individual shrubs are common and standing dead shrub canopy material may be as much as 15% of total woody canopy.
- 
14. **Average percent litter cover (%) and depth ( in):** Within plant interspaces ( $\pm$  80%) and depth (1 to 3 in.)
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** For normal or average growing season (September thru July)  $\pm$  2500 lbs/ac; Spring flooding significantly

affects total production.

---

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Saltcedar, quackgrass, foxtail barley, thistle, annual kochia, hoary cress, tall whitetop and annual, warm-season grasses are invaders on this site. Willow and rubber rabbitbrush are increasers on this site.
- 

17. **Perennial plant reproductive capability:** All functional groups should reproduce in most years.
-