

Ecological site R027XY072NV GRANITIC SLOPE 10-12 P.Z.

Last updated: 6/03/2024
Accessed: 05/13/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 027X–Fallon-Lovelock Area

Physiography

Found in the Great Basin Section of the Basin and Range Province of the Intermontane Plateaus this area is characterized by isolated uplifted fault block mountain ranges trending north to south that are separated by broad, hydrologically closed basins. The entire area occurs in the rain-shadow of the Sierra Nevada mountains and is influenced by Pleistocene Lake Lahontan which reached its most recent high stand about 12,000 years ago. There is substantial evidence suggesting the western Great Basin has been the site of pluvial-interpluvial cycles for at least the past two million years.

The mountains and valleys are dissected by the Humboldt, Truckee, Carson, and Walker Rivers and their tributaries, all of which terminate within MLRA 27. Extensive playas can be found throughout this area and are the result of drying of ancient Lake Lahontan. Elevation generally ranges from 3,300 to 5,900 feet (1,005 to 1,800 meters) in valleys, but on some mountain peaks it is more than 7,870 feet (2,400 meters).

Geology

Landforms and soils of this MLRA have been heavily influenced by fluctuating lake level over the last 40,000 years. There is a level line evident on the higher slopes marking the former extent of glacial Lake Lahontan. Almost half of this area has surface deposits of alluvial valley fill influenced by lacustrine sediment. The rest has andesite and basalt rocks of different ages. Mesozoic and Tertiary intrusives are concentrated along the western border of the area, and Lower Volcanic Rocks (17 to 43 million years old) are common on the eastern side of the area. Also, some scattered outcrops of Mesozoic sedimentary and volcanic rocks and tuffaceous sedimentary rocks are in the mountains within the interior of this MLRA.

Climate

The average annual precipitation is 5 to 10 inches (125 to 255 millimeters) in most of the area but is as much as 19 inches (485 millimeters) on high mountain slopes. Most of the rainfall occurs as high-intensity, convective thunderstorms during the growing season. The amount of precipitation is very low from summer to midautumn. The precipitation in winter occurs mainly as snow. The average annual temperature is 43 to 54 degrees F (6 to 12 degrees C). The freeze-free period averages 155 days and ranges from 110 to 195 days, decreasing in length with elevation.

Water

The amount of precipitation is very low, and water for irrigation is obtained principally from diversions on the four large rivers in the area and from water stored in the Lahontan, Rye Patch, and Weber Reservoirs. Pyramid Lake and Walker Lakes are terminal lakes for the Truckee and Walker Rivers, respectively. Much of the annual flow of both rivers is diverted for irrigation, causing lake levels to fall and levels of dissolved salts to increase causing problems for the native Lahontan cutthroat trout.

Soils

The dominant soil orders are Aridisols and Entisols. The soils in the area are predominantly a mesic temperature

regime, aridic moisture regime, and have a mixed mineralogy. They are generally well drained, loamy or sandy, commonly skeletal, and shallow to very deep. Accumulation of salts, tufa deposits, and eolian sediments with soluble salts over lacustrine deposits influence most of the soils in the basin landforms of this MLRA. Soils on bedrock-controlled landforms are typically comprised of volcanic or tuffaceous sedimentary colluvium over residuum.

Biological Resources

This area supports extensive areas of salt-desert shrub vegetation. Shadscale and Bailey's greasewood are widespread, occurring both individually and together. Grasses are generally sparse, although Indian ricegrass is prominent, especially on the sandy soils. Fourwing saltbush, winterfat, spiny hopsage, wolfberry, ephedra, dalea, and bud sagebrush are common shrubs. Basin wildrye, creeping wildrye, alkali sacaton, saltgrass, black greasewood, rubber rabbitbrush, and big saltbush are important plants on saline bottom lands and terraces. A few marsh areas support cattail, bulrushes, sedges, and rushes. Big sagebrush, along with scattered Utah juniper and singleleaf pinyon, is associated with Thurber needlegrass, desert needlegrass, Sandberg bluegrass, and squirreltail on the higher elevation piedmont slopes and mountains.

Ecological site concept

The Granitic Slope 10-12 P.Z. site occurs on mountains and hills on all aspects. Slope gradients of 4 to 50 percent are typical. Elevations are 4500 to 6800 feet. The soils are shallow and well drained. The soils have formed in residuum from granitic rock sources. The soil surface is coarse textured and normally very gravelly. An argillic horizon occurs within 10 inches of the surface.

Associated sites

R027XY058NV	LOAMY 10-12 P.Z. Soils not from granitic rock sources; more productive site
-------------	---

Similar sites

R027XY065NV	GRANITIC SLOPE 8-10 P.Z. ACSP12 dominant grass
R027XY088NV	GRANITIC LOAM 10-12 P.Z. More productive site
R027XY073NV	GRANITIC SLOPE 12-14 P.Z. More productive site; ARTRW8 rare to absent
R027XY067NV	GRANITIC LOAM 8-10 P.Z. ACSP12 dominant grass
R027XY007NV	LOAMY SLOPE 8-10 P.Z. Soils not from granitic rock sources
R027XY054NV	LOAMY SLOPE 10-12 P.Z. Soils not from granitic rock sources

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> (2) <i>Artemisia tridentata</i> ssp. <i>vaseyana</i>
Herbaceous	(1) <i>Achnatherum thurberianum</i>

Physiographic features

The Granitic Slope 10-12 P.Z. site occurs on mountains and hills on all aspects. Slope gradients of 4 to 50 percent are typical. Elevations are 4500 to 6800 feet.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Mountain
Runoff class	Medium to very high
Elevation	1,372–2,073 m
Slope	4–50%
Water table depth	183 cm
Aspect	Aspect is not a significant factor

Climatic features

The climate is semiarid with cool, moist winters and warm, dry summers. Average annual precipitation is 10 to 12 inches. Mean annual temperatures are 42 to 53 degrees F. The average growing season is about 90 to 120 days. There is no climate station available for this site.

Table 3. Representative climatic features

Frost-free period (average)	120 days
Freeze-free period (average)	
Precipitation total (average)	305 mm

Influencing water features

There are no influencing water features associated with this site.

Soil features

The soils associated with this site are shallow and well drained. These soils have formed in residuum from granitic rock sources. The soil surface is coarse textured and normally very gravelly. An argillic horizon occurs within 10 inches of the surface. Available water capacity is very low. Coarse textured surface soils provide moderately slow to moderate infiltration. Potential for sheet and rill erosion is moderate. The soil series associated with this site include: Acrelane and Wedekind.

Table 4. Representative soil features

Parent material	(1) Residuum–granite (2) Colluvium–granite (3) Residuum–volcanic rock
Surface texture	(1) Very gravelly coarse sandy loam (2) Very gravelly sandy clay loam (3) Gravelly sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately slow to moderate
Soil depth	25–51 cm
Surface fragment cover <=3"	14–33%
Surface fragment cover >3"	2–8%
Available water capacity (0–101.6cm)	3.05–7.37 cm

Calcium carbonate equivalent (0-101.6cm)	0%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	6.1–7.8
Subsurface fragment volume <=3" (Depth not specified)	14–33%
Subsurface fragment volume >3" (Depth not specified)	2–14%

Ecological dynamics

As ecological condition declines, overstory shrubs such as big sagebrush, Anderson's peachbrush, and Douglas' rabbitbrush increase as Sandberg's bluegrass and bottlebrush squirreltail increase in the understory. With further site degradation squirreltail and bluegrass also decline. After wildfire, Douglas' rabbitbrush, littleleaf horsebrush, Anderson's peachbrush, spiny hopsage and green ephedra often dominate the aspect. Species likely to invade this site are annuals such as cheatgrass and filaree.

Fire Ecology:

The fire return interval for Wyoming big sagebrush communities ranges from 10 to 70 years. Fire is the principal means of renewal for decadent stands of Wyoming big sagebrush.

Wyoming big sagebrush is killed by fire and establishes after fire from a seedbank; from seed produced by remnant plants that escaped fire; and from plants adjacent to the burn that seed in.

Mountain big sagebrush is highly susceptible to injury from fire. It is often top-killed by fire and will not resprout.

Spiny hopsage is considered to be somewhat fire tolerant and often survives fires that kill sagebrush. Mature spiny hopsage generally sprout after being burned. Spiny hopsage is reported to be least susceptible to fire during summer dormancy.

Thurber's needlegrass is classified as moderately resistant, but depending on season of burn, phenology, and fire severity, this perennial bunchgrass is moderately to severely damaged by fire. Early season burning is more damaging to this needlegrass than late season burning.

Indian ricegrass can be killed by fire, depending on severity and season of burn. Indian ricegrass reestablishes on burned sites through seed dispersed from adjacent unburned areas.

State and transition model

1: Reference State

Community Phase 1.1

Thurber's needlegrass dominate, big sagebrush sub-dominant. Associated species include spiny hopsage, Indian ricegrass and forbs. Approximate ground cover (basal & ground) is 20-30%.

Community Phase 1.2

Big sagebrush decreases
Perennial bunchgrasses and native forbs increase

Community Phase 1.3

Big sagebrush and other shrubs increase
Perennial herbaceous understory is reduced

T1A

State 2

Community Phase 2.1

Thurber's needlegrass dominate, big sagebrush sub-dominant. Associated species include spiny hopsage, Indian ricegrass and forbs. Approximate ground cover (basal & ground) is 20-30%. Annual non-native species present

Community Phase 2.2

Perennial bunchgrasses increase. Big sagebrush decreases. Sprouting shrubs stable to increasing
Annual non-native species stable to increasing

Community Phase 2.3 (At Risk)

Big sagebrush increases and become dominant
Thurber's needlegrass decrease, bluegrass and squirreltail increase.
Annual non-native species stable to increasing

T2A R3A

T2B

State 3

Community Phase 3.1

Decadent big sagebrush and rabbitbrush dominate overstory
Deep-rooted perennial bunchgrasses trace to absent, bluegrass may or may not be present or dominant
Annual non-native species increase
Bare ground increases

3.1A

3.2A

Community Phase 3.2

Shallow rooted bunchgrass dominates site
Annual non-native species may be present to increasing, but are not dominant
Trace amounts of sagebrush, rabbitbrush, spiny hopsage may be present

State 4

Community Phase 4.1

Annual non-natives, cheatgrass tansy mustard, etc., dominate site

4.1A

4.2A

Community Phase 4.2

Broom snakeweed, rabbitbrush, shadscale, and/or sagebrush dominate overstory
Annual non-natives dominate understory. Bare ground increasing.

T3A

Reference State 1.0: The Reference State 1.0 is a representative of the natural range of variability under pristine conditions. State dynamics are maintained by interactions between climatic patterns and disturbance regimes. Negative feedbacks enhance ecosystem resilience and contribute to the stability of the state. These include the presence of all structural and functional groups, low fine fuel loads, and retention of organic matter and nutrients. Plant community phase changes are primarily driven by fire, periodic drought and/or insect or disease attack.

Community Phase 1.1:

Wyoming/mountain big sagebrush and Thurber's needlegrass dominate the site. Indian ricegrass and spiny hopsage are important associated species. Forbs are present but not abundant. This site is tolerant of dry conditions, but prolonged drought will result in an overall decline, with possible mortality, in the plant community.

Community Phase Pathway 1.1a: Fire would decrease or eliminate the overstory of sagebrush and allow the perennial bunchgrasses to dominate the site. Fires would typically be small and patchy due to low fine fuel loads.

Community Phase Pathway 1.1b: Natural regeneration over time and lack of disturbance such as fire allows for sagebrush to increase and become decadent. Long-term drought, herbivory, or combinations of these would cause a decline in perennial bunchgrasses and fine fuels and lead to a reduced fire frequency allowing big sagebrush to dominate the site.

Community Phase 1.2: This community phase is characteristic of a post-disturbance, early seral community phase. Thurber's needlegrass and other perennial grasses dominate. Depending on fire severity patches of intact sagebrush may remain.

Community Phase Pathway 1.2a: Absence of disturbance over time coupled with natural regeneration allows sagebrush to increase. Patches of mature sagebrush required for a seed source are important for recovery to community phase 1.1.

Community Phase 1.3:

Big sagebrush increases in the absence of disturbance. Decadent sagebrush dominates the overstory and the deep-rooted perennial bunchgrasses in the understory are reduced either from competition with shrubs and/or from herbivory.

Community Phase 1.3a: Fire decreases or eliminates overstory of sagebrush and allow the perennial bunchgrasses to dominate. Fires would typically be low severity resulting in a mosaic pattern due to low fine fuel loads. A fire following an unusually wet spring or a change in management favoring an increase in fine fuels, may be more severe and reduce sagebrush cover to trace amounts. A severe infestation of Aroga moth could also cause a large decrease in sagebrush within the community, giving a competitive advantage to the perennial grasses and forbs.

Community Phase Pathway 1.3b: A low severity/patchy fire reduces the sagebrush overstory and create a sagebrush/grass mosaic with sagebrush and perennial bunchgrasses co-dominant.

T1A: Transition from Reference State 1.0 to Current Potential State 2.0

Trigger: This transition is caused by the introduction of non-native annual weeds, such as cheatgrass and mustards

Slow variables: Over time the annual non-native plants will increase within the community.

Threshold: Any amount of introduced non-native species causes an immediate decrease in the resilience of the site. Annual non-native species cannot be easily removed from the system and have the potential to significantly alter disturbance regimes from their historic range of variation.

Current Potential State 2.0:

This state is similar to the Reference State 1.0. Ecological function has not changed, however the resiliency of the state has been reduced by the presence of non-natives. Non-natives may increase in abundance but will not become dominant within this State. These non-natives can be highly flammable and can promote fire where historically fire had been infrequent. Negative feedbacks enhance ecosystem resilience and contribute to the stability of the state. These feedbacks include the presence of all structural and functional groups, low fine fuel loads and retention of organic matter and nutrients. Positive feedbacks decrease ecosystem resilience and stability of the state. These include the non-natives' high seed output, persistent seed bank, rapid growth rate, ability to cross pollinate and adaptations for seed dispersal.

Community Phase 2.1: Big sagebrush and Thurber's needlegrass dominate the site. This community phase is similar to reference plant community 1.1 with non-native annual species present.

Community Phase Pathway 2.1a: Fire reduces the shrub overstory and allows for perennial bunchgrasses to dominate the site. Fires are typically low severity resulting in a mosaic pattern due to low fuel loads. A fire following an unusually wet spring or a change in management favoring an increase in fine fuels, may be more severe and reduce sagebrush cover to trace amounts. A severe infestation of Aroga moth could also cause a large decrease in sagebrush. In sagebrush cover, reducing competition with perennial grasses and forbs. Annual non-native species are likely to increase after fire.

Community Phase Pathway 2.1b: Natural regeneration over time and lack of disturbance such as fire allows for sagebrush to increase and become decadent. Chronic drought reduces fine fuels and leads to a reduced fire frequency allowing big sagebrush to dominate the site. Inappropriate grazing management reduces the perennial bunchgrass understory; conversely Sandberg bluegrass may increase in the understory depending on grazing management.

Community Phase 2.2: This community phase is characteristic of a post-disturbance, early seral community phase. Thurber's needlegrass and other perennial grasses dominate. Big sagebrush is present in trace amounts. Depending on fire severity or intensity of Aroga moth infestations, patches of intact sagebrush may remain. Rabbitbrush may be sprouting. Annual non-native species generally respond well after fire and may be stable or increasing within the community.

Community Phase Pathway 2.2a: Natural regeneration over time and lack of disturbance and/or grazing management that favors the establishment and growth of sagebrush allows the shrub component to recover. The establishment of big sagebrush can take many years and is dependent on multiple years of favorable weather conditions.

Community Phase 2.3 (at risk): This community is at risk of crossing a threshold to another state. Sagebrush dominates the overstory and perennial bunchgrasses in the understory are reduced, either from competition with shrubs or from inappropriate

grazing management, or both. Rabbitbrush may be a significant component. Sandberg bluegrass may increase and become co-dominant with deep rooted bunchgrasses. Annual non-natives species may be stable or increasing due to lack of competition with perennial bunchgrasses. This site is susceptible to further degradation from excessive grazing, prolonged drought, and/or fire.

Community Phase Pathway 2.3a: A change in grazing management that decreases shrubs would allow the perennial bunchgrasses in the understory to increase. Brush management with minimal soil disturbance would also decrease sagebrush and release the perennial understory. Low intensity/patchy fire may create a sagebrush-grass mosaic. Annual non-native species are present and may increase in the community.

Community Phase Pathway 2.3b: Fire would decrease or eliminate the overstory of sagebrush and allow the perennial bunchgrasses to dominate the site. A fire following an unusually wet spring or a change in management may be more severe and reduce sagebrush cover to trace amounts. A severe infestation of Argo moth could also cause a large decrease in sagebrush within the community, giving a competitive advantage to the perennial grasses and forbs.

T2A: Transition from Current Potential State 2.0 to Shrub State 3.0

Trigger: Inappropriate, long-term grazing of perennial bunchgrasses during the growing season favors an increase in sagebrush.

Slow variables: Long term decrease in deep-rooted perennial grass density.

Threshold: Loss of deep-rooted perennial bunchgrasses changes spatial and temporal nutrient cycling and nutrient redistribution, and reduces soil organic matter.

T2B: Transition from Current Potential State 2.0 to Annual State 4.0

Trigger: To Community Phase 4.1: Severe fire and/or soil disturbing treatments. To Community

Phase 4.2: Inappropriate grazing management that favors shrubs in the presence of non-native species.

Slow variables: Increased production and cover of non-native annual species.

Threshold: Loss of deep-rooted perennial bunchgrasses and shrubs truncates, spatially and temporally, nutrient capture and cycling within the community. Increased, continuous fine fuels from annual non-native plants modify the fire regime by changing intensity, size and spatial variability of fires.

Shrub State 3.0:

This state is a product of many years of heavy grazing during time periods harmful to perennial bunchgrasses. Sandberg bluegrass may increase with a reduction in deep rooted perennial bunchgrass competition and may become the dominant grass or the herbaceous understory may be completely eliminated. Sagebrush cover exceeds site concept and may be decadent, reflecting stand maturity and lack of seedling establishment due to competition with mature plants. The shrub overstory dominates site resources such that soil water, nutrient capture, nutrient cycling and soil organic matter are temporally and spatially redistributed. Bare ground may be significant with soil redistribution occurring between interspace and canopy locations.

Community Phase 3.1:

Big sagebrush dominates overstory and rabbitbrush may be a significant component. Deep-rooted perennial bunchgrasses may be present in trace amounts or absent. Sandberg bluegrass may dominate the understory. Annual nonnative species are present and may be co-dominant. Bare ground is significant.

Community Phase Pathway 3.1a: Fire, heavy fall grazing causing mechanical damage to shrubs, and/or brush treatments with minimal soil disturbance, will greatly reduce the overstory shrubs to trace amounts and allow Sandberg bluegrass to dominate the site.

Community Phase 3.2:

Bluegrass dominates the site; annual non-native species may be present but are not dominant. Trace amounts of sagebrush may be present. Sprouting shrubs such as Nevada ephedra, Utah serviceberry, Anderson's peachbrush or rabbitbrush may be dominant.

Community Phase Pathway 3.2a: Time and lack of disturbance and/or grazing management that favors the establishment and growth of sagebrush allows the shrub component to recover. The re-establishment of big sagebrush can take many years.

T3A: Transition from Shrub State 3.0 to Annual State 4.0

Trigger: Severe/repeated fire and/or soil disturbing treatments. Possible soil disturbing treatments include attempted restoration with drought tolerant perennials, such as crested wheatgrass. Restoration attempts causing soil disturbance will likely initiate a transition to an annual state. Probability of success very low. Inappropriate grazing management in the presence of annual non-native species.

Slow variables: Increased production and cover of non-native annual species.

Threshold: Increased, continuous fine fuels modify the fire regime by changing intensity, size and spatial variability of fires.

Changes in plant community composition and spatial variability of vegetation due to the loss of perennial bunchgrasses and sagebrush truncate energy capture spatially and temporally thus impacting nutrient cycling and distribution.

R3A: Restoration from Shrub State 3.0 to Current Potential State 2.0

Brush management with minimal soil disturbance, coupled with seeding of deep rooted perennial native bunchgrasses.

Probability of success very low.

Annual State 4.0:

This community is characterized by the dominance of annual non-native species such as cheatgrass and tansy mustard in the understory. Sprouting shrubs such as rabbitbrush, ephedra, etc. may be present, but are not contributing to ecological site function. Spatial distribution of nutrient and water resources are being controlled by the dominance of non-native annuals.

Community Phase 4.1

Annual non-native plants such as cheatgrass or tansy mustard dominate the site. Rabbitbrush may or may not be present.

Community Phase Pathway 4.1a: Time and lack of fire allows for the sagebrush to establish. Probability of sagebrush establishment is extremely low.

Community Phase 4.2: Sprouting shrubs such as spiny hopsage and Rabbitbrush dominate overstory. Big sagebrush may be a minor component. Annual non-native species dominate understory. Trace amounts of desirable bunchgrasses may be present. Bare ground is significant.

Community Phase Pathway 4.2a: Fire eliminates shrubs and allows for annual non-native species to dominate the site.

State 1
Reference Plant Community

Community 1.1
Reference Plant Community

The reference plant community is dominated by Thurber's needlegrass and Wyoming big sagebrush. Potential vegetative composition is about 60% grasses, 10% forbs and 30% shrubs. Approximate ground cover (basal and crown) is 20 to 30 percent.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	269	404	538
Shrub/Vine	135	202	269
Forb	45	67	90
Total	449	673	897

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Primary Perennial Grasses			282–404	
	Thurber's needlegrass	ACTH7	<i>Achnatherum thurberianum</i>	269–336	–
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	13–67	–
2	Secondary Perennial Grasses			13–54	
	desert needlegrass	ACSP12	<i>Achnatherum speciosum</i>	3–20	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	3–20	–
	basin wildrye	LECI4	<i>Leymus cinereus</i>	3–20	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	3–20	–
Forb					
3	Perennial Forbs			34–101	
	arrowleaf balsamroot	BASA3	<i>Balsamorhiza sagittata</i>	3–27	–
	tapertip hawksbeard	CRAC2	<i>Crepis acuminata</i>	3–27	–
	buckwheat	ERIOG	<i>Eriogonum</i>	3–27	–
	lupine	LUPIN	<i>Lupinus</i>	3–27	–
Shrub/Vine					
4	Primary Shrubs			114–202	
	mountain big sagebrush	ARTRV	<i>Artemisia tridentata</i> ssp. <i>vaseyana</i>	50–84	–
	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i>	50–84	–
	spiny hopsage	GRSP	<i>Grayia spinosa</i>	13–34	–
5	Secondary Shrubs			13–67	
	Utah serviceberry	AMUT	<i>Amelanchier utahensis</i>	3–20	–
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	3–20	–
	mormon tea	EPVI	<i>Ephedra viridis</i>	3–20	–
	desert peach	PRAN2	<i>Prunus andersonii</i>	3–20	–

Animal community

Livestock Interpretations:

This site is suitable for livestock grazing. Grazing management should be keyed to Thurber's needlegrass production. Thurber's needlegrass species begin growth early in the year and remain green throughout a relatively long growing season. This pattern of development enables animals to use Thurber's needlegrass when many other grasses are unavailable. Cattle prefer Thurber's needlegrass in early spring before fruits have developed as it becomes less palatable when mature. Thurber's needlegrasses are grazed in the fall only if the fruits are softened by rain.

Livestock browse Wyoming big sagebrush, but may use it only lightly when palatable herbaceous species are available.

Mountain big sagebrush is eaten by domestic livestock but has long been considered to be of low palatability, and a competitor to more desirable species.

Spiny hopsage provides a palatable and nutritious food source for livestock, particularly during late winter through spring. Domestic sheep browse the succulent new growth of spiny hopsage in late winter and early spring.

Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year.

Wildlife Interpretations:

Wyoming big sagebrush is preferred browse for wild ungulates. Pronghorn usually browse Wyoming big sagebrush heavily. Sagebrush-grassland communities provide critical sage-grouse breeding and nesting habitats. Meadows surrounded by sagebrush may be used as feeding and strutting grounds. Sagebrush is a crucial component of their diet year-round, and sage-grouse select sagebrush almost exclusively for cover. Sage-grouse prefer mountain big sagebrush and Wyoming big sagebrush communities to basin big sagebrush communities.

Mountain big sagebrush is highly preferred and nutritious winter forage for mule deer.

Spiny hopsage provides a palatable and nutritious food source for big game animals. Spiny hopsage is used as forage to at least some extent by domestic goats, deer, pronghorn, and rabbits.

Thurber needlegrass is valuable forage for wildlife.

Hydrological functions

Runoff is medium to very high. Permeability is moderately slow to moderate.

Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site is used for camping and hiking and has potential for upland and big game hunting.

Other products

Native Americans made tea from big sagebrush leaves. They used the tea as a tonic, an antiseptic, for treating colds, diarrhea, and sore eyes and as a rinse to ward off ticks. Big sagebrush seeds were eaten raw or made into meal.

Native Americans used big sagebrush leaves and branches for medicinal teas, and the leaves as a fumigant. Bark was woven into mats, bags and clothing.

Some Native American peoples traditionally ground parched seeds of spiny hopsage to make pinole flour.

Other information

Wyoming big sagebrush is used for stabilizing slopes and gullies and for restoring degraded wildlife habitat, rangelands, mine spoils and other disturbed sites. It is particularly recommended on dry upland sites where other shrubs are difficult to establish.

Spiny hopsage has moderate potential for erosion control and low to high potential for long-term revegetation projects. It can improve forage, control wind erosion, and increase soil stability on gentle to moderate slopes. Spiny hopsage is suitable for highway plantings on dry sites in Nevada.

Inventory data references

NASIS soil component data.

Type locality

Location 1: Pershing County, NV	
Township/Range/Section	T26N R26E S7
General legal description	Sahwave Mountains, Pershing County, Nevada.

Other references

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>).

Contributors

Approval

Kendra Moseley, 6/03/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/13/2025
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:
-
2. Presence of water flow patterns:
-
3. Number and height of erosional pedestals or terracettes:
-
4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
-
5. Number of gullies and erosion associated with gullies:
-
6. Extent of wind scoured, blowouts and/or depositional areas:
-
7. Amount of litter movement (describe size and distance expected to travel):
-
8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):
-

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-
17. **Perennial plant reproductive capability:**
-