

## Ecological site R028AY024NV SODIC TERRACE 5-8 P.Z.

Accessed: 05/11/2025

---

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	P. Novak-Echenique
Contact for lead author	State Rangeland Management Specialist
Date	10/28/2013
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

- 1. Number and extent of rills:** This site is nearly flat, thus rills are non-existent.  

---
- 2. Presence of water flow patterns:** Water flow patterns are common in areas subjected to summer convection storms. Flow patterns not extensive, short (<3m), meandering and stable.  

---
- 3. Number and height of erosional pedestals or terracettes:** Pedestals are none to rare with occurrence typically limited to areas within water flow patterns.  

---
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare Ground ± 60 to 80%  

---
- 5. Number of gullies and erosion associated with gullies:** None  

---
- 6. Extent of wind scoured, blowouts and/or depositional areas:** Typically none - wind scouring may occur during severe wind events prior to summer convection storms or winter storms.  

---

7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage from grasses and annual & perennial forbs) expected to move distance of slope length during intense summer convection storms or rapid snowmelt events. Persistent litter (large woody material) will remain in place except during large rainfall events.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values should be 3 to 6 on most soil textures found on this site. Areas of this site occurring on soils that have a physical crust will probably have stability values less than 3.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Surface structure is typically fine platy. Soil surface colors are light yellowish browns and soils are typified by an ochric epipedon. Surface textures are loams and silt loams. Organic matter of the surface 2 to 3 inches is less than 1 percent.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Shrub canopy and associated litter break raindrop impact and provide for snow capture on this site. Deep-rooted, perennial grasses (alkali sacaton) enhance infiltration and reduce runoff.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** Compacted layers are none. Platy or massive sub-surface horizons are not to be interpreted as compacted layers.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Reference State: Deep-rooted, warm season perennial bunchgrass (alkali sacaton)
- Sub-dominant: tall shrubs (black greasewood) > associated shrubs > deep-rooted, cool season perennial bunchgrasses > warm-season, rhizomatous grasses = perennial forbs.
- Other: Cool season rhizomatous grasses and annual forbs. Microbiotic crusts
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 35% of total woody canopy; mature bunchgrasses commonly ( $\pm 25\%$ ) have dead centers.
- 
14. **Average percent litter cover (%) and depth (in):** Between plant interspaces (10-20%) and depth ( $< \frac{1}{4}$  in.)
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** For normal or average growing season (thru June)  $\pm 500$  lbs/ac; Favorable years  $\pm 700$  lbs/ac and unfavorable years  $\pm 300$  lbs/ac.

- 
16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Potential invaders include halogeton, annual mustards, Russian thistle, and cheatgrass.
- 

17. **Perennial plant reproductive capability:** All functional groups should reproduce in average and above average growing season years. Reduced growth and reproduction occurs during extended or extreme drought periods.
-