

## **Ecological site R028BY073NV SHALLOW SILTY 5-8 P.Z.**

Last updated: 2/19/2025  
Accessed: 05/11/2025

---

### **General information**

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### **MLRA notes**

Major Land Resource Area (MLRA): 028B—Central Nevada Basin and Range

MLRA 28B occurs entirely in Nevada and comprises about 23,555 square miles (61,035 square kilometers). More than nine-tenths of this MLRA is federally owned. This area is in the Great Basin Section of the Basin and Range Province of the Intermontane Plateaus. It is an area of nearly level, aggraded desert basins and valleys between a series of mountain ranges trending north to south. The basins are bordered by long, gently sloping to strongly sloping alluvial fans. The mountains are uplifted fault blocks with steep sideslopes. Many of the valleys are closed basins containing sinks or playas. Elevation ranges from 4,900 to 6,550 feet (1,495 to 1,995 meters) in the valleys and basins and from 6,550 to 11,900 feet (1,995 to 3,630 meters) in the mountains.

The mountains in the southern half are dominated by andesite and basalt rocks that were formed in the Miocene and Oligocene. Paleozoic and older carbonate rocks are prominent in the mountains to the north. Scattered outcrops of older Tertiary intrusives and very young tuffaceous sediments are throughout this area. The valleys consist mostly of alluvial fill, but lake deposits are at the lowest elevations in the closed basins. The alluvial valley fill consists of cobbles, gravel, and coarse sand near the mountains in the apex of the alluvial fans. Sands, silts, and clays are on the distal ends of the fans.

The average annual precipitation ranges from 4 to 12 inches (100 to 305 millimeters) in most areas on the valley floors. Average annual precipitation in the mountains ranges from 8 to 36 inches (205 to 915 millimeters) depending on elevation. The driest period is from midsummer to midautumn. The average annual temperature is 34 to 52 degrees F (1 to 11 degrees C). The freeze-free period averages 125 days and ranges from 80 to 170 days, decreasing in length with elevation.

The dominant soil orders in this MLRA are Aridisols, Entisols, and Mollisols. The soils in the area dominantly have a mesic soil temperature regime, an aridic or xeric soil moisture regime, and mixed or carbonatic mineralogy. They generally are well drained, loamy or loamyskeletal, and shallow to very deep.

Nevada's climate is predominantly arid, with large daily ranges of temperature, infrequent severe storms and heavy snowfall in the higher mountains. Three basic geographical factors largely influence Nevada's climate: continentality, latitude, and elevation. The strong continental effect is expressed in the form of both dryness and large temperature variations. Nevada lies on the eastern, lee side of the Sierra Nevada Range, a massive mountain barrier that markedly influences the climate of the State. The prevailing winds are from the west, and as the warm moist air from the Pacific Ocean ascend the western slopes of the Sierra Range, the air cools, condensation occurs and most of the moisture falls as precipitation. As the air descends the eastern slope, it is warmed by compression, and very little precipitation occurs. The effects of this mountain barrier are felt not only in the West but throughout the state, as a result the lowlands of Nevada are largely desert or steppes.

The temperature regime is also affected by the blocking of the inland-moving maritime air. Nevada sheltered from maritime winds, has a continental climate with well-developed seasons and the terrain responds quickly to changes in solar heating. Nevada lies within the midlatitude belt of prevailing westerly winds which occur most of the year. These winds bring frequent changes in weather during the late fall, winter and spring months, when most of the precipitation occurs.

To the south of the mid-latitude westerlies, lies a zone of high pressure in subtropical latitudes, with a center over the Pacific Ocean. In the summer, this high-pressure belt shifts northward over the latitudes of Nevada, blocking storms from the ocean. The resulting weather is mostly clear and dry during the summer and early fall, with

occasional thundershowers. The eastern portion of the state receives noteworthy summer thunderstorms generated from monsoonal moisture pushed up from the Gulf of California, known as the North American monsoon. The monsoon system peaks in August and by October the monsoon high over the Western U.S. begins to weaken and the precipitation retreats southward towards the tropics (NOAA 2004).

## Ecological site concept

This site occurs on fan skirts, lake plains and alluvial flats. Slopes range from 0 to 8 percent, but slope gradients of 2 to 4 percent are most typical. Elevations are 4800 to 6700 feet.

The climate associated with this site is semiarid, characterized by cool, moist winters and warm, dry summers. Average annual precipitation is 5 to 8 inches. Mean annual air temperature is 45 to 50 degrees F. The average growing season is about 100 to 120 days.

Soils associated with this site are very deep and well drained. They typically have a restrictive layer at less than 10 inches from the surface. Permeability is slow to moderate and runoff is low to high. Surface textures are usually silts and silt loams. The soils are moderately to very strongly alkaline and calcareous throughout. The potential for sheet and rill erosion is slight.

The reference state is dominated by shadscale. Production ranges from 200 to 400 pounds per acre.

## Associated sites

R028BY017NV	<b>LOAMY 5-8 P.Z.</b> This site occurs on fan skirts. Slopes gradients of 2 to 8 percent are typical. Elevations range from 5000 to 6500 feet. The soils associated with this site are very deep, well drained, and formed in alluvium derived from mixed rocks. Soils are alkaline throughout and characterized by an ochric epipedon. Soil surface structure is typically platy with vesicular pores. The soil moisture regime is typic aridic and the soil temperature regime is mesic. The reference state is dominated by Indian ricegrass, bottlebrush squirreltail, and shadscale. Production ranges from 200 to 400 pounds per acre.
R028BY018NV	<b>SILTY 5-8 P.Z.</b> This site occurs on lakeplains. Slopes gradients are typically less than 4 percent and elevations range from 4500 to 5500 feet. The soils associated with this site are very deep, well drained and formed in mixed alluvium. Soils are characterized by an ochric epipedon, a calcic horizon, and a vesicular surface horizon. Soil temperature regime is typic aridic and the soil moisture regime is mesic. The plant community is dominated by winterfat. Indian ricegrass and bottlebrush squirreltail are other important species. Production ranges from 200 to 500 pounds per acre.
R028BY074NV	<b>SODIC TERRACE 5-8 P.Z.</b> This site occurs on fan skirts, alluvial flats and lakeplains. Slopes range from 0 to 15 percent, but slope gradients of 0 to 8 percent are most typical. Elevations are 4300 to 5600 feet. The soils associated with this site are very deep, moderately well to well drained, and formed over lacustrine sediments. Depth to the lake sediments usually ranges from 30 to 40 inches. Soils are calcareous throughout and moderately to strongly saline. The reference state is dominated by shadscale and black greasewood. Although shadscale makes up most of the annual production, black greasewood is often prevalent enough to dominate the visual aspect. Production ranges from 200 to 600 pounds per acre.

## Similar sites

R028BY074NV	<b>SODIC TERRACE 5-8 P.Z.</b> SAVE4 is more prevalent
R028BY075NV	<b>COARSE GRAVELLY LOAM 6-8 P.Z.</b> KRLA2 and ACHY are more prevalent
R028BY017NV	<b>LOAMY 5-8 P.Z.</b> PIDE4 and ACHY are more prevalent
R028BY007NV	<b>LOAMY 10-12 P.Z.</b> more productive site; typically occurs on inset fans of piedmont slope landscapes

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Atriplex confertifolia</i>

Herbaceous	(1) <i>Elymus elymoides</i> (2) <i>Achnatherum hymenoides</i>
------------	--

Physiographic features

This site occurs on fan skirts, lake plains and alluvial flats. Slopes range from 0 to 8 percent, but slope gradients of 2 to 4 percent are most typical. Elevations are 4800 to 6700 feet.

Table 2. Representative physiographic features

Landforms	(1) Fan skirt (2) Lake plain (3) Alluvial flat
Runoff class	Low to high
Flooding duration	Extremely brief (0.1 to 4 hours)
Flooding frequency	None to rare
Ponding frequency	None
Elevation	4,800–6,700 ft
Slope	0–8%
Water table depth	72 in
Aspect	Aspect is not a significant factor

Climatic features

The climate associated with this site is semiarid, characterized by cold, moist winters and warm, dry summers. Average annual precipitation is about 5-8 inches. Mean annual air temperature ranges from 45 to 50 degrees F. The average number of growing season day is about 100 to 120 days. Average monthly precipitation across the range in which this ecological site occurs is:  
\*The above data is averaged from Lages and Beowawe WRCC stations.

Table 3. Representative climatic features

Frost-free period (average)	93 days
Freeze-free period (average)	116 days
Precipitation total (average)	8 in






Figure 1. Monthly precipitation range



**Figure 2. Monthly average minimum and maximum temperature**



**Figure 3. Annual precipitation pattern**

## Climate stations used

- (1) BEOWAVE 49S U OF N RCH [USC00260800], Eureka, NV
- (2) LAGES [USC00264341], Ely, NV

## Influencing water features

There are no influencing water features associated with this site.

## Soil features

Soils associated with this site are very deep and well drained. They typically have a restrictive layer at less than 10 inches from the surface. Permeability is slow to moderate and runoff is low to high. Surface textures are usually silts and silt loams. The soils are moderately to very strongly alkaline and calcareous throughout. The potential for sheet and rill erosion is slight. The soil moisture regime is typic aridic. The soil series associated with this site include: Hardhat, Katelana, Mysol, Mazuma, Sondoia, and Yelbrick.

A representative soil component is Katelana, level (NV766, MU917), classified as a fine-silty, carbonatic, mesic Typic Torriorthents. An ochric epipedon occurs from the soil surface to 7 inches. Clay content in the particle control section averages 18 to 27 percent, when mixed. Reaction is strongly alkaline. Effervescence is violently effervescent. Lithology consists of limestone over lacustrine sediments.

**Table 4. Representative soil features**

Parent material	(1) Alluvium–limestone
Surface texture	(1) Silt loam (2) Silty clay loam (3) Ashy silt loam (4) Silt
Family particle size	(1) Loamy
Drainage class	Well drained

Permeability class	Slow to moderate
Soil depth	55–72 in
Surface fragment cover ≤3"	0–7%
Surface fragment cover >3"	1–3%
Available water capacity (0–40in)	3.3–7.9 in
Calcium carbonate equivalent (0–40in)	10–20%
Electrical conductivity (0–40in)	4–32 mmhos/cm
Sodium adsorption ratio (0–40in)	13–30
Soil reaction (1:1 water) (0–40in)	8.4–9
Subsurface fragment volume ≤3" (Depth not specified)	3–20%
Subsurface fragment volume >3" (Depth not specified)	1–3%

## Ecological dynamics

An ecological site is the product of all the environmental factors responsible for its development and it has a set of key characteristics that influence a site's resilience to disturbance and resistance to invasives. Key characteristics include 1) climate (precipitation, temperature), 2) topography (aspect, slope, elevation, and landform), 3) hydrology (infiltration, runoff), 4) soils (depth, texture, structure, organic matter), 5) plant communities (functional groups, productivity), and 6) natural disturbance regime (fire, herbivory, etc.) (Caudle 2013). Biotic factors that influence resilience include site productivity, species composition and structure, and population regulation and regeneration (Chambers et al. 2013).

The ecological site is dominated by deep-rooted cool season, perennial bunchgrasses and drought tolerant shrubs with high root to shoot ratios. The dominant shrubs usually root to the full depth of the winter-spring soil moisture recharge, which Fernandez and Caldwell (1975) reported as between 80 and 110 cm for shadscale and winterfat. Shadscale initiates root growth in early April, a few days to a week prior to aerial plant parts and shadscale in particular exhibits active root growth for several weeks after termination of shoot growth (Fernandez and Caldwell 1975). Continued root growth, even for established plants that are not exploring new areas of the soil, facilitates water absorption particularly in low soil moisture conditions (Gardner 1960). Fernandez and Caldwell (1975) concluded that the ability of shadscale to explore the soil volume at greater depths with a more profuse system of small branching lateral roots than winterfat or sagebrush may play a role in its ability to remain photosynthetically active longer into the summer season. Although shadscale exhibits the ability to withstand drought conditions on a short-term basis the forty year photographic record (1951-1990) from the Raft River Valley of south-central Idaho visually demonstrates the impact of multiple years of drought on shadscale communities (Sharp et al. 1990). Scale insects have also been implicated in the death of shadscale (Sharp et al. 1990) however the data on this subject remains inconclusive (Nelson et al. 1990). Interestingly, periods of above normal springtime precipitation are also linked to shadscale die-off. Nelson et al. (1990) investigated areas of severe shadscale die-off that were, for the most part, located in low areas in valley bottoms or upland depressions that apparently incurred prolonged high soil moisture during a wet period. The high soil moisture appeared to be correlated with increased pythiaceous fungi leading to rootlet mortality and plant stress (Nelson et al. 1990). The authors suggest that depending on the degree and duration of plant stress, injury could range from a sustained disease to rapid death.

Shadscale is a densely clumped, rounded, compact native shrub. It generally attains heights of 8 to 32 inches and widths of 12 to 68 inches (Blaisdell and Holmgren 1984). Shadscale is considered an evergreen to partially deciduous shrub, since a small percentage of leaves are dropped in the winter (Smith and Nobel 1986). Shadscale possesses wider ecological amplitude than most *Atriplex* species (Crofts and Van Epps 1975), and shows ploidy levels from diploid (2x) to decaploid (10x). The extensive polyploidy of shadscale is an important consideration when implementing revegetation projects because ploidy levels are usually associated with distinct habitats

(Sanderson et al. 1990). Diploid individuals are unlikely to perform as well in areas where tetraploids are more common. Diploid individuals generally occur above Pleistocene lake levels, whereas lake floors are usually occupied by autotetraploids. Overall, tetraploids are the most widespread throughout its range (Carlson 1984). Bud sagebrush, a common shrub to this ecological site, is a native, summer-deciduous shrub. It is low growing, spinescent, aromatic shrub with a height of 4 to 10 inches and a spread of 8 to 12 inches (Chambers and Norton 1993).

The perennial bunchgrasses that are sub-dominant with the shrubs include Indian ricegrass and bottlebrush squirreltail. The dominant grass within this site, is Indian ricegrass a hardy, cool-season, densely tufted, native perennial bunchgrass that grows from 4 to 24 inches in height (Blaisdell and Holmgren 1984). Squirreltail is a competitive, short-lived, perennial grass that readily establishes from seed. These species generally have somewhat shallower root systems than the shrubs, but root densities are often as high or higher than those of the shrubs in the upper 0.5m of the soil profile. General differences in root depth distributions between grasses and shrubs results in resource partitioning these shrub – grass systems.

The invasibility of plant communities is often linked to resource availability. Disturbance can decrease resource uptake due to damage or mortality of the native species and depressed competition or can increase resource pools by the decomposition of dead plant material following disturbance. Historically, shadscale dominant salt-desert shrub communities were free of exotic invaders; however, excessive grazing pressure during settlement and into the 20th century has increased the overall presence of cheatgrass, halogeton, Russian thistle and annual non-native mustard species (Peters and Bunting 1994). The presence of exotic annual plants within these ecosystems decreases ecosystem resilience and resistance to disturbance through competition for limited resources. Dobrowolski et al. (1990) cite multiple authors on the extent of the soil profile exploited by the competitive exotic annual cheatgrass. Specifically, the depth of rooting is dependent on the size the plant achieves and in competitive environments cheatgrass roots were found to penetrate only 15 cm whereas isolated plants and pure stands were found to root at least 1 m in depth with some plants rooting as deep as 1.5 to 1.7 m.

The ecological site has low resilience to disturbance and resistance to invasion. Increased resilience increases with elevation, aspect, increased precipitation and increased nutrient availability. Four possible stable states have been identified for this site.

#### Fire Ecology:

The lack of continuous fuels to carry fires made fire rare to nonexistent in shadscale communities (Young and Tipton 1990), thus it is not surprising that shadscale and bud sagebrush are both fire intolerant (Banner 1992, West 1994). Shadscale does not readily recover from fire, except for establishment through seed (West 1994). The slow reestablishment allows for easy invasion by cheatgrass and other non-native weedy species (Sanderson et al. 1990). The increased presence of exotic annual grasses has greatly altered fire regimes in areas of the Intermountain West where shadscale is a major vegetational component. Exotic annuals increase fire frequency under wet to near-normal summer moisture conditions and repeated, frequent fire has converted large expanses of shadscale rangeland to annual non-native plant communities (Knapp 1998).

The effect of fire on bunchgrasses relates to culm density, culm-leaf morphology, and the size of the plant. The initial condition of bunchgrasses within the site along with seasonality and intensity of the fire factor into individual species' responses. For most forbs and grasses the growing points are located at or below the soil surface providing relative protection from disturbances which decrease above ground biomass, such as grazing or fire. Thus, fire mortality is more correlated to duration and intensity of heat which is related to culm density, culm-leaf morphology, size of plant and abundance of old growth (Wright 1971, Young 1983). However, season and severity of the fire and post-fire soil moisture availability will influence plant response.

Indian ricegrass is a deep-rooted, cool season perennial bunchgrass that is adapted primarily to sandy soils. A prominent grass on this site, it is fairly fire tolerant (Wright 1985), which is likely due to its low culm density and below ground plant crowns. Vallentine (1989) cites several studies in the sagebrush zone that classified Indian ricegrass as being slightly damaged from late summer burning. Indian ricegrass has also been found to reestablish on burned sites through seed dispersed from adjacent unburned areas (Young 1983, West 1994). Thus the presence of surviving, seed producing plants facilitates the reestablishment of Indian ricegrass. Grazing management following fire to promote seed production and establishment of seedlings is important.

Bottlebrush squirreltail is considered one of the most fire resistant bunchgrasses due to its small size, coarse stems, and sparse leafy material (Britton et al. 1990). Postfire regeneration occurs from surviving root crowns and from on- and off-site seed sources. Bottlebrush squirreltail has the ability to produce large numbers of highly germinable seeds, with relatively rapid germination (Young and Evans 1977) when exposed to the correct

environmental cues. It exhibits the ability to germinate in the late fall and very early spring at a wide range of temperatures making it a strong competitor with cheatgrass (USDA NRCS Plant Fact Sheet). Early spring growth and ability to grow at low temperatures contribute to the persistence of bottlebrush squirreltail among cheatgrass dominated ranges (Hironaka and Tisdale 1973).

Needle and thread a minor component on this site is a fine leaf grass and is considered sensitive to fire (Akinsoji 1988, Bradley et al. 1992, Miller et al. 2013). In a study by Wright and Klemmedson (1965), season of burn rather than fire intensity seemed to be the crucial factor in mortality for needle and thread grass. Early spring season burning was seen to kill the plants while August burning had no effect. Thus, under wildfire scenarios needle and thread is often present in the post-burn community.

Sandberg's bluegrass also a minor component on this ecological site, has been found to increase following fire likely due to its low stature and productivity (Daubenmire 1975). Sandberg bluegrass may retard reestablishment of deeper rooted bunchgrass.

Rehabilitation following fire will have limited success. Observations from one hundred and seven separate plantings within the shadscale zone in Utah and Nevada indicate a very low success rate (Bleak et al. 1965). Seed from 148 native and non-native grasses, forbs and shrubs were planted from 1937 to 1962 across ten locations. Good seedling stands were obtained with introduced wheatgrasses, but most perished during the first summer. A few plantings of crested, fairway and Siberian wheatgrass along with Russian wildrye maintained stands for 10 or more years but eventually declined to a very few plants (Bleak et al. 1965). The primary cause of seeding failures appeared to be the arid climate.

Livestock/wildlife grazing interpretations:

Traditionally, shadscale plant communities provided good winter forage for the expanding sheep and cattle industry in the arid west. Shadscale is a valuable browse species for a wide variety of wildlife and livestock (Blaisdell and Holmgren 1984). The spinescent growth habit of shadscale lends to its browsing tolerance with no more than 15 to 20% utilization by sheep being reported (Blaisdell and Holmgren 1984) and significantly less utilization by cattle. Increased presence of shadscale within grazed versus ungrazed areas is generally a result of the decreased competition from more heavily browsed associates (Cibils et al. 1998). Reduced competition from more palatable species in heavily grazed areas may increase shadscale germination and establishment. Chambers and Norton (1993) found shadscale establishment higher under spring than winter browsing as well as heavy compared to light browsing. During years of below average precipitation, shadscale has been found very susceptible to grazing pressure regardless of season (Chambers and Norton 1993). Following fire, grazing exclusion for 2 or more years is beneficial for revegetation of shadscale communities as first year shadscale seedlings lack spines and are highly susceptible to browsing. Spines develop in the second year (Zielinski 1994).

Bud sagebrush is also a palatable, nutritious forage for upland game birds, small game, big game and domestic sheep in winter, particularly late winter (Johnson 1978), however it can be poisonous or fatal to calves when eaten in quantity (Stubbendieck et al. 1992). Bud sagebrush is highly susceptible to effects of browsing. It decreases under browsing due to year-long palatability of its buds and is particularly susceptible to browsing in the spring when it is physiologically most active (Chambers and Norton 1993, Harper et al. 1990). Heavy browsing (>50%) may kill bud sagebrush rapidly (Wood and Brotherson 1986). Winterfat, a highly nutritious winter feed shows similar results to bud sagebrush with significant declines in density with late winter or early spring grazing (Harper et al. 1990). Interestingly the same 54 year study also showed winterfat density decreasing in the ungrazed plots.

Indian ricegrass is a preferred forage species for livestock and wildlife (Cook 1962, Booth et al. 2006). This species is often heavily utilized in winter because it cures well (Booth et al. 2006). It is also readily utilized in early spring, being a source of green feed before most other perennial grasses have produced new growth (Quinones 1981). Booth et al. (2006) note that the plant does well when utilized in winter and spring. Cook and Child (1971) however, found that repeated heavy grazing reduced crown cover, which may reduce seed production, density, and basal area of these plants. Additionally, heavy early spring grazing reduces plant vigor and stand density (Stubbendieck 1985). In eastern Idaho, productivity of Indian ricegrass was at least 10 times greater in undisturbed plots than in heavily grazed ones (Pearson 1965). Cook and Child (1971) found significant reduction in plant cover even after 7 years of rest from heavy (90%) and moderate (60%) spring use. The seed crop may be reduced where grazing is heavy (Bich et al. 1995). Tolerance to grazing increases after May, thus spring deferment may be necessary for stand enhancement (Pearson 1964, Cook and Child 1971); however, utilization of less than 60% is recommended. In summary, adaptive management is required to manage this bunchgrass well.

Bottlebrush squirreltail generally increases in abundance when moderately grazed or protected (Hutchings and Stewart 1953). It is considered fair to good forage for cattle, horses and sheep in the spring prior to seed development and in the late fall after seed shatter. In addition, moderate trampling by livestock in big sagebrush

rangelands of central Nevada enhanced bottlebrush squirreltail seedling emergence compared to untrampled conditions. Heavy trampling however was found to significantly reduce germination sites (Eckert et al. 1987). Squirreltail is more tolerant of grazing than Indian ricegrass but all bunchgrasses are sensitive to over utilization within the growing season.

Needle and thread a minor component on these sites, is most commonly found on warm/dry soils (Miller et al. 2013). It is not grazing tolerant and will be one of the first grasses to decrease under heavy grazing pressure (Smoliak et al. 1972, Tueller and Blackburn 1974). Heavy grazing is likely to reduce basal area of these plants (Smoliak et al. 1972).

In summary, overgrazing causes a decrease in Indian ricegrass along with bud sagebrush, while shadscale may initially increase. Spring grazing year after year can be detrimental to bud sagebrush and bunchgrasses. Continued abusive grazing leads to increased bare ground and invasion by annual weeds (e.g., cheatgrass, halogeton, and tansy mustard). Shadscale may become dominant with an annual understory. With further deterioration, shadscale declines, bare ground increases, soil redistribution accelerates and site productivity decreases. On some soils, erosion can result in increased surface salts and development of desert pavement. Reestablishment of perennials is limited in areas of extensive desert pavement. Fire is a very infrequent and patchy event in these salt desert shrub communities; however, where it has occurred the shrub community is greatly reduced and annual exotic weeds will increase if present. Repeated fire within a 10 to 20 year timeframe has the potential to convert this site to an annual weed dominated system. Knowledge of successful rehabilitation strategies in these droughty plant communities is limited.

## State and transition model




Figure 5. State and Transition Model



Reference State 1.0

Transition T1A: Introduction of non-native annual species such as halogeton.

Current Potential State 2.0 Community Phase Pathways

2.1a: Long-term drought and/or inappropriate grazing management

2.2a: Release from drought and/or lack of disturbance allows for an increase in perennial grasses. Extreme growing season moisture may reduce shadscale.

Transition T2A: Inappropriate grazing management and/or long-term drought.

Transition T2B: Soil disturbing treatments (drill seeding, roller chopper, Lawson aerator etc.), fire, and/or unusually wet spring.

Transition T3A: Soil disturbing treatments (drill seeding, roller chopper, Lawson aerator etc.), fire, and/or unusually wet spring.

Figure 6. Legend

## Animal community

### Livestock Interpretations:

This site is suitable for livestock grazing. Grazing management considerations include timing, duration, frequency, and intensity of grazing.

Traditionally, shadscale plant communities provided good winter forage for the expanding sheep and cattle industry in the arid West. Indian ricegrass is the dominant grass on this site and is a preferred forage species for livestock and wildlife (Cook 1962, Booth et al. 2006). Indian ricegrass is highly palatable to all classes of livestock in both green and cured condition. It supplies a source of green forage before most other native grasses have produced much new growth. This species is often heavily utilized in winter because it cures well (Booth et al. 2006). It is also readily utilized in early spring, being a source of green feed before most other perennial grasses have produced new growth (Quinones 1981). Booth et al. (2006) note that the plant does well when utilized in winter and spring. Cook and Child (1971) however, found that repeated heavy grazing reduced crown cover, which may reduce seed production, density, and basal area of these plants. Additionally, heavy early spring grazing reduces plant vigor and stand density (Stubbendieck 1985). In eastern Idaho, productivity of Indian ricegrass was at least 10 times greater in undisturbed plots than in heavily grazed ones (Pearson 1965). Cook and Child (1971) found significant reduction in plant cover even after 7 years of rest from heavy (90%) and moderate (60%) spring use. The seed crop may be reduced where grazing is heavy (Bich et al. 1995). Tolerance to grazing increases after May, thus spring deferment may be necessary for stand enhancement (Pearson 1964, Cook and Child 1971); however, utilization of less than 60% is recommended. In summary, adaptive management is required to manage this bunchgrass well.

Shadscale is a valuable browse species for a wide variety of wildlife and livestock (Blaisdell and Holmgren 1984). Shadscale provides good browse for domestic sheep. Shadscale leaves and seeds are an important component of domestic sheep and cattle winter diets. The spinescent growth habit of shadscale lends to its browsing tolerance with no more than 15 to 20% utilization by sheep being reported (Blaisdell and Holmgren 1984) and significantly less utilization by cattle. Increased presence of shadscale within grazed versus ungrazed areas is generally a result of the decreased competition from more heavily browsed associates (Cibils et al. 1998). Reduced competition from more palatable species in heavily grazed areas may increase shadscale germination and establishment. Chambers and Norton (1993) found shadscale establishment higher under spring than winter browsing as well as heavy compared to light browsing. During years of below average precipitation, shadscale has been found very susceptible to grazing pressure regardless of season (Chambers and Norton 1993). Following fire, grazing exclusion for 2 or more years is beneficial for revegetation of shadscale communities as first year shadscale seedlings lack spines and are highly susceptible to browsing. Spines develop in the second year (Zielinski 1994). Greenmolly provides excellent forage for sheep and cattle, and is often used as a winter forage, when it is high in

protein.

Needle and thread also a minor component on these sites, is most commonly found on warm/dry soils (Miller et al. 2013). It is not grazing tolerant and will be one of the first grasses to decrease under heavy grazing pressure (Smoliak et al. 1972, Tueller and Blackburn 1974). Heavy grazing is likely to reduce basal area of these plants (Smoliak et al. 1972). With the reduction in competition from deep rooted perennial bunchgrasses, the rhizomatous galleta grass bluegrass will likely increase (Smoliak et al. 1972).

In summary, overgrazing causes a decrease in Indian ricegrass along with winterfat and bud sagebrush, while shadscale may initially increase. Spring grazing year after year can be detrimental to bud sagebrush and the perennial bunchgrasses. Continued abusive grazing leads to increased bare ground and invasion by annual weeds (e.g., cheatgrass, halogeton, and tansy mustard). Shadscale may become dominant with an annual understory. With further deterioration, shadscale declines, bare ground increases, soil redistribution accelerates and site productivity decreases. On some soils, erosion can result in increased surface salts and development of desert pavement. Reestablishment of perennials is limited in areas of extensive desert pavement. Fire is a very infrequent and patchy event in these salt-desert shrub communities; however, where it has occurred the shrub community is greatly reduced and annual exotic weeds will increase if present. Repeated fire within a 10 to 20 year timeframe has the potential to convert this site to an annual weed dominated system. Knowledge of successful rehabilitation strategies in these droughty plant communities is limited. Cook and Child (1971) found significant reduction in plant cover even after 7 years of rest from heavy (90%) and moderate (60%) spring use. The seed crop may be reduced where grazing is heavy (Bich et al. 1995). Tolerance to grazing increases after May, thus spring deferment may be necessary for stand enhancement (Pearson 1964, Cook and Child 1971); however, utilization of less than 60% is recommended. Adaptive management is required to manage this bunchgrass well.

Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine-tuned by the client by adaptive management through the year and from year to year.

#### Wildlife interpretations:

Salt-desert shrub communities provide valuable habitat for a number of species.

Shadscale is a valuable browse species, providing a source of palatable, nutritious forage for a wide variety of wildlife particularly during spring and summer before the hardening of spiny twigs. (Jameson 1952, Welch et al. 1987). It supplies browse, seed, and cover for birds, small mammals, rabbits, deer, and pronghorn antelope. Shadscale also provides feed for wild ungulates: mule deer (*Odocoileus hemionus*) browse shadscale, especially during winter (Bartmann 1983). Although it is not preferred, shadscale is also browsed in winter by pronghorn (*Antilocapra americana*) (Beal and Smith 1970). Shadscale habitats throughout northeastern Nevada are important home ranges for small mammals. The chisel-toothed kangaroo rat (*Dipodomys microps*) feed on shadscale foliage and use shadscale habitats during the spring, summer, and fall. Deer mice (*Peromyscus maniculatus*) use shadscale habitats all year (O'farrell and Clark 1986). Shadscale leaves and seeds are preferred forage for jackrabbits (*Lepus californicus*) (Currie and Goodwin 1966). The Great Basin kangaroo rat (*Dipodomys ordii*) also feeds on shadscale foliage (Kenagy 1973).

Several bird species will eat shadscale seeds and use shadscale habitats for cover and nesting sites. The horned lark (*Eremophila alpestris*) occurs throughout shadscale communities. Although less commonly apparent the Brewer's sparrow (*Spizella breweri*) and sage thrasher (*Oreoscoptes montanus*) also occur in shadscale habitat. Other species, observed occasionally throughout breeding season in shadscale habitat include: northern harrier (*Circus cyaneus*), red-tailed hawk (*Buteo jamaicensis*), ferruginous hawk (*Buteo regalis*), golden eagle (*Aquila chrysaetos*), American kestrel (*Falco sparverius*), prairie falcon (*Falco mexicanus*), mourning dove (*Zenaidura macroura*), burrowing owl (*Athene cunicularia*), short-eared owl (*Asio flammeus*), violet-green swallow (*Tachycineta thalassina*), cliff swallow (*Petrochelidon*), barn swallow (*Hirundo rustica*), common raven (*Corvus corax*), loggerhead shrike (*Lanius ludovicianus*), vesper sparrow (*Pooecetes gramineus*), black-throated sparrow (*Amphispiza bilineata*), and western meadowlark (*Sturnella neglecta*) (Medin 1990).

It should be noted the loss of shadscale and associated shrubs has a negative effect on golden eagle habitat. The golden eagle is listed as a threatened species throughout the United States. Areas of shadscale shrub-steppe provide cover and forage for black-tailed jackrabbits, which are a major food source of golden eagles. Shadscale should be maintained within 1.9 miles of golden eagle nests in order to maintain the species (Kochert et al. 1999).

## Hydrological functions

Runoff is low to high. Permeability is slow to moderate. These sites are subject to brief ponding and flooding after summer convection storms. As a result, water flow patterns may be numerous in areas. Flow patterns may be fairly

long, meandering and stable. Pedestals are rare with occurrence typically limited to areas within water flow paths. Terracettes are non-existent. Sparse shrub canopy and associated litter provide some protection from raindrop impact and allow for some snow capture on this site. Medium to fine textured surface soils have moderate to slow infiltration and medium runoff. The vesicular crust also impedes infiltration

## Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site is used for camping and hiking and has potential for upland and big game hunting.

## Other products

Seeds of shadscale were used by Native Americans for bread and mush. Indian ricegrass was traditionally eaten by some Native American peoples. The Paiutes used the seeds as a reserve food source.

## Other information

Bottlebrush squirreltail is tolerant of disturbance and is well-suited for revegetation. Indian ricegrass is well-suited for surface erosion control and desert revegetation although it is not highly effective in controlling sand movement.

## Inventory data references

NASIS soil component data.

## Type locality

Location 1: Elko County, NV	
Township/Range/Section	T37N R69E S10
Latitude	41° 6' 20"
Longitude	114° 8' 50"
General legal description	NE ¼ NE ¼, Approximately 12 miles south of Montello, Pilot Creek Valley area, Elko County, Nevada. This site also occurs in Eureka and White Pine Counties, Nevada.

## Other references

Akinsoji, A. 1988. Postfire vegetation dynamics in a sagebrush steppe in southeastern Idaho, USA. *Vegetatio* 78:151-155.

Banner, R. E. 1992. Vegetation Types of Utah. *Rangelands* 14:109-114.

Bich, B. S., J. L. Butler, and C. A. Schmidt. 1995. Effects of Differential Livestock Use on Key Plant Species and Rodent Populations within Selected *Oryzopsis hymenoides*/*Hilaria jamesii* Communities of Glen Canyon National Recreation Area. *The Southwestern Naturalist* 40:281-287.

Blaisdell, J. P. and R. C. Holmgren. 1984. Managing Intermountain rangelands - salt-desert shrub ranges. General Technical Report INT-163, USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, UT.

Booth, D. T., C. G. Howard, and C. E. Mowry. 2006. 'Nezpar' Indian ricegrass: description, justification for release, and recommendations for use. *Rangelands Archives* 2:53-54.

Bradley, A. F., N. V. Noste, and W. C. Fischer. 1992. Gen. Tech. Rep. INT-287: Fire ecology of forests and woodlands in Utah. . U.S. Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, UT.

Britton, C. M., G. R. McPherson, and F. A. Sneva. 1990. Effects of burning and clipping on five bunchgrasses in eastern Oregon. *Western North American Naturalist* 50:115-120.

- Carlson, J. 1984. Atriplex cultivar development. Pages 176-182 in Proceedings - symposium on the biology of Atriplex and related chenopods Gen. Tech. Rep. INT-172. USDA, Forest Service, Intermountain Forest and Range Experiment Station, Provo, UT.
- Chambers, J., B. Bradley, C. Brown, C. D'Antonio, M. Germino, J. Grace, S. Hardegree, R. Miller, and D. Pyke. 2013. Resilience to Stress and Disturbance, and Resistance to *Bromus tectorum* L. Invasion in Cold Desert Shrublands of Western North America. *Ecosystems*:1-16.
- Chambers, J. C. and B. E. Norton. 1993. Effects of grazing and drought on population dynamics of salt desert species on the Desert Experimental Range, Utah. *Journal of Arid Environments*:261-275.
- Cibils, A. F., S. M. David, and D. E. McArthur. 1998. Plant-Herbivore Interactions in Atriplex: Current State of Knowledge. General Technical Report RMRS-GTR-14, USDA: FS, Rocky Mountain Research Station, Ogden, UT.
- Cook, C. W. 1962. An Evaluation of Some Common Factors Affecting Utilization of Desert Range Species. *Journal of Range Management* 15:333-338.
- Cook, C. W. and R. D. Child. 1971. Recovery of Desert Plants in Various States of Vigor. *Journal of Range Management* 24:339-343.
- Crofts, K. and G. Van Epps. 1975. Use of shadscale in revegetation of arid disturbed sites. Pages 151-152 in *Wildland shrubs: symposium and workshop proceedings*. Brigham Young University, Provo, UT.
- Daubenmire, R. 1975. Plant succession on abandoned fields, and fire influences in a steppe area in southeastern Washington. *Northwest Science* 49:36-48.
- Dobrowolski, J. P., M. M. Caldwell, and J. H. Richards. 1990. Basin hydrology and plant root systems. Pages 243-292 *Plant Biology of the Basin and Range*. Springer.
- Eckert, R. E., Jr., F. F. Peterson, and F. L. Emmerich. 1987. A study of factors influencing secondary succession in the sagebrush [*Artemisia* spp. L.] type. Pages 149-168 in *Proceedings: Seed and seedbed ecology of rangeland plants*. U. S. Department of Agriculture, Agricultural Research Service, Tucson, A.Z.
- Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).
- Harper, K.T., F.J. Wagstaff and W.P. Clary. 1990. Shrub mortality over a 54-year period in shadscale desert, west-central Utah. Pages 119-126 in *Proceedings-Symposium on Cheatgrass Invasion, Shrub Die-off, and Other Aspects of Shrub Biology and Management*. Gen. Tech. Rep. INT-GTR-276. USDA, Forest Service, Intermountain Research Station, Ogden, UT.
- Hironaka, M. and E. Tisdale. 1973. Growth and development of *Sitanion hystrix* and *Poa sandbergii*. Research Memorandum RM 72-24. U.S. International Biological Program, Desert Biome.
- Houghton, J.G., C.M. Sakamoto, and R.O. Gifford. 1975. Nevada's Weather and Climate, Special Publication 2. Nevada Bureau of Mines and Geology, Mackay School of Mines, University of Nevada, Reno, NV.
- Hutchings, S. S. and G. Stewart. 1953. Increasing forage yields and sheep production on intermountain winter ranges. Circular No. 925. U.S. Department of Agriculture, Washington, D.C.
- Johnson, K. L. 1978. Wyoming shrublands: Proceedings, 7th Wyoming shrub ecology workshop. Page 58. University of Wyoming, Agricultural Extension Service, Rock Spring WY.
- Knapp, P. A. 1998. Spatio-temporal patterns of large grassland fires in the Intermountain West, U.S.A. *Global Ecology & Biogeography Letters* 7:259-272.
- Miller, R. F., J. C. Chambers, D. A. Pyke, F. B. Pierson, and C. J. Williams. 2013. A review of fire effects on vegetation and soils in the Great Basin Region: response and ecological site characteristics.

National Oceanic and Atmospheric Administration. 2004. The North American Monsoon. Reports to the Nation. National Weather Service, Climate Prediction Center. Available online: <http://www.weather.gov/>.

Nelson, D.L., D.J. Weber and S.C. Garvin. 1990. The possible role of plant disease in the recent wildland shrub dieoff in Utah. Pages 84-90 in Proceedings-Symposium on Cheatgrass Invasion, Shrub Die-off, and Other Aspects of Shrub Biology and Management. Gen. Tech. Rep. INT-GTR-276. USDA, Forest Service, Intermountain Research Station, Ogden, UT.

Nelson, C.R., B.A. Haws and D.L. Nelson. 1990. Mealybugs and related homoptera of shadscale: Possible agents in the dieoff problem in the intermountain west. Pages 152-165 in Proceedings-Symposium on Cheatgrass Invasion, Shrub Die-off, and Other Aspects of Shrub Biology and Management. Gen. Tech. Rep. INT-GTR-276. USDA, Forest Service, Intermountain Research Station, Ogden, UT.

Pearson, L. 1964. Effect of harvest date on recovery of range grasses and shrubs. *Agronomy Journal* 56:80-82.

Pearson, L. C. 1965. Primary Production in Grazed and Ungrazed Desert Communities of Eastern Idaho. *Ecology* 46:278-285.

Peters, E. F. and S. Bunting. 1994. Fire conditions pre- and post-occurrence of annual grasses on the Snake River Plain. Pages 31-36 in Proceedings--Ecology and Management of Annual Rangelands Gen. Tech. Rep. INT-GTR-313. USDA, Forest Service, Intermountain Research Station, Boise, ID.

Quinones, F. A. 1981. Indian ricegrass evaluation and breeding. Bulletin 681. Page 19. New Mexico State University, Agricultural Experiment Station, Las Cruces, NM.

Sanderson, S. C., H. C. Stutz, and E. D. McArthur. 1990. Geographic Differentiation in *Atriplex confertifolia*. *American Journal of Botany* 77:490-498.

Smith, S. D. and P. S. Nobel. 1986. Deserts. Pages 13-62 in *Photosynthesis in contrasting environments*. Elsevier Science Publishers, Amsterdam, The Netherlands.

Smoliak, S., J. F. Dormaar, and A. Johnston. 1972. Long-Term Grazing Effects on *Stipa-Bouteloua* Prairie Soils. *Journal of Range Management* 25:246-250.

Stringham, T.K., P. Novak-Echenique, P. Blackburn, C. Coombs, D. Snyder and A. Wartgow. 2015. Final Report for USDA Ecological Site Description State-and-Transition Models, Major Land Resource Area 28A and 28B Nevada. University of Nevada Reno, Nevada Agricultural Experiment Station Research Report 2015-01. p. 1524.

Stubbendieck, J. L. 1985. Nebraska Range and Pasture Grasses: (including Grass-like Plants). University of Nebraska, Department of Agriculture, Cooperative Extension Service, Lincoln, NE.

Stubbendieck, J. L., S. L. Hatch, and C. H. Butterfield. 1992. North American range plants. University of Nebraska Press, Lincoln, NE.

Tueller, P. T. and W. H. Blackburn. 1974. Condition and Trend of the Big Sagebrush/Needle and thread Habitat Type in Nevada. *Journal of Range Management* 27:36-40.

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>).

Vallentine, J. F. 1989. Range development and improvements. Academic Press, Inc.

West, N. E. 1994. Effects of fire on salt-desert shrub rangelands. in Proceedings--Ecology and Management of Annual Rangelands, General Technical Report INT-313. USDA Forest Service, Intermountain Research Station, Boise, ID.

Wood, B. W. and J. D. Brotherson. 1986. Ecological adaptation and grazing response of budsage (*Artemisia*

spinescens) Pages 75-92 in Proceedings-- symposium on the biology of Artemisia and Chrysothamnus. Gen. Tech. Rep. INT-200. U. S. Department of Agriculture, Forest Service, Intermountain Research Station, Provo, UT.  
Wright, H. A. 1971. Why Squirreltail Is More Tolerant to Burning than Needle and thread. Journal of Range Management 24:277-284.

Wright, H. A. 1985. Effects of fire on grasses and forbs in sagebrush-grass communities. Pages 12-21 in Rangeland Fire Effects; A Symposium: Boise, ID, USDI-BLM.

Wright, H. A. and J. O. Klemmedson. 1965. Effect of Fire on Bunchgrasses of the Sagebrush-Grass Region in Southern Idaho. Ecology 46:680-688.

Young, J. A. and R. A. Evans. 1977. Squirreltail Seed Germination. Journal of Range Management 30:33-36.

Young, J. A. and F. Tipton. 1990. Invasion of cheatgrass into arid environments of the Lahontan Basin. Pages 37-40 in Proceedings- Symposium on cheatgrass invasion, shrub die-off, and other aspects of shrub biology and management. Gen. Tech. Rep. INT-276. USDA, Forest Service, Intermountain Research Station, Las Vegas, NV.

Young, R. P. 1983. Fire as a vegetation management tool in rangelands of the intermountain region. Pages 18-31 in Managing intermountain rangelands - improvement of range and wildlife habitats. USDA, Forest Service.

Zielinski, M. J. 1994. Controlling erosion on lands administered by the Bureau of Land Management, Winnemucca District, Nevada. Pages 143-146 in Proceedings - ecology and management of annual rangelands. Gen. Tech. Rep. INT-GTR-313. USDA, Forest Service, Intermountain Research Station, Ogden UT.

## Contributors

MD/RK  
T Stringham  
P NovakEchenique

## Approval

Kendra Moseley, 2/19/2025

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	P NOVAK-ECHENIQUE
Contact for lead author	State Rangeland Management Specialist
Date	05/15/2013
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:** This site is nearly level, thus rills are non-existent.
-

2. **Presence of water flow patterns:** These sites are subject to brief ponding and flooding after summer convection storms. As a result, water flow patterns may be numerous in areas. Flow patterns may be fairly long, meandering and stable.
- 
3. **Number and height of erosional pedestals or terracettes:** Pedestals are rare with occurrence typically limited to areas within water flow paths. Terracettes are non-existent.
- 
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare Ground 70-80%
- 
5. **Number of gullies and erosion associated with gullies:** None
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:** None - wind scouring may occur from a severe wind event during summer convection storms.
- 
7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage from grasses and annual & perennial forbs) expected to move distance of slope length during intense summer convection storms or rapid snowmelt events. Persistent litter (large woody material) will remain in place except during large rainfall or ponding events.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values should be 4 to 6 on most soil textures found on this site.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Surface structure is typically fine to medium platy or prismatic. Soil surface colors are pale browns and grays and soils are typified by an ochric epipedon. Surface textures are typically silt loams. A thick vesicular crust is common. Organic matter of the surface 2 to 3 inches is less than 1 percent.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Sparse shrub canopy and associated litter provide some protection from raindrop impact and allow for some snow capture on this site. Medium to fine textured surface soils have moderate to slow infiltration and medium runoff. The vesicular crust also impedes infiltration.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** Compacted layers are none. Massive sub-surface horizons or subsoil calcic horizons are not to be interpreted as compacted layers.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Reference State: Salt desert shrubs (shadscale)

Sub-dominant: shallow-rooted, cool season, perennial bunchgrasses > deep-rooted, cool season, perennial bunchgrasses > deep-rooted, cool season perennial forbs > associated salt-desert shrubs > fibrous, shallow-rooted, cool season, perennial forbs = annual forbs

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 35% of total woody canopy; mature bunchgrasses commonly ( $\pm 25\%$ ) have dead centers.
- 

14. **Average percent litter cover (%) and depth ( in):** Between plant interspaces (10-15%) and depth ( $< \frac{1}{4}$  in.)
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** For normal or average growing season (thru June)  $\pm 300$  lbs/ac; Favorable years  $\pm 400$  lbs/ac and unfavorable years  $\pm 200$  lbs/ac
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Potential invaders include halogeton, Russian thistle, annual mustards, and cheatgrass.
- 

17. **Perennial plant reproductive capability:** All functional groups should reproduce in average and above average growing season years. Little growth or reproduction occurs in extended or extreme drought conditions.
-