

Ecological site R029XY061NV FAN COLLAR 12-16 P.Z.

Last updated: 2/20/2025
Accessed: 05/11/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 029X–Southern Nevada Basin and Range

The Southern Nevada Basin and Range MLRA (29) represents the transition from the Mojave Desert to the Great Basin. It is cooler and wetter than the Mojave. It is warmer and typically receives more summer precipitation than the Great Basin. This area is in Nevada (73 percent), California (25 percent), and Utah (2 percent). It makes up about 26,295 square miles (68,140 square kilometers). Numerous national forests occur in the area, including the San Bernardino, Angeles, Sequoia, Inyo, Humboldt-Toiyabe, and Dixie National Forests. Portions of Death Valley National Monument, the Nuclear Regulatory Commission's Nevada Test Site, the Hawthorne Ammunition Depot, and the Nellis Air Force Range in Nevada and the China Lake Naval Weapons Center in California also are in this MLRA. The northeast part of the Paiute Indian Reservation and the southern third of the Walker River Indian Reservation are in the part of this MLRA in Nevada, and the Lone Pine, Fort Independence, and Big Pine Indian Reservations are in the part in California.

Physiography:

The entire area is in the Great Basin Section of the Basin and Range Province of the Intermontane Plateaus. The area of broad, nearly level, aggraded desert basins and valleys between a series of mountain ranges trending north to south. The basins are bordered by sloping fans and pluvial lake terraces. The mountains are uplifted fault blocks with steep side slopes and not well dissected due to limited annual precipitation. Most of the valleys in this MLRA are closed basins or bolsons containing sinks or playa lakes.

Geology:

The mountains are dominated by Pliocene and Miocene andesite and basalt rocks, Paleozoic and Precambrian carbonate rocks prominent in some areas. Scattered outcrops of older Tertiary intrusives and very young tuffaceous sediments (Pliocene and Miocene) are in the western and eastern thirds of this MLRA. The valleys consist mostly of alluvial fill and playa deposits at the lowest elevations in the closed basins.

Climate:

The average annual precipitation is 3 to 12 inches (75 to 305 millimeters) in most of this area. It may be as high as 29 inches (735 millimeters), on the higher mountain slopes. Most of the rainfall occurs as high-intensity, convective thunderstorms during the growing season. Summers are dry, but sporadic storms are common in July and August.

Water Resources:

Water resources are scarce. Ground water and surface water sources are limited. Streams are small and intermittent. Quality of surface water is naturally degraded as streams cross area of valley fill effected by dissolved salts. Irrigation water may raise the levels of dissolved salts and suspended sediments causing contamination.

Soils:

Dominant soil orders include Entisols and Aridisols.

Ecological site concept

This site is on alluvial fan collars. Slopes range from 8 to 15 percent. Elevations are 6,000 to 7,000 feet. The soils are deep and well drained.

Associated sites

F029XY095NV	PIMO-JUOS WSG 0R0501 12 to 16 This site is on mountain summits and sideslopes on all aspects. Slopes range from 15 to 75 percent. Elevations are 5,500 to about 8,600 feet. Soils of this site are typically shallow to moderately deep, deep and well drained. These soils have extreme amounts of very large, exfoliating ignimbritic boulders on the surface. Available water capacity is very low, but trees and shrubs extend their roots into fractures in the bedrock allowing them to utilize deep moisture.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata</i> var. <i>vaseyana</i> (2) <i>Purshia glandulosa</i>
Herbaceous	(1) <i>Leymus cinereus</i>

Physiographic features

This site occurs on alluvial fan collars. Slopes range from 8 to 15 percent. Elevations are 6,000 to 7,000 feet.

Table 2. Representative physiographic features

Landforms	(1) Alluvial fan
Runoff class	Low
Flooding duration	Very brief (4 to 48 hours)
Flooding frequency	Rare
Elevation	6,000–7,000 ft
Slope	8–15%
Water table depth	72 in
Aspect	Aspect is not a significant factor

Climatic features

The climate associated with this site is arid, characterized by cool, moist winters and hot, dry summers. Average annual precipitation is 12 to 16 inches. Mean annual air temperature is 45 to 50 degrees F. The average growing season is about 90 to 120 days. No climate stations are associated with this site.

Table 3. Representative climatic features

Frost-free period (average)	105 days
Freeze-free period (average)	
Precipitation total (average)	14 in




Figure 1. Monthly precipitation range




Figure 2. Monthly average minimum and maximum temperature




Figure 3. Annual precipitation pattern

Influencing water features

There are no influencing water features associated with this site.

Soil features

The soils are deep and well drained to well drained. The available water capacity is low. These soils have high amounts of stones, cobbles, and gravels throughout the soil profile, which reduce the available water holding capacity. The soils are usually moist in winter and spring and dry in summer and fall, except for 10 to 20 days between July and October due to convection storms. The soil moisture regime is aridic bordering on xeric and the soil temperature regime is mesic. Runoff is low. Soil series associated with this site include: Raster.

Table 4. Representative soil features

Parent material	(1) Alluvium
Surface texture	(1) Very gravelly coarse sand (2) Very stony sandy loam

Family particle size	(1) Sandy
Drainage class	Well drained
Permeability class	Moderately rapid
Soil depth	60 in
Surface fragment cover <=3"	22–65%
Surface fragment cover >3"	0–6%
Available water capacity (0-40in)	1.2–1.7 in
Calcium carbonate equivalent (0-40in)	0%
Electrical conductivity (0-40in)	0–2 mmhos/cm
Sodium adsorption ratio (0-40in)	0–2
Soil reaction (1:1 water) (0-40in)	6.6–7.8
Subsurface fragment volume <=3" (Depth not specified)	23–32%
Subsurface fragment volume >3" (Depth not specified)	4–36%

Ecological dynamics

As ecological condition deteriorates, Anderson peachbrush, mountain big sagebrush, and rubber rabbitbrush increase while basin wildrye, Utah serviceberry and desert bitterbrush decrease. Species likely to invade this site are annual forbs and grasses. Singleleaf pinyon and Utah juniper will increase on this site where it occurs adjacent to these woodlands.

Fire Ecology:

Pre-settlement fire return intervals in mountain big sagebrush communities varied from 15 to 25 years. Mountain big sagebrush is highly susceptible to injury from fire. It is often top-killed by fire and will not resprout. Desert bitterbrush is considered fire tolerant. Desert bitterbrush recovers from fire by sprouting from undamaged root crowns below the soil surface and by establishing from seeds cached by rodents. Desert bitterbrush often sprouts vigorously following fire, even in dry conditions. Aboveground parts of Utah serviceberry may be killed or consumed under fire conditions with sufficient flame lengths. Utah serviceberry may be slightly harmed by fire, depending on moisture conditions, but is generally considered to be fire tolerant. Utah serviceberry sprouts from the root crown following fire. Soil moisture is important to aid sprouting. Basin wildrye is top-killed by fire. Older basin wildrye plants with large proportions of dead material within the perennial crown can be expected to show higher mortality due to fire than younger plants having little debris. Basin wildrye is generally tolerant of fire but may be damaged by early season fire combined with dry soil conditions. Sand dropseed is usually killed or top killed by fire. Sand dropseed has the potential for postfire regeneration and seedling establishment as seeds within burned areas may remain viable. However postfire regeneration responses may differ according to relative abiotic and biotic site characteristics. Galleta is a rhizomatous perennial which can resprout after top-kill by fire.

The reference plant community is dominated by mountain big sagebrush, desert bitterbrush and basin wildrye. Potential vegetative composition is about 35 percent grasses, 5 percent forbs, 55 percent shrubs, and 5 percent trees. Approximate ground cover (basal and crown) is 35 to 55 percent.

State and transition model




Figure 5. DRAFT STM

State 1: Representative of the reference conditions prior to Euro-American settlement in the west.

- 1.1A: fire or other disturbance that removes sagebrush canopy
- 1.1B: absence of disturbance and natural regeneration over time
- 1.2A: absence of disturbance and natural regeneration over time
- 1.3A: fire or other disturbance that removes sagebrush canopy
- T1A: introduction of non-native species

State 2: Representative of the current potential with the presence of non-native annuals. Non-native annuals have the ability to significantly change disturbance regimes and nutrient cycling dynamics.

- 2.1A: fire or other disturbance that removes sagebrush canopy
- 2.1B: absence of disturbance and natural regeneration over time, may be coupled with inadequate rest and recovery from defoliation
- 2.2A: absence of disturbance and natural regeneration over time
- 2.3A: fire or other disturbance that removes sagebrush canopy
- T2A: long term absence of fire and natural regeneration of pinyon & juniper trees
- T2B: reoccurring severe fire

State 3: Dominated by pinyon and/or juniper trees. Changes in disturbance return intervals over the long term allows for pinyon and/or juniper to dominate the site by controlling site resources.

- 3.1A: absence of disturbance and natural regeneration over time
- T3A: reoccurring severe fire

State 4: Dominated by non-native annuals. Changes in disturbance return intervals and nutrient dynamics creating a positive feedback loop.

- 4.1A: absence of disturbance and natural regeneration over time
- 4.2A: fire or other disturbance that removed shrub canopy

Figure 6. DRAFT STM LEGEND

Animal community

Livestock Interpretations:

This site is suited to livestock grazing. Grazing management should be keyed to perennial grass production. The early growth and abundant production of basin wildrye make it a valuable source of forage for livestock. It is important forage for cattle and is readily grazed by cattle and horses in early spring and fall. Though coarse-

textured during the winter, basin wildrye may be utilized more frequently by livestock and wildlife when snow has covered low shrubs and other grasses. Sand dropseed provides fair to good forage for livestock. Sand dropseed's value as livestock forage is regional and dependent upon season. If fall rains are adequate, sand dropseed may have a period of renewed growth, producing new shoots in old sheaths. The persistent green base throughout winter makes sand dropseed an important desert winter range plant. In general, sand dropseed provides fair winter forage for domestic sheep and is most preferred by cattle of dune rangelands. When actively growing, galleta provides good to excellent forage for cattle and horses and fair forage for domestic sheep. Although not preferred, all classes of livestock may use galleta when it is dry. Domestic sheep show greater use in winter than summer months and typically feed upon central portions of galleta tufts, leaving coarser growth around the edges. Galleta may prove somewhat coarse to domestic sheep. Mountain big sagebrush is eaten by domestic livestock but has long been considered to be of low palatability, and a competitor to more desirable species. Desert bitterbrush is an important forage species for livestock, especially in winter. Utah serviceberry provides good browse for domestic sheep and domestic goats. In the spring, Utah serviceberry provides fair forage for cattle and good to excellent browse for domestic sheep and goats. Utah serviceberry provides good forage late in winter and in early spring, because it leafs out and blooms earlier than associated species.

Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year.

Wildlife Interpretations:

Mountain big sagebrush is highly preferred and nutritious winter forage for mule deer and elk. Sagebrush-grassland communities provide critical sage-grouse breeding and nesting habitats. Open Wyoming sagebrush communities are preferred nesting habitat. Meadows surrounded by sagebrush may be used as feeding and strutting grounds. Sagebrush is a crucial component of their diet year-round, and sage-grouse select sagebrush almost exclusively for cover. Lek sites are often located on low sagebrush sites, grassy openings, dry meadows, ridgetops, and disturbed sites. Sage-grouse prefer mountain big sagebrush and Wyoming big sagebrush communities to basin big sagebrush communities. Desert bitterbrush is an important forage species for deer and pronghorn, especially in winter. Upright growth forms of desert bitterbrush provide cover for game animals. Utah serviceberry is a very important species for mule deer in the Great Basin. Porcupines and desert bighorn sheep also use Utah serviceberry. Utah serviceberry fruit is preferred by many birds. It can be an important winter food for birds since berries stay on the shrub throughout the winter. In Nevada, sage grouse eat the fruit of Utah serviceberry. Basin wildrye provides winter forage for mule deer, though use is often low compared to other native grasses. Basin wildrye provides summer forage for black-tailed jackrabbits. Because basin wildrye remains green throughout early summer, it remains available for small mammal forage for longer time than other grasses. Sand dropseed provides poor forage for wildlife. Large mammals in general show little use of sand dropseed. Sand dropseed is not preferred by pronghorn, elk, and deer. Small mammals and birds utilize sand dropseed to a greater extent than large mammals. Galleta provides moderately palatable forage when actively growing and relatively unpalatable forage during dormant periods. Galleta provides poor cover for most wildlife species.

Hydrological functions

Runoff is low. Permeability is moderately rapid to very rapid.

Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site is used for camping and hiking and has potential for upland and big game hunting.

Other products

Native peoples used big sagebrush leaves and branches for medicinal teas, and the leaves as a fumigant. Bark was woven into mats, bags and clothing. Utah serviceberry fruits were used by Native Americans and early European explorers in North America for food and medicine. Basin wildrye was used as bedding for various Native American ceremonies, providing a cool place for dancers to stand. Sand dropseed is an edible grass used by Native Americans.

Other information

Utah serviceberry has been used to revegetate big game winter range and for surface stabilization. It grows slowly from seed and therefore transplanting may be more successful than seeding for revegetation projects. Basin wildrye is useful in mine reclamation, fire rehabilitation and stabilizing disturbed areas. Its usefulness in range seeding, however, may be limited by initially weak stand establishment. Sand dropseed is recommended as a component of grass seed mixtures for sandy and heavy to semi-sandy soils. Good results are seen reseeding dry low lands receiving less than 9 inches (230 mm) of precipitation within rangelands of Nevada.

Inventory data references

NASIS data used for abiotic narratives and tables.

Type locality

Location 1: Nye County, NV	
Township/Range/Section	T9N R51E S21
Latitude	38° 37' 34"
Longitude	116° 14' 13"
General legal description	NE¼ Section 21, T9N. R51E. MDBM. About 4 air miles southwest of Moores Station (Site), 3-way exclosure on Morey Bench, Hot Creek Range, Nye County, Nevada.

Other references

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).

United States Department of Agriculture, Natural Resources Conservation Service. 2022. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture, Agriculture Handbook 296.

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>)

Contributors

GED

Approval

Kendra Moseley, 2/20/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	State Rangeland Management Specialist
Date	10/01/1985
Approved by	Kendra Moseley
Approval date	

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-