

Ecological site R029XY077NV SHALLOW GRAVELLY LOAM 8-10 P.Z.

Last updated: 2/20/2025
Accessed: 05/11/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	P NOVAK-ECHENIQUE
Contact for lead author	State Rangeland Management Specialist
Date	07/12/2012
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- Number and extent of rills:** Rills are none to rare. A few rills (<1/10 m or 30 ft) will occur, especially following summer convective storms due to very shallow skeletal soils and steep slopes.
- Presence of water flow patterns:** None to rare. A few (<1/10 m or 30 ft) water flow patterns may occur in interspaces between shrubs, rarely connected. These should be limited to times following intense summer storms on steeper slopes or to natural drainages within the ecological site.
- Number and height of erosional pedestals or terracettes:** None to rare. Should only occur when associated with rills or water flow patterns on steeper slopes or natural drainages. Height less than 0.5 inch.
- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare Ground 5 to 20 percent. Soil is mostly covered by gravels, vegetation and some lichens and mosses. When patches of bare ground (3 foot diameter) occur, they should be associated with rodent burrow activity.
- Number of gullies and erosion associated with gullies:** None
- Extent of wind scoured, blowouts and/or depositional areas:** None

-
7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage from grasses and annual and perennial forbs) expected to move distance of slope length (<10 feet) during intense summer convection storms or rapid snowmelt events. Persistent litter (large woody material) will remain in place except during intense summer storms.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values are typically 1 to 3 in the interspaces and 4 to 6 under canopy.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** A-horizon thickness can range from 0 to 3 inches. Surface structure is typically moderate thin to medium platy. Soil surface colors are very pale brown to brown and soils are typified by an ochric epipedon. Organic matter of the surface horizon is typically less than 1 percent dropping off quickly below. Organic matter content can be more or less depending on micro-topography.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Soils have moderate permeability. Deep-rooted bunchgrasses and shrub canopy break raindrop impact, slow runoff and increase infiltration. Shrub canopy and standing dead grasses provide some opportunity for snow catch on this site.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** Compacted layers are none. Subsoil calcic and petrocalcic horizons are not to be interpreted as compacted.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: non-sprouting evergreen shrubs
- Sub-dominant: deep-rooted, cool-season, perennial bunchgrasses = deciduous shrubs > shallow-rooted cool-season perennial bunchgrasses > deep-rooted, cool-season perennial forbs <> annual forbs <> succulents
- Other: warm-season, perennial grasses, biological soil crust
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 30 percent of total woody canopy; some of the mature bunchgrasses (<20 percent) have dead centers, especially with multi-year droughts.
-
14. **Average percent litter cover (%) and depth (in):** Reference Plant Community: Litter cover is high under shrubs and

grasses and is between 10 to 15 percent in plant interspaces. Litter depth, where it occurs, is 0.25 inches.

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** For normal or average growing season (through mid-June) is ± 500 lbs/ac, Favorable years ± 700 lbs/ac, unfavorable years ± 300 lbs/ac
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Red brome, Mediterranean grass, red-stem filaree, and cheatgrass invade and persist on this site.
-
17. **Perennial plant reproductive capability:** All functional groups should reproduce in average (or normal) and above average growing season years. Less reproduction, although, rarely none, will occur in below-average precipitation years.
-