

Ecological site R029XY168NV SHALLOW ERODED SLOPE 8-10

Accessed: 05/14/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	P NOVAK-ECHENIQUE
Contact for lead author	State Rangeland Management Specialist
Date	05/20/2013
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:** Rills are common on steeper slopes and most frequently occur in areas subjected to summer convection storms or rapid spring snowmelt.

- 2. Presence of water flow patterns:** Water flow patterns are rare to slight.

- 3. Number and height of erosional pedestals or terracettes:** Pedestals are rare to slight and most frequently occur in areas subjected to summer convection storms

- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare Ground 50-60%; surface rock fragments less than 10%; shrub canopy <15%; foliar cover for perennial herbaceous plants <25%.

- 5. Number of gullies and erosion associated with gullies:** None

- 6. Extent of wind scoured, blowouts and/or depositional areas:** None

7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage from grasses and annual & perennial forbs) expected to move distance of slope length during intense summer storms. Persistent litter (large woody material) will remain in place except during large rainfall events.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values should be 1 to 4 on the coarse surface soil textures found on this site. (To be field tested.)
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Surface structure is typically single grain. Soil surface colors are pale browns and soils are typified by an ochric epipedon. Organic matter of the surface 2 to 3 inches is typically less than 1 percent dropping off quickly below. Organic matter content can be more or less depending on micro-topography.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Sparse shrub canopy and associated litter provide some protection from raindrop impact.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: low statured shrubs (snakeweed, black sagebrush)
- Sub-dominant: shallow-rooted, cool season, perennial grasses = deep-rooted, cool season, perennial forbs = fibrous, shallow-rooted, cool season, annual and perennial forbs > shallow-rooted, warm season, perennial grasses
- Other: evergreen trees
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 30% of total woody canopy; some of the mature bunchgrasses (to 25%) have dead centers.
-
14. **Average percent litter cover (%) and depth (in):** Between plant interspaces 20-30%; depth < ¼ inch
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** For normal or average growing season (March thru May) ± 125 lbs/ac; Favorable years ±200 lbs/ac and unfavorable years ± 75 lbs/ac
-

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Potential invaders include cheatgrass, halogeton, Russian thistle, red-stem filaree, and annual mustards.
-

17. **Perennial plant reproductive capability:** All functional groups should reproduce in average (or normal) and above average growing season years. Little reproduction or growth occurs during drought years.
-