

Ecological site R030XB031NV SHALLOW LIMY 5-7 P.Z.

Last updated: 2/18/2025
Accessed: 05/13/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 030X–Mojave Basin and Range

The Mojave Desert Major Land Resource Area (MLRA 30) is found in southern California, southern Nevada, the extreme southwest corner of Utah and northwestern Arizona within the Basin and Range Province of the Intermontane Plateaus. The Mojave Desert is a transitional area between hot deserts and cold deserts where close proximity of these desert types exert enough influence on each other to distinguish these desert types from the hot and cold deserts beyond the Mojave. Kottek et. al 2006 defines hot deserts as areas where mean annual air temperatures are above 64 F (18 C) and cold deserts as areas where mean annual air temperatures are below 64 F (18 C). Steep elevation gradients within the Mojave create islands of low elevation hot desert areas surrounded by islands of high elevation cold desert areas.

The Mojave Desert receives less than 10 inches of mean annual precipitation. Mojave Desert low elevation areas are often hyper-arid while high elevation cold deserts are often semi-arid with the majority of the Mojave being an arid climate. Hyper-arid areas receive less than 4 inches of mean annual precipitation and semi-arid areas receive more than 8 inches of precipitation (Salem 1989). The western Mojave receives very little precipitation during the summer months while the eastern Mojave experiences some summer monsoonal activity.

In summary, the Mojave is a land of extremes. Elevation gradients contribute to extremely hot and dry summers and cold moist winters where temperature highs and lows can fluctuate greatly between day and night, from day to day and from winter to summer. Precipitation falls more consistently at higher elevations while lower elevations can experience long intervals without any precipitation. Lower elevations also experience a low frequency of precipitation events so that the majority of annual precipitation may come in only a couple precipitation events during the whole year. Hot desert areas influence cold desert areas by increasing the extreme highs and shortening the length of below freezing events. Cold desert areas influence hot desert areas by increasing the extreme lows and increasing the length of below freezing events. Average precipitation and temperature values contribute little understanding to the extremes which govern wildland plant communities across the Mojave.

Arid Eastern Mojave Land Resource Unit (XB)

LRU notes

The Mojave Desert is currently divided into 4 Land Resource Units (LRUs). This ecological site is within the Arid Eastern Mojave LRU where precipitation is bi-modal, occurring during the winter months and summer months. The Arid Eastern Mojave LRU is designated by the 'XB' symbol within the ecological site ID. This LRU is found across the eastern half of California, much of the mid-elevations of Nevada, the southernmost portions of western Utah, and the mid-elevations of northwestern Arizona. This LRU is essentially equivalent to the Eastern Mojave Basins and Eastern Mojave Low Ranges and Arid Footslopes of EPA Level IV Ecoregions

Elevations range from 1650 to 4000 feet and precipitation is between 4 to 8 inches per year. This LRU is

distinguished from the Arid Western Mojave (XA) by the summer precipitation, falling between July and September, which tends to support more warm season plant species. The 'XB' LRU is generally east of the Mojave River and the 117 W meridian (Hereford et. al 2004). Vegetation includes creosote bush, burrobush, Nevada jointfir, ratany, Mojave yucca, Joshua tree, cacti, big galleta grass and several other warm season grasses. At the upper portions of the LRU, plant production and diversity are greater and blackbrush is a common dominant shrub.

Ecological site concept

This site is found on active alluvial fans in the northern Mojave Desert where average minimum temperatures for December and January are below 32 degrees F. Alluvium is derived from volcanic parent material.

Associated sites

R030XB010NV	LOAMY SLOPE 5-7 P.Z.
R030XB029NV	SHALLOW GRAVELLY LOAM 5-7 P.Z.

Similar sites

R030XB074NV	COBBLY LOAM 5-7 P.Z. ATCO rare to absent
R030XB005NV	Arid Active Alluvial Fans MESP2 typically rare to absent
R030XB041NV	SHALLOW DROUGHTY LOAM 5-7 P.Z. GRSP codominant shrub; more productive site
R030XB006NV	LOAMY 5-7 P.Z. MESP2 typically rare to absent

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Atriplex confertifolia</i> (2) <i>Menodora spinescens</i>
Herbaceous	(1) <i>Achnatherum hymenoides</i>

Physiographic features

This site occurs on sideslopes and summits of fan remnants on all exposures. Slopes range from 0 to 30 percent, but slope gradients of 4 to 8 percent are most typical. Elevations are 2000 to 4000 feet.

Table 2. Representative physiographic features

Landforms	(1) Fan remnant
Elevation	610–1,219 m
Slope	0–30%
Aspect	Aspect is not a significant factor

Climatic features

The climate of the Mojave Desert has extreme fluctuations of daily temperatures, strong seasonal winds, and clear skies. The climate is arid and is characterized with cool, moist winters and hot, dry summers. Most of the rainfall falls between November and April. Summer convection storms from July to September may contribute up to 25 percent of the annual precipitation. Average annual precipitation is 5 to 7 inches. Mean annual air temperature is 58 to 65 degrees F. The average growing season is about 180 to 250 days.

Table 3. Representative climatic features

Frost-free period (average)	250 days
Freeze-free period (average)	
Precipitation total (average)	178 mm




Figure 1. Monthly average minimum and maximum temperature

Influencing water features

There are no influencing water features associated with this site.

Soil features

The soils associated with this site are shallow to a duripan or petrocalcic layer. They are well drained, have very high runoff, and have moderate permeability. These soils have typically formed in alluvium from ignimbritic and basalt parent material. Available water holding capacity is very low. The soil series associated with this site is Alko.

Table 4. Representative soil features

Parent material	(1) Alluvium–ignimbrite
Surface texture	(1) Gravelly sandy loam (2) Coarse sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderate
Soil depth	25–51 cm
Surface fragment cover <=3"	10–20%
Surface fragment cover >3"	0–3%
Available water capacity (0-101.6cm)	3.05–3.3 cm
Calcium carbonate equivalent (0-101.6cm)	5–15%
Electrical conductivity (0-101.6cm)	0–4 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–5
Soil reaction (1:1 water) (0-101.6cm)	7.9–9
Subsurface fragment volume <=3" (Depth not specified)	13–24%

Subsurface fragment volume >3" (Depth not specified)	2–3%
---	------


Ecological dynamics

As ecological condition deteriorates, shadscale, creosotebush, wolfberry and white bursage increase while Indian ricegrass, big galleta and spiny menodora decrease. Species likely to invade this site are annuals such as red brome.


Fire Ecology:
 Fires in the Mojave Desert are infrequent and of low severity because production of annual and perennial herbs seldom provides a fuel load capable of sustaining fire. Fire generally kills white bursage. Shadscale communities are usually unaffected by fire because of low fuel loads, although a year of exceptionally heavy winter rains can generate fuels by producing a heavy stand of annual forbs and grasses. The mean fire return interval for shadscale communities range from 35 to 100 years. Increased presence of non-native annual grasses, such as cheatgrass, can alter fire regimes by increasing fire frequency under wet to near-normal summer moisture conditions. Spiny menodora often survives fire because its foliage does not readily burn. Nevada ephedra is top-killed by fire. Underground regenerative structures commonly survive when aboveground vegetation is consumed by fire. Nevada ephedra generally sprouts after fire damages aboveground vegetation and may increase in plant cover. Bud sagebrush is killed by fire. Fire kills many creosotebush. Creosotebush is poorly adapted to fire because of its limited sprouting ability. Creosotebush survives some fires that burn patchily or are of low severity. Indian ricegrass can be killed by fire, depending on severity and season of burn. Indian ricegrass reestablishes on burned sites through seed dispersed from adjacent unburned areas. Fire most likely top-kills big galleta. Big galleta sprouts from rhizomes following fire. Damage to big galleta from fire varies, depending on whether big galleta is dormant when burned. If big galleta is dry, damage may be severe. However, when plants are green, fire will tend to be less severe and damage may be minimal, with big galleta recovering quickly. Desert needlegrass has persistent dead leaf bases, which make it susceptible to burning. Fire removes the accumulation; a rapid, cool fire will not burn deep into the root crown and surviving tufts will resprout.

State and transition model


Ecosystem states



State 1 submodel, plant communities



State 2 submodel, plant communities



State 1 Reference State

The reference state is representative of the natural range of variability under pristine conditions. Plant communities are dynamic in response to changes in disturbance regimes and weather patterns. Plant community phase changes are primarily driven by long-term drought. Historically, fire had little impact in this system due to low fuel loading and widely spaced vegetation.

Community 1.1

Reference Plant Community

The reference plant community is dominated by shadscale and spiny menodora. Other important species associated with this site include white bursage, Indian ricegrass, big galleta and desert needlegrass. Potential vegetative composition is about 15% annual and perennial grasses, 5% forbs and 80% shrubs. Approximate ground cover (basal and crown) is 10 to 15 percent.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Shrub/Vine	135	269	448
Grass/Grasslike	26	50	84
Forb	8	17	28
Total	169	336	560

State 2

Representative State

Introduced annuals such as red brome, schismus and redstem stork's bill have invaded the reference plant community and have become a dominant component of the herbaceous cover. Dominant species persist after invasion by non-native annuals but non-native annuals are extremely difficult to remove from the system. When dried, non-natives annuals can be highly flammable and may increase fire frequency where fires historically have been infrequent. Establishment of non-natives is attributed to a combination of factors including; surface disturbances, changes in the kinds of animals and their grazing patterns, drought, and changes in fire history.

Community 2.1

Representative Plant Community

This plant community is similar to the reference plant community with a trace of non-natives in the understory. Ecological function has been not compromised at this time. Ecological resilience is reduced by the presence of non-native species and this plant community phase will respond differently following a disturbance when compared to non-invaded plant communities.

Community 2.2

Plant Community 2.2

This plant community is characteristic of a post-disturbance plant community. It is initially dominated by herbaceous vegetation, woody perennials are increasing. Short lived and pioneering shrubs such as cattle saltbush, rabbit brush, California buckwheat, spiny hopsage, and burrobrush provide favorable microsites for the establishment of long lived shrub seedlings.

Pathway 2.1a

Community 2.1 to 2.2

Frequent and repeated surface disturbances, wildfire, disease, insect attack, or any other type of incomplete vegetation removal.

Pathway 2.2a

Community 2.2 to 2.1

Absence from disturbance and natural regeneration over time.

Transition T1 State 1 to 2

Introduction of non-native species due to a combination of factors including; surface disturbance, changes in the kinds of animals and their grazing patterns, drought, changes in fire history or any other type of vegetation removal.

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Primary Perennial Grasses			24–81	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	7–27	–
	low woollygrass	DAPU7	<i>Dasyochloa pulchella</i>	3–20	–
	big galleta	PLRI3	<i>Pleuraphis rigida</i>	7–17	–
	desert needlegrass	ACSP12	<i>Achnatherum speciosum</i>	7–17	–
2	Secondary Perennial Grasses			7–17	
	Mormon needlegrass	ACAR14	<i>Achnatherum aridum</i>	2–7	–
3	Annual			1–10	
Forb					
4	Perennial			7–27	
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	2–7	–
	princesplume	STANL	<i>Stanleya</i>	2–7	–
5	Annual			1–10	
Shrub/Vine					
6	Primary shrubs			128–269	
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	50–101	–
	spiny menodora	MESP2	<i>Menodora spinescens</i>	34–67	–
	burrobush	AMDU2	<i>Ambrosia dumosa</i>	17–34	–
	Nevada jointfir	EPNE	<i>Ephedra nevadensis</i>	7–17	–
	creosote bush	LATR2	<i>Larrea tridentata</i>	7–17	–
	desert-thorn	LYCIU	<i>Lycium</i>	7–17	–
7	Secondary shrubs			7–50	
	spiny hopsage	GRSP	<i>Grayia spinosa</i>	3–10	–
	burrobrush	HYSA	<i>Hymenoclea salsola</i>	3–10	–
	Fremont's dalea	PSFR	<i>Psoralea fremontii</i>	3–10	–
	yucca	YUCCA	<i>Yucca</i>	3–10	–

Animal community

Livestock Interpretations:

This site is suited to livestock grazing. Grazing management should be keyed to perennial grass and palatable shrub production. Indian ricegrass has good forage value for domestic sheep, cattle and horses. It supplies a source of green feed before most other native grasses have produced much new growth. Big galleta is considered a valuable forage plant for cattle and domestic sheep. Its coarse, rigid culms make it relatively resistant to heavy grazing and trampling. Desert needlegrass produces considerable basal foliage and is good forage while young.

Young desert needlegrass is palatable to all classes of livestock. Mature herbage is moderately grazed by horses and cattle but rarely grazed by sheep. Shadscale provides good browse for domestic sheep and goats. Shadscale leaves and seeds are an important component of domestic sheep and cattle winter diets. Shadscale tends to be browse tolerant. Heavy grazing during the winter and/or spring reduces shadscale. Die-off can also occur during extended periods of high precipitation. Shadscale is tolerant of early spring light-intensity browsing. Cattle will graze the stems of spiny menodora in the spring before the stems become woody and spiny. Spiny menodora has lower palatability than the other shrubs but is consumed during early spring before spines mature. White bursage is of intermediate forage value. It is fair to good forage for horses and fair to poor for cattle and sheep. However, because there is often little other forage where white bursage grows, it is often highly valuable to browsing animals and is sensitive to browsing. Nevada ephedra is important winter range browse for domestic cattle, sheep and goats. Budsage is palatable and nutritious forage for domestic sheep in the winter and spring although it is known to cause mouth sores in lambs. Budsage can be poisonous or fatal to calves when eaten in quantity. Budsage, while desired by cattle in spring, is poisonous to cattle when consumed alone. Creosotebush is unpalatable to livestock. Consumption of creosotebush may be fatal to sheep.

Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year.

Wildlife Interpretations:

Shadscale is a valuable browse species providing a source of palatable, nutritious forage for a wide variety of wildlife. The fruits and leaves are a food source for deer, desert bighorn sheep and pronghorn antelope. Elk will graze the stems of spiny menodora in the spring before the stems become woody and spiny. White bursage is an important browse species for wildlife. Mule deer, bighorn sheep, and pronghorn browse Nevada ephedra, especially in spring and late summer when new growth is available. Mountain quail eat ephedra seeds. Budsage is palatable, nutritious forage for upland game birds, small game and big game in winter. Budsage is rated as valuable forage for mule deer in Nevada in winter and is utilized by bighorn sheep in summer, but the importance of budsage in the diet of bighorns is not known. Bud sage comprises 18 – 35% of a pronghorn's diet during the spring where it is available. Chukar will utilize the leaves and seeds of bud sage. Budsage is highly susceptible to effects of browsing. It decreases under browsing due to year-long palatability of its buds and is particularly susceptible to browsing in the spring when it is physiologically most active. Creosotebush is unpalatable to most browsing wildlife. Indian ricegrass is eaten by pronghorn in moderate amounts whenever available. In Nevada it is consumed by desert bighorns. A number of heteromyid rodents inhabiting desert rangelands show preference for seed of Indian ricegrass. Indian ricegrass is an important component of jackrabbit diets in spring and summer. In Nevada, Indian ricegrass may even dominate jackrabbit diets during the spring through early summer months. Indian ricegrass seed provides food for many species of birds. Doves, for example, eat large amounts of shattered Indian ricegrass seed lying on the ground. Desert bighorn sheep and feral horses and burros will graze desert needlegrass.

Hydrological functions

Runoff is low. Permeability is moderate.

Other products

Indian ricegrass was traditionally eaten by some Native Americans. The Paiutes used seed as a reserve food source. Seeds of shadscale were used by Native Americans for bread and mush. White bursage is a host for sandfood, a parasitic plant. Sandfood was a valuable food supply for Native Americans. Some Native American tribes steeped the twigs of Nevada ephedra and drank the tea as a general beverage. Creosotebush has been highly valued for its medicinal properties by Native Americans. Twigs and leaves may be boiled as tea, steamed, pounded into a powder, pressed into a poultice, or heated into an infusion.

Other information

Big galleta's clumped growth form stabilizes blowing sand. Desert needlegrass may be used for groundcover in areas of light disturbance, but it is susceptible to excessive trampling. White bursage may be used to revegetate disturbed sites in southwestern deserts. Once established, creosotebush may improve sites for annuals that grow under its canopy by trapping fine soil, organic matter, and symbiont propagules. It may also increase water

infiltration and storage.

Type locality

Location 1: Lincoln County, NV	
Township/Range/Section	T5S R60E S1
General legal description	Approximately 1 mile east of Hiko Junction on north side of US Highway 93, Lincoln County, Nevada. This site also occurs in Clark County, Nevada.

Other references

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).

Hereford, R., R.H. Webb and C. I. Longpre. 2004. Precipitation history of the Mojave Desert region, 1893-2001 (No. 117-03).

Kottek, M., Grieser, J., Beck, C., Rudolf, B., & Rubel, F. (2006). World map of the Köppen-Geiger climate classification updated. Meteorologische Zeitschrift, 15(3), 259-263.

Salem, B. B. (1989). Arid zone forestry: a guide for field technicians (No. 20). Food and Agriculture Organization (FAO).

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>).

Contributors

RRK/BO'D
Dustin Detweiler

Approval

Kendra Moseley, 2/18/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/13/2025
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
