

## Ecological site R030XB052NV RUBBLY OUTWASH

Last updated: 2/24/2025  
Accessed: 05/13/2025

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	P NOVAK-ECHENIQUE
Contact for lead author	State Rangeland Management Specialist
Date	04/08/2010
Approved by	Sarah Quistberg
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

- 1. Number and extent of rills:** Rills are none to rare.  

---
- 2. Presence of water flow patterns:** Water flow patterns few to common.  

---
- 3. Number and height of erosional pedestals or terracettes:** Pedestals are rare with occurrence typically limited to areas within water flow patterns.  

---
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare Ground to 20%; surface rock fragments to 85%; shrub canopy to 15%; basal area for perennial herbaceous plants (Trace).  

---
- 5. Number of gullies and erosion associated with gullies:** None  

---
- 6. Extent of wind scoured, blowouts and/or depositional areas:** None  

---

7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage from grasses and annual & perennial forbs) expected to move distance of slope length during intense summer convection storms or rapid snowmelt events. Persistent litter (large woody material) will remain in place except during catastrophic events.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values should be 2 to 4 on most soil textures found on this site. Areas of this site occurring on soils that have a physical crust will probably have stability values less than 3. (To be field tested.)
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Surface structure is typically weak coarse platy. Soil surface colors are light and soils are typified by an ochric epipedon. Organic matter of the surface 2 to 3 inches is less than 0.5 percent.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Sparse shrub canopy and associated litter break raindrop impact.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Reference Plant Community: Mojave Desert shrubs
- Sub-dominant: deep-rooted, warm-season, bunchgrasses > perennial forbs > deep-rooted, cool-season, bunchgrasses > annual forbs > annual grasses
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 25% of total woody canopy.
- 
14. **Average percent litter cover (%) and depth ( in):** Between plant interspaces <5% and depth ( $\pm\frac{1}{4}$ -inch).
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** For normal or average growing season  $\pm 300$ lbs/ac.
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if**

their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Invaders on this site include annual grasses and forbs, such as red brome, Mediterranean grass, and filaree.

---

17. **Perennial plant reproductive capability:** All functional groups should reproduce in above average growing season years.
-