

Ecological site R030XB207AZ Granitic Hills 6-9" p.z.

Last updated: 10/21/2024
Accessed: 05/11/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 030X–Mojave Basin and Range

AZ LRU 30-2 – Middle Mohave Desert

Elevations range from 1500 to 3200 feet and precipitation averages 6 to 9 inches per year. Vegetation includes creosotebush, white bursage, yucca, prickly pear and cholla species, Mormon tea, flattop buckwheat, ratany, winterfat, bush muhly, threeawns, and big galleta. The soil temperature regime is thermic and the soil moisture regime is typic aridic. This unit occurs within the Basin and Range Province and is characterized by broad basins, valleys, and old lakebeds. Widely spaced mountains trending north to south occur throughout the area. Isolated, short mountain ranges are separated by an aggraded desert plain. The mountains are fault blocks that have been tilted up. Long alluvial fans coalesce with dry lakebeds between some of the ranges.

LRU notes

AZ LRU 30-2 – Middle Mohave Desert

Elevations range from 1500 to 3200 feet and precipitation averages 6 to 9 inches per year. Vegetation includes creosotebush, white bursage, yucca, prickly pear and cholla species, Mormon tea, flattop buckwheat, ratany, winterfat, bush muhly, threeawns, and big galleta. The soil temperature regime is thermic and the soil moisture

regime is typic aridic.

Ecological site concept

This ecological site is located on steeply sloping uplands (slopes 15%-65%). Soils are very shallow to shallow to granite or gneiss bedrocks.

Associated sites

R030XB214AZ	Limy Upland 6-9" p.z. Limy Upland, 6 to 10
R030XB226AZ	Sandy Loam Upland 6-9" p.z. Fine Sandy Loam Upland, fine, 6 to 10

Similar sites

R030XA104AZ	Granitic Hills 3-6" p.z. Granitic Hills, 3 to 6
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Ambrosia dumosa</i> (2) <i>Encelia farinosa</i>
Herbaceous	Not specified

Physiographic features

This ecological site is found in an upland position on shoulders, sideslopes and summits of hills of granitic parent material. Granitic or gneiss rock outcrop is common in association with this site. This site occurs on all aspects.

Table 2. Representative physiographic features

Landforms	(1) Hill
Flooding frequency	None
Ponding frequency	None
Elevation	2,150–3,800 ft
Slope	20–65%

Climatic features

The climate is arid and warm. Annual precipitation ranges from 6 to 9 inches. About 65 percent of the rainfall comes from October through May as gentle rain from Pacific storms which may last for a couple of days. The rest of the rainfall comes during the summer monsoon season from July through September as spotty, brief, intense thunderstorms. Snow rarely falls, and only remains on the ground a few hours at most. Annual air temperature ranges from 59 to 70 degrees F. The average frost-free period ranges from 156 to 259 days.

Table 3. Representative climatic features

Frost-free period (average)	259 days
Freeze-free period (average)	290 days
Precipitation total (average)	9 in

Influencing water features

Soil features

The soil of this ecological site is very shallow to shallow and is of granitic alluvium and colluvium parent materials. The subsoil is very gravelly sandy loam. The available water holding capacity is very low. The soil's erosion hazard by water is moderate to very severe and by wind is slight. The soil is non-sodic, non-saline with 7.6-7.8 pH. The soil moisture regime is typic aridic and soil temperature is thermic. Rock outcrop is associated with this site. Depth to hard granite or gneiss is 6-13 inches.

A typical soil profile is:

A-0 to 1 inches; very gravelly sandy loam

C-1 to 5 inches; gravelly coarse sandy loam

2Cr-5 to 8 inches; weathered bedrock

2R-8 inches; unweathered bedrock

The soil taxonomic classification is Loamy-skeletal, mixed, superactive, nonacid, thermic, shallow Typic Torriorthents

This ecological site has been correlated to map unit 627134, Wikieup soil, Mohave County, AZ, Southern Part SSA.

Table 4. Representative soil features


Surface texture	(1) Very gravelly sandy loam
Family particle size	(1) Loamy
Drainage class	Somewhat excessively drained
Permeability class	Moderately rapid
Soil depth	5–15 in
Surface fragment cover <=3"	45–70%
Surface fragment cover >3"	0–10%
Available water capacity (0-40in)	0.5–0.9 in
Calcium carbonate equivalent (0-40in)	0%
Electrical conductivity (0-40in)	0 mmhos/cm
Sodium adsorption ratio (0-40in)	0
Soil reaction (1:1 water) (0-40in)	7.4–8.4
Subsurface fragment volume <=3" (Depth not specified)	20–45%
Subsurface fragment volume >3" (Depth not specified)	0–10%

Ecological dynamics


Granitic Hills, 6"-9" p.z., is a shrub dominated ecological site. Sparse perennial grasses and forbs are occasionally encountered. Annual forbs and grasses flourish following rainfall. Natural disturbances are rare. Following introduction of non-native annuals (forbs and/or grasses), an increased seasonal herbaceous production following periods of rain. Dominant shrubs are creosote and white bursage. Assorted half-shrubs are widely scattered.

State and transition model


Ecosystem states



State 1 submodel, plant communities



State 2 submodel, plant communities



State 1
Historic Climax Plant Community

Community 1.1
Historic Climax Plant Community

The dominant aspect of this ecological site is a desert shrub. This site is dominated by creosotebush and white bursage. In some areas, blackbrush or white brittlebush are dominant. Perennial grasses are present in minor amounts. This site has a poor soil-plant moisture relationship. Steep slopes and rock hinder large animal movement. Degradation occurs only because of catastrophic events such as fire or man-caused disturbances (mining). If shrubs are lost, they would return very slowly leaving the soil exposed and susceptible to erosion. The site's total annual production by weight (air-dried) is comprised of: grasses and grasslike plants - 5 to 15 percent, forbs 5 to 10 percent, and shrubs 70 to 80 percent.

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Shrub/Vine	86	215	381
Grass/Grasslike	6	35	71
Forb	6	25	48
Total	98	275	500

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	0-2%
Grass/grasslike foliar cover	0%
Forb foliar cover	0%
Non-vascular plants	0%
Biological crusts	0%

Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 7. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	—	0-2%
>0.5 <= 1	—	—	0-2%	—
>1 <= 2	—	4-6%	—	—
>2 <= 4.5	—	—	—	—
>4.5 <= 13	—	—	—	—
>13 <= 40	—	—	—	—
>40 <= 80	—	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

Figure 5. Plant community growth curve (percent production by month). AZ3022, 30.2 6-9" p.z. upland sites. Growth begins in the late winter, most growth occurs in the spring..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	2	19	33	18	7	7	11	3	0	0	0

State 2

Historic Native Plant Community plus Introduced Annuals

Community 2.1

Non-native Annuals

This plant community resembles the historic native plant community, but exotic annuals have been introduced. Non-native species include red brome, Mediterranean grass (*Schismus* spp.), and filaree. The flourish of non-native annuals that occurs following rainfalls may preclude native annuals.

Transition T1A

State 1 to 2

Introduction of non-native annual forb and grass seed.

Restoration pathway R2A

State 2 to 1

None known.

Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
-------	-------------	--------	-----------------	--------------------------------	---------------------

Grass/Grasslike				
1				3–14
	big galleta	PLRI3	<i>Pleuraphis rigida</i>	3–14
2				6–14
	desert needlegrass	ACSP12	<i>Achnatherum speciosum</i>	6–14
3				3–8
	Parish's threeawn	ARPUP5	<i>Aristida purpurea</i> var. <i>parishii</i>	0–8
4				3–8
	Grass, perennial	2GP	<i>Grass, perennial</i>	3–8
5				3–14
	Grass, annual	2GA	<i>Grass, annual</i>	3–14
Forb				
6				3–11
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	3–11
7				3–14
	Forb, annual	2FA	<i>Forb, annual</i>	7–14
	Forb, annual	2FA	<i>Forb, annual</i>	3–14
8				0–3
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–3
Shrub/Vine				
9				14–41
	creosote bush	LATR2	<i>Larrea tridentata</i>	14–41
10				28–55
	burrobush	AMDU2	<i>Ambrosia dumosa</i>	28–55
11				14–28
	blackbrush	CORA	<i>Coleogyne ramosissima</i>	14–28
12				8–14
	Nevada jointfir	EPNE	<i>Ephedra nevadensis</i>	8–14
13				8–14
	littleleaf ratany	KRER	<i>Krameria erecta</i>	8–14
14				3–8
	rayless goldenhead	ACSP	<i>Acamptopappus sphaerocephalus</i>	3–8
15				8–14
	Eastern Mojave buckwheat	ERFAP	<i>Eriogonum fasciculatum</i> var. <i>polifolium</i>	8–14
16				0–6
	water jacket	LYAN	<i>Lycium andersonii</i>	0–6
17				0–6
	pricklypear	OPUNT	<i>Opuntia</i>	0–6
18				0–14
	Parish's goldeneye	VIPA14	<i>Viguiera parishii</i>	0–14
19				28–55
	Eastern Mojave buckwheat	ERFAP	<i>Eriogonum fasciculatum</i> var. <i>polifolium</i>	28–55

20				0–14	
	button brittlebush	ENFR	<i>Encelia frutescens</i>	0–14	–
21				0–14	
	Shrub, other	2S	<i>Shrub, other</i>	0–14	–

Contributors

Harmon Hodgkinson
Harmon S. Hodgkinson
Larry D. Ellicott
Stephen Cassady

Approval

Kendra Moseley, 10/21/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/11/2025
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. **Extent of wind scoured, blowouts and/or depositional areas:**
-
7. **Amount of litter movement (describe size and distance expected to travel):**
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state**

for the ecological site:

17. **Perennial plant reproductive capability:**
