

## Ecological site R034AY138WY Saline Lowland Green River and Great Divide Basins (SL)

Last updated: 2/24/2025  
Accessed: 05/12/2025

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	K. Clause, J. Haverkamp, E. Bainter
Contact for lead author	karen.clause@wy.usda.gov or 307-367-2257
Date	03/16/2007
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

- Number and extent of rills:** Rare to nonexistent.  

---
- Presence of water flow patterns:** Barely observable.  

---
- Number and height of erosional pedestals or terracettes:** Rare to nonexistent.  

---
- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground can range from 5-30%.  

---
- Number of gullies and erosion associated with gullies:** Active gullies should not be present.  

---
- Extent of wind scoured, blowouts and/or depositional areas:** Minimal to nonexistent.  

---
- Amount of litter movement (describe size and distance expected to travel):** Herbaceous litter not expected to move.

- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil Stability Index ratings range from 2 (interspaces) to 6 (under plant canopy), but average values should be 3.5 or greater.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Typically an A-horizon of 3-9 inches (7-23 cm) with weak to medium platy or sometimes granular structure and color hues of 10YR or 2.5Y, values of 6-8, and chromas of 2-4. Sometimes a shallow E-horizon of 2 inches (6 cm) with platy structure will replace the A-horizon. Organic matter typically ranges from .5 to 1%.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Plant community consists of 50-70% grasses, 10% forbs, and 20-40% shrubs. Dense plant canopy (70-90%) and litter plus moderate infiltration rates result in minimal to nonexistent runoff. Basal cover is typically 5-10% for this site and does effectively reduce runoff on this site.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer exists.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional: perennial shrubs>cool season rhizomatous grasses>warm season bunchgrasses>mid-size, cool season bunchgrasses=tall, cool season bunchgrasses>perennial forbs=warm season rhizomatous grasses
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Minimal decadence, typically associated with shrub component.
- 
14. **Average percent litter cover (%) and depth ( in):** Litter ranges from 10-30% of total canopy measurement with total litter (including beneath the plant canopy) from 70-95% expected. Herbaceous litter depth typically ranges from 10-25 mm. Woody litter can be up to a couple inches (4-6 cm).
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** English: 800-2000 lb/ac (1200 lb/ac average); Metric: 896-2240 kg/ha (1344 kg/ha average).
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize**

degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is **NOT** expected in the reference state for the ecological site: Bare ground greater than 40% is the most common indicator of a threshold being crossed. Greasewood, inland saltgrass, and alkali bluegrass are common increasers. Annual weeds such as halogeton, kochia, and Russian thistle are common invasive species in disturbed sites.

---

17. **Perennial plant reproductive capability:** All species are capable of reproducing, except in drought years.
-