

Ecological site R034AY274WY Subirrigated Foothills and Basins West (Sb)

Accessed: 05/12/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	K. Clause, J. Haverkamp, E. Bainter
Contact for lead author	karen.clause@wy.usda.gov or 307-367-2257
Date	03/16/2007
Approved by	E. Bainter
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- Number and extent of rills:** Rare to nonexistent.

- Presence of water flow patterns:** Water flow patterns sometimes evident in floodplain zone where this site occurs.

- Number and height of erosional pedestals or terracettes:** Rare to nonexistent.

- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground is typically less than 5%.

- Number of gullies and erosion associated with gullies:** Active gullies should not be present.

- Extent of wind scoured, blowouts and/or depositional areas:** Minimal to nonexistent.

- Amount of litter movement (describe size and distance expected to travel):** Herbaceous litter exhibits slight movement only associated with water flow patterns.

-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil Stability Index ratings typically 6.0.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface highly variable, from 2 to 20 inches (5-50 cm) in depth with OM of 1-3%.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Plant community consists of 55-75% grasses, 15% forbs, and 10-30% shrubs. Dense plant canopy (75-100%) and litter plus moderate to moderately rapid infiltration rates result in minimal to nonexistent runoff until soils are saturated. Basal cover is typically greater than 5% for this site and effectively reduces runoff on this site.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer exists.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional: mid-size, cool season bunchgrasses>tall, cool season bunchgrasses>perennial shrubs>rhizomatous grass-like>perennial forbs>cool season rhizomatous grasses
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Minimal decadence, typically associated with shrub component.
-
14. **Average percent litter cover (%) and depth (in):** Litter ranges from 1-25% of total canopy measurement with total litter (including beneath the plant canopy) from 75-100% expected. Herbaceous litter depth typically ranges from 10-25 mm. Woody litter can be up to a couple inches (4-6cm).
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** English: 3000-4300 lb/ac (3700 lb/ac average); Metric: 3360-4816 kg/ha (4144 kg/ha average).
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not**

invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Bare ground greater than 20% and presence of noxious weeds or Kentucky bluegrass are the most common indicators of a threshold being crossed. Baltic rush, inland sedge, herbaceous cinquefoil, pussytoes, and shrubby cinquefoil are common increasers. Kentucky bluegrass and Canada thistle are common invasive species.

17. **Perennial plant reproductive capability:** All species are capable of reproducing, except in drought years.
-