

Ecological site DX035X02B611 Clay Loam Upland 13-17" p.z. Gravelly (PIED, JUOS)

Accessed: 05/12/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 035X–Colorado Plateau

AZ CRA 35.6 - Colorado Plateau Pinyon-Juniper-Sagebrush

Elevations range from 5500 to 7000 feet and precipitation averages 13 to 17 inches per year. Vegetation includes pinyon, juniper, big sagebrush, cliffrose, Mormon tea, muttongrass, prairie junegrass, squirreltail, western wheatgrass, and blue grama. The soil temperature regime is mesic and the soil moisture regime is aridic ustic. This unit occurs within the Colorado Plateau Physiographic Province and is characterized by a sequence of flat to gently dipping sedimentary rocks eroded into plateaus, valleys and deep canyons. Sedimentary rock classes dominate the plateau with volcanic fields occurring for the most part near its margin.

Table 1. Dominant plant species

Tree	(1) Pinus edulis(2) Juniperus osteosperma
Shrub	(1) Artemisia tridentata ssp. wyomingensis(2) Purshia stansburiana
Herbaceous	(1) Poa fendleriana (2) Bouteloua gracilis

Legacy ID

F035XF611AZ

Physiographic features

This ecological site is located on summits, footslopes, shoulders and fan terraces of hills, mesas and plateaus. The soil is shallow to deep to any plant root restricting layer. The soil surface is gravelly to extremely gravelly loams and fine sandy loams. Slope of the ecological site ranges from 1 to 20 percent.

Landforms	(1) Hill(2) Mesa(3) Plateau
Flooding frequency	None
Ponding frequency	None
Elevation	5,700–6,600 ft
Slope	1–20%
Aspect	Aspect is not a significant factor

Table 2. Representative physiographic features

Climatic features

The climate of this land resource unit is semiarid with warm summers and cool winters. The mean annual precipitation ranges from 13 – 17 inches, but it is very erratic, often varying substantially from year to year. The majority of the precipitation comes from October through April. This precipitation comes as gentle rain or snow from frontal storms coming out of the Pacific Ocean. Snow is common from November through February. Generally no more than a few inches of snow accumulates, melting within a few days, but may last a week or more. The remaining precipitation comes from July through September as spotty, unreliable and sometimes violent thunderstorms. The moisture for this precipitation originates in the Gulf of Mexico (and the Pacific Ocean in the fall) and flows into the area on the north end of the Mexican monsoon. Late May through late June is generally a dry period. The mean annual air temperature ranges from 47 to 49 degrees Fahrenheit (F). The frost-free period (air temperature > 32 degrees F) ranges from 113 to 144 days (@ 50 percent probability). Strong winds are common, especially in the spring.

Table 3. Representative climatic features

Frost-free period (average)	144 days
Freeze-free period (average)	160 days
Precipitation total (average)	17 in

Influencing water features

Soil features

Soils are shallow to deep with surface textures of gravelly fine sandy loam, extremely gravelly loam, very gravelly loam, loam, cobbly silty clay loam, silty clay loam, very cobbly clay loam, and very cobbly silty clay loam. Subsoil textures are loam, clay loam, clay, gravelly clay, silty clay, gravelly clay loam, gravelly loam, very gravelly clay loam, very gravelly loam, stony clay and very cobbly clay. Parent materials are alluvium, colluvium and residuum from limestone, basalt and sandstone.

Typical taxonomic units include:

SSA-623 Shivwits Area MU's 21 Disterheff, 21 & 95 Natank, 69 Thunderbird, 72 Springerville, 69 & 70 Showlow, 72

Delenbaw;

SSA-625 Mohave county NE part MU's 57,58,59 & 60 Showlow;

SSA-699 Hualapai/Havasupai Area MU's 11 Disterheff, 27 Natank, 48 Toqui, 50 Tovar;

SSA-701 Grand Canyon Area MU's 4 Aridic haplustalfs, 8 Bilburc GrVL, 9 & 170 Bilburc, 9 Binsin, 22 Houserock family, 23 Toqui family, 35 Disterheff, 55 Kellypoint, 94, 125, 131 & 175 Toqui, 124 Toqui GrL.

Table 4. Representative soil features

Parent material	(1) Alluvium–limestone and sandstone(2) Colluvium–basalt
Surface texture	(1) Gravelly fine sandy loam(2) Extremely gravelly loam(3) Cobbly silty clay loam
Family particle size	(1) Clayey
Drainage class	Well drained
Permeability class	Slow to moderately slow
Soil depth	10–60 in
Surface fragment cover <=3"	0–85%
Surface fragment cover >3"	0–35%
Available water capacity (0-40in)	2.5–10 in
Calcium carbonate equivalent (0-40in)	15–45%
Electrical conductivity (0-40in)	0 mmhos/cm
Sodium adsorption ratio (0-40in)	0
Soil reaction (1:1 water) (0-40in)	6.1–8.4
Subsurface fragment volume <=3" (Depth not specified)	0–10%
Subsurface fragment volume >3" (Depth not specified)	0–5%

Ecological dynamics

The plant communities found on an ecological site are naturally variable. Composition and production will vary with yearly conditions, location, aspect, and the natural variability of the soils. The Historical Climax Plant Community represents the natural potential plant communities found on relict or relatively undisturbed sites. Other plant communities described here represent plant communities that are known to occur when the site is disturbed by factors such as fire, grazing, or drought.

Production data provided in this site description is standardized to air dry weight at the end of the summer growing season. The plant communities described in this site description are based on near normal rainfall years.

NRCS uses a Similarity Index to compare existing plant communities to the plant communities described here. Similarity index is determined by comparing the production and composition of a plant community to the production and composition of a plant community described in this site description. To determine Similarity index, compare the production (air dry weight) of each species to that shown in the plant community description. For each species, count no more than the maximum amount shown for the species, and for each group, count no more than the maximum amount shown for each group. Divide the resulting total by the total normal year production shown in the plant community description. If the rainfall has ben significantly above or below normal, use the total production shown for above or below normal years. If field data is not collected at the end of the summer growing season, then the field data must be corrected to the end of the year production before comparing it to the site description. The growth curve can be used as a guide for estimating production at the end of the summer growing season.

State and transition model



State 1 Historic Climax Plant Community

Community 1.1 Historic Climax Plant Community

The potential plant community consists of 50% tree canopy cover over an understory of grasses, forbs, shrubs and small trees. In the herbaceous/shrub stage,, grasses and some shrubs dominate the site. A few forbs and tree seedlings are present. In the immature stage, grasses and shrubs will dominate with medium sized trees; forbs are few. In the mature/climax stage, trees dominate the site; grasses and shrubs are in balance with tree competition and shade.

Figure 4. Plant community growth curve (percent production by month). AZ3506, 35.6 13-17" p.z. blue grama. Growth occurs mostly in summer and early fall during the rainy season..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	5	10	15	25	30	15	0	0	0

Figure 5. Plant community growth curve (percent production by month). AZ3561, 35.6 13-17" p.z. all sites. Growth begins in the spring and continues into the fall..

Jan	Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	1	5	16	17	15	15	15	11	5	0	0

Figure 6. Plant community growth curve (percent production by month). AZ3602, 35.6 13-17" p.z. muttongrass. Most growth occurs in early to mid spring, plants may be green in the fall. Seed set occurs by summer..

Jan	Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	10	20	30	15	5	5	10	5	0	0

Figure 7. Plant community growth curve (percent production by month). AZ3603, 35.6 13-17" p.z. Wyoming big sagebrush. Most growth occurs in the summer. Seed set occurs in the fall.

Jan	Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	5	15	20	20	20	10	10	0	0

Figure 8. Plant community growth curve (percent production by month). AZ3929, 35.6 13-17" p.z. turbinella oak. Growth begins in spring and continues through the summer. Seed set occurs in the fall..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	10	15	20	15	15	15	10	0	0

Figure 9. Plant community growth curve (percent production by month). AZ3930, 35.6 13-17" p.z. banana yucca. Most leaf growth occurs in late spring. Flowers in summer. Seed set occurs in late summer to fall..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	10	10	20	20	20	10	10	0	0

Additional community tables

 Table 5. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)	Diameter (In)	Basal Area (Square Ft/Acre)
Tree							
Utah juniper	JUOS	Juniperus osteosperma	Native	1–16	20–35	2–13	_
twoneedle pinyon	PIED	Pinus edulis	Native	4–27	20–35	3–14	_

Table 6. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)	
Grass/grass-like (Graminoids)						
blue grama	BOGR2	Bouteloua gracilis	Native	0.1–1	5–8	
squirreltail	ELELE	Elymus elymoides ssp. elymoides	Native	0.1–2	3–8	
Fendler's threeawn	ARPUF	Aristida purpurea var. fendleriana	Native	0.1–1.2	2–5	
muttongrass	POFE	Poa fendleriana	Native	0.1–1.5	2–4	
prairie Junegrass	KOMA	Koeleria macrantha	Native	0.1–1.5	1–3	
Shrub/Subshrub						
broom snakeweed	GUSA2	Gutierrezia sarothrae	Native	0.2–0.6	5–10	
Siler's penstemon	PELIS	Penstemon linarioides ssp. sileri	Native	0.2–2	2–10	
Wyoming big sagebrush	ARTRW8	Artemisia tridentata ssp. wyomingensis	Native	1–4	5–10	
Fremont's mahonia	MAFR3	Mahonia fremontii	Native	1–7	4–9	
pointleaf manzanita	ARPU5	Arctostaphylos pungens	Native	1–6	4–8	
Gambel oak	QUGA	Quercus gambelii	Native	1–18	2–7	
mormon tea	EPVI	Ephedra viridis	Native	0.3–3	2–5	
Stansbury cliffrose	PUST	Purshia stansburiana	Native	2–7	1–5	
Whipple cholla	CYWH	Cylindropuntia whipplei	Native	0.3–3	1–5	
New Mexico locust	RONE	Robinia neomexicana	Native	1–8	1–4	
skunkbush sumac	RHTR	Rhus trilobata	Native	0.5–4	1–3	
sulphur-flower buckwheat	ERUM	Eriogonum umbellatum	Native	0.1–0.6	1–3	
Sonoran scrub oak	QUTU2	Quercus turbinella	Native	1–6	1–3	
banana yucca	YUBA	Yucca baccata	Native	0.1–1.5	1–2	
Tree						
twoneedle pinyon	PIED	Pinus edulis	Native	4–25	10–30	
Utah juniper	JUOS	Juniperus osteosperma	Native	2–13	10–25	

Animal community

Trees are thick making the site only fairly suitable for grazing by cattle, sheep and horses in the summer and fall. Management considerations include grazing systems, proper grazing use and snow covering forage.

Springs are found on this site. Elk and deer use the site for winter cover. Food is adequte for native wildlife species. Competition with livestock can be a problem.`

Recreational uses

Recreational activities include hiking, hunting, wildlife observations and potography. Landscape has good aesthetic appeal.

Type locality

Location 1: Coconino County, AZ				
Township/Range/Section	T29 N. R7 W. S12			
General legal description	National Tank quad 4 miles SW of National Tank; Sec. 12, T. 29 N., R. 7 W.; Hualapai Indian Reservation, Arizona.			

Contributors

Harmon Hodgkinson Harmon S. Hodgkinson Larry D. Ellicott Stephen Cassady Steve Barker

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 17. Perennial plant reproductive capability: