

Ecological site DX035X03B625 Loamy Upland (PIED, JUOS) 13-17" p.z. (Provisional)

Accessed: 05/12/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 035X–Colorado Plateau

This ecological site occurs in Land Resource Unit 35.6 - Colorado Plateau Pinyon-Juniper-Sagebrush

The Land Resource Unit occurs within the Colorado Plateau Physiographic Province. Elevations range from 5800 to 7300 feet and precipitation averages 14 to 18 inches per year. Vegetation includes pinyon, juniper, big sagebrush, cliffrose, Mormon tea, muttongrass, prairie junegrass, squirreltail, western wheatgrass, and blue grama. The soil temperature regime is mesic and the soil moisture regime is aridic ustic. This unit occurs within the Colorado Plateau Physiographic Province and is characterized by a sequence of flat to gently dipping sedimentary rocks eroded into plateaus, valleys and deep canyons. Sedimentary rock classes dominate the plateau with volcanic fields occurring for the most part near its margin.

Ecological site concept

"ATTENTION: This ecological site meets the requirements for PROVISIONAL (if not more). A provisional ecological site is established after ecological site concepts are developed and an initial state-and-transition model is drafted. A provisional ecological site typically will include literature reviews, land use history information, legacy data (prior approved range site descriptions, forage suitability groups, woodland suitability groups, etc.,), and includes some soils data, and estimates for canopy and/or species composition by weight,. A provisional ecological site provides the conceptual framework of soil-site correlation for the development of the ESD. For more information about this

Associated sites

DX035X03B628	Sandy Loam Upland (JUOS, PIED) 13-17" p.z.
	Deep to very deep upland forest sites with coarse textured soils

Table 1. Dominant plant species

Tree	(1) Pinus edulis (2) Juniperus osteosperma
Shrub	(1) Artemisia tridentata ssp. wyomingensis(2) Gutierrezia sarothrae
Herbaceous	(1) Poa fendleriana (2) Bouteloua gracilis

Legacy ID

F035XF625AZ

Physiographic features

This site occurs mainly on summits, mesas, and plateaus derived from eolian deposits, slope alluvium and/or residuum from sandstone and shale. Slopes are generally 1 to 15%, but can occasionally range up to 25 percent on sideslopes and shoulders of hills and plateaus. Soils are moderately deep to deep to any plant root restrictive layer. The soil surface textures range from fine sandy loam to clay loam.

Landforms	(1) Mesa(2) Plateau(3) Hill
Flooding frequency	None
Ponding frequency	None
Elevation	5,800–7,300 ft
Slope	1–15%
Aspect	Aspect is not a significant factor

Table 2. Representative physiographic features

Climatic features

The climate of this land resource unit is semiarid with warm summers and cool winters. The mean annual precipitation ranges from 13-17 inches, but it is very erratic, often varying substantially from year to year. The majority of the precipitation comes from October through April. This precipitation comes as gentle rain or snow from frontal storms coming out of the Pacific Ocean. Snow is common from November through February. Generally no more then a few inches of snow accumulates, melting within a few days, but may last a week or more. The remaining precipitation comes from July through September as spotty, unreliable and sometimes violent thunderstorms. The moisture for this precipitation originates in the Gulf of Mexico (and the Pacific Ocean in the fall) and flows into the area on the north end of the Mexican monsoon. Late May through late June is generally a dry period. The mean annual air temperature ranges from 47 to 49 degrees Fahrenheit (F). The frost-free periods (air temperature > 32 degrees F) range from 113 to 144 days (@ 50 percent probability). Strong winds are common, especially in the spring.

Table 3. Representative climatic features

Frost-free period (average)	144 days
-----------------------------	----------

Freeze-free period (average)	160 days
Precipitation total (average)	17 in

Influencing water features

Soil features

The soils on this site are moderately deep to deep. Surface textures range from loam to clay loam. Parent materials are eolian deposits and/or residuum weathered from sandstone and shale. There may be up to 15% gravels on the surface. Some soils may have a gravelly substratum. Effervescence increases with depth. Depth to restrictive layers is 26 to 60 inches.

Typical taxonomic units on this site include:

SSA-701 Grand Canyon Area MU's 130 Tovar loam and 131 Tovar; SSA-692 McKinley County, NM MU's 310,320,350,354,360,555,575 Parkelei; 310,320,350,555 Bryway; 310,320,555 Evpark SSA-712 Canyon de Chelly NM MU 9 Evpark family; SSA-713 Chinle Area MU 25 Kachina & Evpark; SSA-715 Fort Defiance Area AZ/NM MU's 31 Evpark, 32 Evpark, 42 Nomrah, 75, 76, 77 & 86 Parkelei family.

Table 4.	Representative	soil	features
----------	----------------	------	----------

Parent material	(1) Eolian deposits–sandstone(2) Residuum–calcareous sandstone(3) Slope alluvium–sandstone and shale
Surface texture	(1) Loam(2) Fine sandy loam(3) Clay loam
Family particle size	(1) Loamy
Drainage class	Moderately well drained to well drained
Permeability class	Very slow to moderately rapid
Soil depth	26–60 in
Surface fragment cover <=3"	0–15%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	3.9–10.2 in
Calcium carbonate equivalent (0-40in)	0–15%
Electrical conductivity (0-40in)	0–2 mmhos/cm
Sodium adsorption ratio (0-40in)	0–2
Soil reaction (1:1 water) (0-40in)	6.6–8.3
Subsurface fragment volume <=3" (Depth not specified)	0–15%
Subsurface fragment volume >3" (Depth not specified)	0–5%

Ecological dynamics

An ecological site is not a precise assemblage of species for which the proportions are the same from place to place or from year to year. In all plant communities, variability is apparent in productivity and occurrence of individual species. Spatial boundaries of the communities; however, can be recognized by characteristic patterns of species composition, association, and community structure. The historic climax plant community for this ecological site has been described by sampling relict or relatively undisturbed sites and/or reviewing historic records. The historic climax plant community is the plant community that evolved over time with the soil forming process and long term changes in climatic conditions of the area. It is the plant community that was best adapted to the unique combination of environmental factors associated with the site.

Natural disturbances, such as drought, fire, grazing of native fauna, and insects, are inherent in the development and maintenance of these plant communities. The effects of these disturbances are part of the range of characteristics of the ecological site. Fluctuations in plant community structure and function caused by the effects of natural disturbances help establish the boundaries and characteristics of an ecological site. They are accounted for as part of the range of characteristics of the ecological site. Recognizable plant community phases are identified in the reference state of the ecological site. Some sites may have a small range of variation, while others have a large range. Some plant community phases may exist for long periods of time, while others may only occur for a couple of years after a disturbance.

Deterioration of the plant community, hydrology, or soil site stability on an ecological site can result in crossing a threshold or potentially irreversible boundary to another state, or equilibrium. This can occur as a result of the loss of soil surface through erosion, the loss of the stability of the site due to disturbances that cause active erosion on the site, increases in the amounts and/or patterns or runoff from rainstorms, changes in availability of surface and subsurface water, significant changes in plant structural and functional types, or the introduction of non-native species. When these thresholds are crossed, the potential of the ecological site to return to the historic climax plant community can be lost, or restoration will require significant inputs . There may be multiple states possible for an ecological site, determined by the type and or severity of disturbance.

The known states and transition pathways for this ecological site are described in the state and transition model. Within each state, there may be one or more known plant community phases. These community phases describe the different plant community that can be recognized and mapped across this ecological site. The state and transition model is intended to help land users recognize the current plant community on the ecological site, and the management options for improving the plant community to the desired plant community.

Plant production information in this site description is standardized to the annual production on an air-dry weight basis in near normal rainfall years.

State and transition model

35.6AZ Loamy Upland 13-17' p.z. (PIED, JUOS) (F035XH625AZ)



Figure 4. 35.6 Loamy Upaland 13-17"p.z. (PIED, JUOS)

State 1 Reference State

This is a forest site. Utah juniper and pinyon are the dominant overstory species. The understory is dominated by a grass/shrub mixture. When disturbance occurs the shrub and tree component can increase while grasses decrease. Non-native plant species may be present in minor amounts.

Community 1.1 Reference Community - Juniper Pinyon Forest Overstory

This forested site has an overstory canopy cover of 35% to 55% of pinyon and juniper species. The plant community understory is a grass and shrub mixture with small amounts of forbs and trees shorter than 4.5'. The understory plant community composition is comprised of approximately 45-55% grasses, 25-35% shrubs, 5-15% forbs and 5-15% trees (under 4.5 feet tall). Common grasses include blue grama, muttongrass, Indian ricegrass and squirreltail. Dominant shrubs include big sagebrush, cliffrose, snakeweed and Bigelow sage.

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	200	275	355
Shrub/Vine	110	165	225
Tree	30	55	85
Forb	35	55	85
Total	375	550	750

Community 1.2 Light Forest Overstory

This plant community is characterized by a reduced amount of overstory tree species. The crown canopy is less than 35%, ranging from 25-35%. The understory has more production than the reference plant community (1.1) with an increase of grasses and forbs and reduced shrubs. Common grasses include blue grama, muttongrass, Indian ricegrass, squirreltail and galleta. Dominant shrubs include big sagebrush, cliffrose, snakeweed and rabbitbrush.

Community 1.3 Dense Tree Overstory



Figure 6. Loamy Upland 13-17" - Community 1.3

This site has a denser overstory of tree species with a crown canopy greater than 55%. The understory has less production than the reference plant community (1.1) with an decrease of grasses, forbs and shrubs. The shrub component moves to become co-dominanat to dominant. Dominant shrubs include big sagebrush, broom snakeweed, Bigelow sage and cliffrose. Common grasses include muttongrass, Indian ricegrass, squirreltail and bluegrama.

Community 1.4 Reduced Tree Overstory-Grassland/Shrubland

This plant community is characterized by a reduced overstory of tree species. The crown canopy is usually less than 20%, ranging from 5-25%. The understory has more production than the reference plant community with an increase of grasses, forbs and shrubs. The understory plant community is dominated by grasses, then shrubs, forbs and trees (under 4.5 feet tall). Common grasses include muttongrass, blue grama, Indian ricegrass and squirreltail. Dominant shrubs include big sagebrush, snakeweed and Bigelow sagebrush.

Pathway 1.1b Community 1.1 to 1.2

Drought, insect infestation, and/or woodcutting

Pathway 1.1a Community 1.1 to 1.3

Fire exclusion/Lack of natural fire, favorable precipitation, unmanaged grazing.

Pathway 1.2a Community 1.2 to 1.1

Favorable precipitation, trees compete with shrubs/grass, fire exclusion.

Pathway 1.3a Community 1.3 to 1.1

Drought, insect infestation, fire

Pathway 1.3b Community 1.3 to 1.4

Drought in combination with severe disturbance such as major insect infestation and/or intense fire.

Pathway 1.4a Community 1.4 to 1.2

Favorable precipitation, seed source for tree regeneration

Additional community tables

 Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)			
Grass	Grass/Grasslike							
1	Dominant Grasses			138–275				
	muttongrass	POFE	Poa fendleriana	55–138	_			
	blue grama	BOGR2	Bouteloua gracilis	28–110	_			
	Indian ricegrass	ACHY	Achnatherum hymenoides	28–83	_			
2	Miscellaneous Grasses			28–83				
	western wheatgrass	PASM	Pascopyrum smithii	28–55	-			
	squirreltail	ELEL5	Elymus elymoides	28–55	_			
	prairie Junegrass	KOMA	Koeleria macrantha	2–28	-			
	Fendler's threeawn	ARPUF	Aristida purpurea var. fendleriana	6–28	-			
	James' galleta	PLJA	Pleuraphis jamesii	6–28	-			
	Sandberg bluegrass	POSE	Poa secunda	6–28	_			
	Grass, annual	2GA	Grass, annual	0–17	-			
	Grass, perennial	2GP	Grass, perennial	0–17	_			
	sand dropseed	SPCR	Sporobolus cryptandrus	0–11	-			
Forb								
3	Forbs			28–85				
	Forb, annual	2FA	Forb, annual	0–11	-			
	Forb, perennial	2FP	Forb, perennial	0–11	_			
	ragwort	SENEC	Senecio	0–11	_			
	Carruth's sagewort	ARCA14	Artemisia carruthii	n_11				

	ounding ougonon	/		v		
	sandwort	ARENA	Arenaria	0–11	-	
	Wright's bird's beak	COWR2	Cordylanthus wrightii	0–11	_	
	redroot buckwheat	ERRA3	Eriogonum racemosum	0–11	-	
	pingue rubberweed	HYRI	Hymenoxys richardsonii	0–11	_	
	tansyaster	MACHA	Machaeranthera	0–11	_	
	dwarf lousewort	PECE	Pedicularis centranthera	0–6	_	
	beardtongue	PENST	Penstemon	0–6	_	
	dense mountain phlox	PHAUD	Phlox austromontana ssp. densa	0–6	-	
	longleaf phlox	PHLO2	Phlox longifolia	0–6	_	
	flatspine stickseed	LAOC3	Lappula occidentalis	0–6	_	
	Wright's deervetch	LOWR	Lotus wrightii	0–6	-	
	cryptantha	CRYPT	Cryptantha	0–6	-	
	western tansymustard	DEPI	Descurainia pinnata	0–6	-	
	wedgeleaf draba	DRCUC	Draba cuneifolia var. cuneifolia	0–6	-	
	fleabane	ERIGE2	Erigeron	0–6	-	
	milkvetch	ASTRA	Astragalus	0–6	-	
	curveseed butterwort	CETE5	Ceratocephala testiculata	0–6	_	
	Esteve's pincushion	CHST	Chaenactis stevioides	0–6	-	
	scarlet globemallow	SPCO	Sphaeralcea coccinea	0–6	_	
	onion	ALLIU	Allium	0–6	_	
	Townsend daisy	TOWNS	Townsendia	0–1	-	
	Shrub/Vine					
Shrub	/Vine					
Shrub 4	/Vine Dominant Shrubs			83–193		
Shrub 4	/Vine Dominant Shrubs Wyoming big sagebrush	ARTRW8	Artemisia tridentata ssp. wyomingensis	83–193 55–110		
Shrub 4	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed	ARTRW8 GUSA2	Artemisia tridentata ssp. wyomingensis Gutierrezia sarothrae	83–193 55–110 28–55		
Shrub 4 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs	ARTRW8 GUSA2	Artemisia tridentata ssp. wyomingensis Gutierrezia sarothrae	83–193 55–110 28–55 28–83	-	
Shrub 4 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage	ARTRW8 GUSA2 ARBI3	Artemisia tridentata ssp. wyomingensis Gutierrezia sarothrae Artemisia bigelovii	83–193 55–110 28–55 28–83 0–55	-	
Shrub 4 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose	ARTRW8 GUSA2 ARBI3 PUST	Artemisia tridentata ssp. wyomingensis Gutierrezia sarothrae Artemisia bigelovii Purshia stansburiana	83–193 55–110 28–55 28–83 0–55 0–28	-	
55	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush	ARTRW8 GUSA2 ARBI3 PUST PUTR2	Artemisia tridentata ssp. wyomingensis Gutierrezia sarothrae Artemisia bigelovii Purshia stansburiana Purshia tridentata	83–193 55–110 28–55 28–83 0–55 0–28 0–28	-	
5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m)	ARTRW8 GUSA2 ARBI3 PUST PUTR2 2SHRUB	Artemisia tridentata ssp. wyomingensis Gutierrezia sarothrae Artemisia bigelovii Purshia stansburiana Purshia tridentata Shrub (>.5m)	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–17	-	
5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m) black sagebrush	ARTRW8 GUSA2 ARBI3 PUST PUTR2 2SHRUB ARNO4	Artemisia tridentata ssp. wyomingensis Gutierrezia sarothrae Artemisia bigelovii Purshia stansburiana Purshia tridentata Shrub (>.5m) Artemisia nova	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–17 0–11	-	
5 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m) black sagebrush alderleaf mountain mahogany	ARTRW8 GUSA2 ARBI3 PUST PUTR2 2SHRUB ARNO4 CEMO2	Artemisia tridentata ssp. wyomingensis Gutierrezia sarothrae Artemisia bigelovii Purshia stansburiana Purshia tridentata Shrub (>.5m) Artemisia nova Cercocarpus montanus	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–17 0–11 0–11	-	
5 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m) black sagebrush alderleaf mountain mahogany Cutler's jointfir	ARTRW8 GUSA2 ARBI3 PUST PUTR2 2SHRUB ARNO4 CEMO2 EPCU	Artemisia tridentata ssp. wyomingensis Gutierrezia sarothrae Artemisia bigelovii Purshia stansburiana Purshia tridentata Shrub (>.5m) Artemisia nova Cercocarpus montanus Ephedra cutleri	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–17 0–11 0–11		
5 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m) black sagebrush alderleaf mountain mahogany Cutler's jointfir mormon tea	ARTRW8 GUSA2 ARBI3 PUST PUTR2 2SHRUB ARNO4 CEMO2 EPCU EPVI	Artemisia tridentata ssp. wyomingensisGutierrezia sarothraeGutierrezia sarothraeArtemisia bigeloviiPurshia stansburianaPurshia tridentataShrub (>.5m)Artemisia novaCercocarpus montanusEphedra cutleriEphedra viridis	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–17 0–11 0–11 0–11		
5 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m) black sagebrush alderleaf mountain mahogany Cutler's jointfir mormon tea pricklypear	ARTRW8 GUSA2 ARBI3 PUST PUTR2 2SHRUB ARNO4 CEMO2 EPCU EPVI OPUNT	Artemisia tridentata ssp. wyomingensis Gutierrezia sarothrae Artemisia bigelovii Purshia stansburiana Purshia tridentata Shrub (>.5m) Artemisia nova Cercocarpus montanus Ephedra cutleri Ephedra viridis Opuntia	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–17 0–11 0–11 0–11 0–11		
5 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m) black sagebrush alderleaf mountain mahogany Cutler's jointfir mormon tea pricklypear banana yucca	ARTRW8 GUSA2 ARBI3 PUST PUTR2 2SHRUB ARNO4 CEMO2 EPCU EPVI OPUNT YUBA	Artemisia tridentata ssp. wyomingensisGutierrezia sarothraeGutierrezia sarothraeArtemisia bigeloviiPurshia stansburianaPurshia tridentataShrub (>.5m)Artemisia novaCercocarpus montanusEphedra cutleriEphedra viridisOpuntiaYucca baccata	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–17 0–11 0–11 0–11 0–11 0–11		
5 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m) black sagebrush alderleaf mountain mahogany Cutler's jointfir mormon tea pricklypear banana yucca longflower rabbitbrush	ARTRW8 GUSA2 ARBI3 PUST PUTR2 2SHRUB ARNO4 CEMO2 EPCU EPVI OPUNT YUBA CHDE2	Artemisia tridentata ssp. wyomingensisGutierrezia sarothraeGutierrezia sarothraeArtemisia bigeloviiPurshia stansburianaPurshia tridentataShrub (>.5m)Artemisia novaCercocarpus montanusEphedra cutleriEphedra viridisOpuntiaYucca baccataChrysothamnus depressus	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–17 0–11 0–11 0–11 0–11 0–11		
5 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m) black sagebrush alderleaf mountain mahogany Cutler's jointfir mormon tea pricklypear banana yucca longflower rabbitbrush hedgehog cactus	ARTRW8 GUSA2 ARBI3 PUST PUTR2 2SHRUB ARNO4 CEMO2 EPCU EPVI OPUNT YUBA CHDE2 ECHIN3	Artemisia tridentata ssp. wyomingensisGutierrezia sarothraeGutierrezia sarothraeArtemisia bigeloviiPurshia stansburianaPurshia tridentataShrub (>.5m)Artemisia novaCercocarpus montanusEphedra cutleriEphedra viridisOpuntiaYucca baccataChrysothamnus depressusEchinocereus	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–28 0–17 0–11 0–11 0–11 0–11 0–11 0–11		
Shrub 4 5 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m) black sagebrush alderleaf mountain mahogany Cutler's jointfir mormon tea pricklypear banana yucca longflower rabbitbrush hedgehog cactus	ARTRW8 GUSA2 PUST PUTR2 2SHRUB ARNO4 CEMO2 EPCU EPVI OPUNT YUBA CHDE2 ECHIN3	Artemisia tridentata ssp. wyomingensisGutierrezia sarothraeGutierrezia sarothraeArtemisia bigeloviiPurshia stansburianaPurshia tridentataShrub (>.5m)Artemisia novaCercocarpus montanusEphedra cutleriEphedra viridisOpuntiaYucca baccataChrysothamnus depressusEchinocereus	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–28 0–17 0–11 0–11 0–11 0–11 0–11 0–11		
Shrub 4 5 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m) black sagebrush alderleaf mountain mahogany Cutler's jointfir mormon tea pricklypear banana yucca longflower rabbitbrush hedgehog cactus Trees (<4.5')	ARTRW8 GUSA2 ARBI3 PUST PUTR2 2SHRUB ARNO4 CEMO2 EPCU EPVI OPUNT YUBA CHDE2 ECHIN3	Artemisia tridentata ssp. wyomingensisGutierrezia sarothraeGutierrezia sarothraeArtemisia bigeloviiPurshia stansburianaPurshia tridentataShrub (>.5m)Artemisia novaCercocarpus montanusEphedra cutleriEphedra viridisOpuntiaYucca baccataChrysothamnus depressusEchinocereus	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–28 0–17 0–11 0–11 0–11 0–11 0–11 0–11 0–11		
Shrub 4 5 5	/Vine Dominant Shrubs Wyoming big sagebrush broom snakeweed Miscellaneous Shrubs Bigelow sage Stansbury cliffrose antelope bitterbrush Shrub (>.5m) black sagebrush alderleaf mountain mahogany Cutler's jointfir mormon tea pricklypear banana yucca longflower rabbitbrush hedgehog cactus Trees (<4.5') twoneedle pinyon	ARTRW8 GUSA2 ARBI3 PUST PUTR2 2SHRUB ARNO4 CEMO2 EPCU EPVI OPUNT YUBA CHDE2 ECHIN3 PIED	Artemisia tridentata ssp. wyomingensisGutierrezia sarothraeGutierrezia sarothraeArtemisia bigeloviiPurshia stansburianaPurshia tridentataShrub (>.5m)Artemisia novaCercocarpus montanusEphedra cutleriEphedra viridisOpuntiaYucca baccataChrysothamnus depressusEchinocereusPinus edulis	83–193 55–110 28–55 28–83 0–55 0–28 0–28 0–28 0–17 0–11 0–11 0–11 0–11 0–11 0–11 0–11		

Gambel oakQUGAQuercus gambelii0–11	_
------------------------------------	---

Inventory data references

Updates and revisions for this ESD were conducted as part of a 2007-2012 Interagency Technical Assistance Agreement between the Bureau of Indian Affairs–Navajo Region and the NRCS-Arizona.

Other references

Updates and revisions for this ESD were conducted as part of a 2007-2012 Interagency Technical Assistance Agreement between the Bureau of Indian Affairs–Navajo Region and the NRCS-Arizona.

Contributors

Dan Carroll Karlynn Huling Kenneth Gishi Kevin Williams Larry D. Ellicott Peter Lefebvre

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values):
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if

their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability: