

Ecological site R035XB035NM Sandy Upland 6-10"

Accessed: 05/12/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) Ephedra ×arenicola
Herbaceous	(1) Achnatherum hymenoides

Physiographic features

This site occurs as treads of high stream terraces, fan terraces, and stable dunes of undulating plateaus and structural benches. It does not benefit from run-in moisture from adjacent areas, nor does it suffer from excessive loss from runoff. It occurs on all exposures. Slopes range from 0 to 8 percent. Elevations range from 4,700 to 5,900 feet.

Landforms	(1) Stream terrace(2) Dune(3) Cuesta
Flooding frequency	None
Ponding frequency	None

Elevation	1,433–1,798 m
Slope	0–8%
Water table depth	152 cm
Aspect	Aspect is not a significant factor

Climatic features

Mean annual precipitation varies from 7 to 10 inches. About 60 percent of this moisture comes as rain from April through October. May and June are the driest months. Most of the moisture from November through March comes as snow. Winds of high velocity during late winter and early spring are common.

Mean temperature for the hottest month, July, is about 83 degrees F. Mean temperature for the coldest month, January, is about 27 degrees F. Extreme temperatures of 104 degrees F and –17 degrees F have been recorded. The frost-free period ranges from 140 to 160 days.

The cool-season plants start growth in March and end with plant maturity and seed dissemination about mid-June. During June, July, August, and September, the warm-season plants make optimum growth taking advantage of the warm temperature and moisture from tropical air out of the Gulf of Mexico. About 40 percent of the total precipitation is received during these summer months. The other 60 percent received during the fall-winter-spring months influences cool-season plants.

Table 3. Representative climatic features

Frost-free period (average)	160 days
Freeze-free period (average)	165 days
Precipitation total (average)	254 mm

Influencing water features

This site is not influenced by water from wetland or stream.

Soil features

The soils are deep to very deep and somewhat excessively to excessively well drained. They are formed in eolium, residuum, and alluvium derived from sandstone. Surface textures include sand, fine sand, loamy fine sand, and loamy sand. The subsoil has textures of loamy fine sand, channery fine sand, loamy sand, and fine sand. Permeability is moderately rapid to rapid. Available water holding capacity to a depth of 40 inches is very low to low. Runoff is negligible to low, and the hazard of water erosion is slight. The hazard of soil blowing is severe. The soils are neutral to strongly alkaline (pH 6.6-9.0), nonsodic to slightly sodic (SAR 0-10), and nonsaline to slightly saline (EC 0-4).

Characteristic taxonomic units are: Shiprock SSA: 115-Denazar-Farb fine sands (Denazar part) 120-Nageezi-Denazar Association (Denazar part) 133-Razito sand 145-Razito-Huerfano Complex (Razito part) 275-Razito loamy sand Other soils included are:

Table 4. Representative soil features

Surface texture	(1) Sand
	(2) Fine sand
	(3) Loamy fine sand

Family particle size	(1) Sandy
Drainage class	Somewhat excessively drained to excessively drained
Permeability class	Moderately rapid to rapid
Soil depth	102–157 cm
Surface fragment cover <=3"	0–10%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	7.62–10.16 cm
Calcium carbonate equivalent (0-101.6cm)	0–30%
Electrical conductivity (0-101.6cm)	0–4 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–10
Soil reaction (1:1 water) (0-101.6cm)	6.6–9
Subsurface fragment volume <=3" (Depth not specified)	0–15%
Subsurface fragment volume >3" (Depth not specified)	0–30%

Ecological dynamics

This site has a plant community made up primarily of grasses with some shrubs and minor amounts of forbs. In the historic climax plant community, there is a mixture of both cool- and warm-season grasses. Plant species most likely to invade or increase on this site when it deteriorates are annual bursage (flatspine bur ragweed), annual mustard spp., sandhill muhly, Greene's rabbitbrush, sandy jointfir, (sand Mormon tea), and Cutler's jointfir (Cutler Mormon tea). Continuous livestock grazing during winter and spring decreases the cool-season grasses, which are replaced by lower forage value species.

State and transition model

Ecosystem states



State 1 submodel, plant communities

1.1. Historic Climax Plant Community

State 1 Historic Climax Plant Community

Community 1.1 Historic Climax Plant Community This site has a plant community made up primarily of grasses with some shrubs and minor amounts of forbs. In the historic climax plant community, there is a mixture of both cool- and warm-season grasses. Plant species most likely to invade or increase on this site when it deteriorates are annual bursage (flatspine bur ragweed), annual mustard spp., sandhill muhly, Greene's rabbitbrush, sandy jointfir, (sand Mormon tea), and Cutler's jointfir (Cutler Mormon tea). Continuous livestock grazing during winter and spring decreases the cool-season grasses, which are replaced by lower forage value species.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	196	314	392
Shrub/Vine	71	112	140
Forb	13	22	28
Total	280	448	560

Figure 5. Plant community growth curve (percent production by month). NM0383, R035XB035NM-Sandy Upland 6 to 10 inch-Reference Community. R035XB035NM-Sandy Upland 6 to 10 inch-Reference Community.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
6	6	7	6	6	5	11	14	12	12	8	7

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)		
Grass	Grass/Grasslike						
1	Indian ricegrass		90–135				
	Indian ricegrass	ACHY	Achnatherum hymenoides	90–135	_		
2	galleta			22–45			
	James' galleta	PLJA	Pleuraphis jamesii	22–45	_		
3	sand dropseed	-	·	4–22			
	sand dropseed	SPCR	Sporobolus cryptandrus	4–22	_		
4	spike dropseed			0–13			
	spike dropseed	SPCO4	Sporobolus contractus	0–13	_		
5	mesa dropseed			4–22			
	mesa dropseed	SPFL2	Sporobolus flexuosus	4–22	_		
6	giant dropseed			0–9			
	giant dropseed	SPGI	Sporobolus giganteus	0–9	_		
7	blue grama	-	·	4–22			
	blue grama	BOGR2	Bouteloua gracilis	4–22	_		
8	Fendler's threeawn	-	·	0-4			
	Fendler's threeawn	ARPUF	Aristida purpurea var. fendleriana	0-4	_		
9	sandhill muhly	-	·	4–22			
	sandhill muhly	MUPU2	Muhlenbergia pungens	4–22	_		
10	needle and thread	-	·	0–9			
	needle and thread	HECO26	Hesperostipa comata	0–9	_		
11	other perennial grass	ses		0–13			
	Grass, perennial	2GP	Grass, perennial	0–13	_		

	1	1			
Forb	-				
12	globemallow			0-4	
	globemallow	SPHAE	Sphaeralcea	0-4	_
13	rose heath (smallflo	wer aster)		0-4	
	rose heath	CHER2	Chaetopappa ericoides	0-4	_
14	other perennial forb	S		0–9	
	Forb, perennial	2FP	Forb, perennial	0–9	_
15	annual forbs	•	•	0–9	
	Forb, annual	2FA	Forb, annual	0–9	_
Shru	b/Vine	•	•		
16	sandy jointfir (sand Mormon tea)			22–45	
17	Cutler's jointfir (Cutl	er Mormo	n tea)	0–9	
	Cutler's jointfir	EPCU	Ephedra cutleri	0–9	_
18	broom snakeweed		·	0–22	
	broom snakeweed	GUSA2	Gutierrezia sarothrae	0–22	_
19	Greene's rabbitbrus	h	·	0–9	
	Greene's rabbitbrush	CHGR6	Chrysothamnus greenei	0–9	_
20	plains pricklypear	•	•	0-4	
	plains pricklypear	OPPO	Opuntia polyacantha	0-4	_
21	fourwing saltbush	•	•	0–9	
	fourwing saltbush	ATCA2	Atriplex canescens	0–9	_
22	winterfat			0-4	
	winterfat	KRLA2	Krascheninnikovia lanata	0-4	_
23	narrowleaf yucca	-		0-4	
	narrowleaf yucca	YUAN2	Yucca angustissima	0-4	_
24	other shrubs			0–9	

Animal community

Fair to good plant diversity makes this site suitable for grassland wildlife species.

Hydrological functions

This site normally receives about 7-10 inches of annual precipitation. Most summer rainfall occurs as brief, sometimes heavy, thunderstorms. Slopes range from 0-8%. The soils are deep to very deep and somewhat excessively to excessively well drained. Surface and subsurface horizons are coarse-textured. Permeability is moderately rapid to rapid. Available water holding capacity is very low to low. Runoff is negligible to low, and the hazard of water erosion is slight.

Recreational uses

Hunting, horseback riding, and wildlife observation are occasional recreation activities on this site. The grassy aspect of this site provides aesthetic appeal.

Wood products

This site has no significant value for wood products.

Other products

Grazing: This site is suitable for yearlong grazing by all classes of livestock. Grazing systems adapt well to this site and should be used. Soils on this site have a high wind erosion hazard; when the site is deteriorated, erosion occurs on overgrazed areas, roads, cattle trails, and high-use areas.

Inventory data references

The potential historic climax plant community has been determined by study of range relict areas, or areas protected from excessive grazing. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures and historical accounts have also been used.

Type locality

Location 1: San Juan County, NM			
Township/Range/Section	T23N R15W S17		
General legal description	Between Hunter Wash and Chaco River – Section 17, Township 23N, Range 15W – Navajo Reservation, NM.		

Contributors

David Trujillo John Tunberg Michael Carpinelli Unknown

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:

- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values):
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):

- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 17. Perennial plant reproductive capability: