

Ecological site R035XH823AZ Sandstone Upland 17-25" p.z.

Accessed: 05/12/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 035X–Colorado Plateau

This ecological site occurs in Common Resource Area 35.8 - the Colorado Plateau Ponderosa Pine Forests

The Common Resource Area occurs within the Colorado Plateau Physiographic Province. Elevations range from 6800 to 8500 feet and precipitation averages 17 to 25 inches per year. Vegetation includes ponderosa pine, white fir, aspen, pinyon, juniper, Gambel oak, big sagebrush, ceanothus, blue elderberry, muttongrass, upland sedge, and big wildrye, mountain muhly, Arizona fescue, pine dropseed, and blue grama. The soil temperature regime ranges from mesic to frigid and the soil moisture regime is typic ustic. This unit occurs within the Colorado Plateau Physiographic Province and is characterized by a sequence of flat to gently dipping sedimentary rocks eroded into plateaus, valleys and deep canyons. Sedimentary rock classes dominate the plateau with volcanic fields occurring for the most part near its margin.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Koeleria macrantha</i> (2) <i>Bouteloua gracilis</i>

Physiographic features

This site occurs on structural benches and knolls of undulating plateaus. The soils are shallow (<20") over sandstone bedrock. Surface textures are loamy. Slopes are less than 15 percent. This site does not benefit significantly from run-on moisture.

Table 2. Representative physiographic features

Landforms	(1) Structural bench (2) Knoll (3) Plateau
Flooding frequency	None
Ponding frequency	None
Elevation	7,800–8,500 ft
Slope	0–15%
Aspect	Aspect is not a significant factor

Climatic features

Winter-Summer moisture ratios are typically 70:30 on the west side of this common resource area and shift to 60:40 on the east side. Late spring is usually the driest period and early fall moisture can be sporadic. Summer rains fall from June through September; moisture originates in the Gulf of Mexico and creates convective, usually brief, intense thunderstorms. Cool season moisture from October through May tends to be frontal; it originates in the Pacific and the Gulf of California and falls in widespread storms with longer duration and lower intensity. Precipitation generally comes as snow from October into April. Snowpack can persist for 3-4 months, although it may disappear in exposed areas during prolonged dry weather. Summer daytime temperatures are typically 80-90 F but can exceed 95 F. Winter temperatures around 0 F are common and can reach -25 F.

Table 3. Representative climatic features

Frost-free period (average)	100 days
Freeze-free period (average)	130 days
Precipitation total (average)	25 in

Influencing water features

The soil moisture on this ecological site comes from precipitation. The site does not benefit significantly from run-on moisture. Shallow bedrock areas will concentrate water in deeper soil pockets, where most of the vegetation production occurs. Shrubs will also harvest water from deep cracks in the bedrock. Because of the shallow soils, rapid snowmelt and larger rainfall events may not be entirely captured by the site. This site contributes runoff to other ecological sites.

Soil features

The soils are well drained, moderately permeable soils which are very shallow to sandstone bedrock. Surface textures are loamy. Subsurface textures are sandy loam to loam. There is not an argillic layer above the sandstone bedrock.

Typical soil series for this site occurs:

SSA-715 Ft. Defiance Area AZ/NM MU 17 Berland family;

SSA-717 Shiprock Area AZ/NM MU 702 Berland.

Table 4. Representative soil features

Parent material	(1) Alluvium—sandstone
Surface texture	(1) Loam
Family particle size	(1) Sandy
Drainage class	Moderately well drained to well drained
Permeability class	Moderately slow to moderate
Soil depth	5–15 in
Surface fragment cover ≤3"	0–5%
Surface fragment cover >3"	0–15%
Electrical conductivity (0–40in)	0–8 mmhos/cm
Soil reaction (1:1 water) (0–40in)	5.6–6.5
Subsurface fragment volume ≤3" (Depth not specified)	0–5%

Ecological dynamics

An ecological site is not a precise assemblage of species for which the proportions are the same from place to place or from year to year. In all plant communities, variability is apparent in productivity and occurrence of individual species. Spatial boundaries of the communities; however, can be recognized by characteristic patterns of species composition, association, and community structure. The historic climax plant community for this ecological site has been described by sampling relict or relatively undisturbed sites and/or reviewing historic records. The historic climax plant community is the plant community that evolved over time with the soil forming process and long term changes in climatic conditions of the area. It is the plant community that was best adapted to the unique combination of environmental factors associated with the site.

Natural disturbances, such as drought, fire, grazing of native fauna, and insects, are inherent in the development and maintenance of these plant communities. The effects of these disturbances are part of the range of characteristics of the ecological site. Fluctuations in plant community structure and function caused by the effects of natural disturbances help establish the boundaries and characteristics of an ecological site. They are accounted for as part of the range of characteristics of the ecological site. Recognizable plant community phases are identified in the reference state of the ecological site. Some sites may have a small range of variation, while others have a large range. Some plant community phases may exist for long periods of time, while others may only occur for a couple of years after a disturbance.

Deterioration of the plant community, hydrology, or soil site stability on an ecological site can result in crossing a threshold or potentially irreversible boundary to another state, or equilibrium. This can occur as a result of the loss of soil surface through erosion, the loss of the stability of the site due to disturbances that cause active erosion on the site, increases in the amounts and/or patterns or runoff from rainstorms, changes in availability of surface and subsurface water, significant changes in plant structural and functional types, or the introduction of non-native species. When these thresholds are crossed, the potential of the ecological site to return to the historic climax plant community can be lost, or restoration will require significant inputs. There may be multiple states possible for an ecological site, determined by the type and or severity of disturbance.

The known states and transition pathways for this ecological site are described in the state and transition model. Within each state, there may be one or more known plant community phases. These community phases describe the different plant community that can be recognized and mapped across this ecological site. The state and transition model is intended to help land users recognize the current plant community on the ecological site, and the management options for improving the plant community to the desired plant community.

Plant production information in this site description is standardized to the annual production on an air-dry weight basis in near normal rainfall years.

State and transition model

35.8 Shallow Loamy 17-25"p.z.

(R035XH823AZ)




Figure 4. STM - R035XH823AZ

State 1

Reference State

This is a grassland site with some forbs. Non-native annual grass (cheatgrass) may or may not be found on this site.

Community 1.1

Blue grama - Kentucky bluegrass and Pussytoes

The dominant aspect of this site is grasses, grasslikes and forbs. Blue grama is the dominant grass species along with prairie Junegrass and carex. The dominant forb is pussytoes followed by hairy false goldenaster and silver cinquefoil.

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	240	440	640
Forb	80	160	240
Total	320	600	880

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	0%
Grass/grasslike foliar cover	15-25%
Forb foliar cover	5-10%
Non-vascular plants	0-10%
Biological crusts	0%
Litter	20-40%

Surface fragments >0.25" and <=3"	0-15%
Surface fragments >3"	0-10%
Bedrock	0-10%
Water	0%
Bare ground	20-40%

Figure 6. Plant community growth curve (percent production by month). AZ3581, 35.8 17-25" p.z. all sites. Growth begins in the spring, most growth occurs during the summer rainy season..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	4	10	24	21	23	13	5	0	0

Figure 7. Plant community growth curve (percent production by month). AZ3902, 35.8 17-25" p.z. Arizona fescue. Growth begins in the late spring and extends through the summer rainy season..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	0	5	20	30	30	10	5	0	0

Community 1.2

Kentucky bluegrass/ Perennial and Annual Forbs



Figure 8. 1.2 Plant Community

This plant community has a perennial grass and forb plant community. There are moderate amounts of annual and perennial forbs in this community.

Pathway 1.1a

Community 1.1 to 1.2

Unmanaged continuous season-long grazing, drought

Pathway 1.2a

Community 1.2 to 1.1

Prescribed grazing with rest or no grazing.

Additional community tables

Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass/Grasslike					
1	Grasses			240–640	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	80–120	–
	mountain muhly	MUMO	<i>Muhlenbergia montana</i>	0–120	–
	needleleaf sedge	CADU6	<i>Carex duriuscula</i>	40–80	–
	Arizona fescue	FEAR2	<i>Festuca arizonica</i>	16–80	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	40–80	–
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–40	–
	squirreltail	ELELE	<i>Elymus elymoides</i> ssp. <i>elymoides</i>	0–24	–
Forb					
2	Forbs			80–240	
	rosy pussytoes	ANRO2	<i>Antennaria rosea</i>	40–80	–
	tarragon	ARDR4	<i>Artemisia dracuncululus</i>	0–40	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	0–40	–
	milkvetch	ASTRA	<i>Astragalus</i>	0–40	–
	geranium	GERAN	<i>Geranium</i>	0–40	–
	hairy false goldenaster	HEVI4	<i>Heterotheca villosa</i>	0–40	–
	silver cinquefoil	POAR8	<i>Potentilla argentea</i>	24–40	–
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–40	–
	common yarrow	ACMI2	<i>Achillea millefolium</i>	0–40	–
	Forb, annual	2FA	<i>Forb, annual</i>	0–20	–

Animal community

Site is favorable for grazing throughout most of the year except when snow cover restricts availability of forage. Planned grazing systems adapt well to use on this site.

The potential plant community produced by this site provides food for those species of wildlife that utilize grass as a major portion of their diet.

Recreational uses

Winters are cold, however, relatively mild summer months are attractive to recreationists.

Activities include hunting, cross-country riding, photography, hiking, rock collecting, and wildlife observation

Type locality

Location 1: San Juan County, NM	
Township/Range/Section	T22N R20W S13
General legal description	Approximately 7 miles southwest of Toadlena, Navajo Indian Reservation, NM

Other references

Updates and revisions for this ESD were conducted as part of a 2007-2012 Interagency Technical Assistance Agreement between the Bureau of Indian Affairs–Navajo Region and the NRCS-Arizona.

Contributors

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Kenneth Gishi
Contact for lead author	NRCS State Rangeland Management Specialist, Arizona State Office, Phoenix, AZ
Date	08/24/2012
Approved by	Steve Barker
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:** None expected, except for steeper slopes where an occasional rills may occur and cover less than 5% of the area.

- 2. Presence of water flow patterns:** Uncommon due to moderate vegetation and rock cover. Some flow patterns may be evident follow large storm events, but should be very short and not connected.

- 3. Number and height of erosional pedestals or terracettes:** None expected.

- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 20-40 percent bare ground.

- 5. Number of gullies and erosion associated with gullies:** None.

- 6. Extent of wind scoured, blowouts and/or depositional areas:** None.

- 7. Amount of litter movement (describe size and distance expected to travel):** Fine litter movement will be short and most will remain in place. Most litter movement will be transported by water.

- 8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of**

values): The expected soil stability rating under plant canopies is 4 to 5 and a rating in the interspaces of 3 to 4. The surface textures loams. When well vegetated and with adequate litter cover these soils have a moderate resistance to erosion.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface thickness ranges from 2-4". Soil structure is typically moderate thick platy structure parting to moderate medium granular.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** This site is characterized by a dominance of perennial grasses with forbs. The distribution of herbaceous and rock fragment cover helps minimizes erosion and aids in reducing surface flow to allow for infiltration.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. Some soils may have subsurface prismatic horizon structure with 6 inches of the surface. These are not compacted layers but may difficult to excavate and mistaken for a compacted layer
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Cool season grasses > warm season grasses >
- Sub-dominant: Perennial forbs > Annual forbs
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** All plants functional groups are adapted to survival in all but the most severe droughts. Severe winter droughts affect shrubs the most. Severe summer droughts affect grasses the most.
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Average annual production on this site is expected to be 500 to 700 lbs/ac. in a year of average annual precipitation.
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not**

invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Kentuckly bluegrass is a naturalized plant that can increase and dominate. Other plants that can increase with disturbance are Larkspur, lupine, rubberweed, snakeweed, goldenaster, burr buttercup, knotweed, leafy spurge and cheatgrass.

17. **Perennial plant reproductive capability:** All plants native to this site are adapted and are capable of producing seeds, stolons and rhizomes in all but the most severe drought.
-