

Ecological site R040XC320AZ Sandy Loam Upland 3"-7" p.z.

Accessed: 05/12/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Womack, Dan Robinett, Emilio Carrillo.
Contact for lead author	NRCS Tucson Area Office
Date	03/07/2005
Approved by	S. Cassady
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- Number and extent of rills:** Rills are common, widely spaced, and continuous in absence of high gravel cover.

- Presence of water flow patterns:** Water flow patterns are common, continuous and occupy 15-20% of area.

- Number and height of erosional pedestals or terracettes:** Shrubs have symmetrical mounds caused by actions of splash, erosion, and rodent activity. There are no pedestals on rock or gravel fragments and no terracettes.

- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground is 10-60%. Expect low values in dry years.

- Number of gullies and erosion associated with gullies:** None.

- Extent of wind scoured, blowouts and/or depositional areas:** No evidence.

- Amount of litter movement (describe size and distance expected to travel):** Herbaceous litter can move by wind and water. Woody litter remains under shrub canopies.

-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil surface resistance to erosion is good under shrub canopies to moderate in interspaces due to crusts formed by raindrop impact.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Weak thin platy to granular; 7.5-10YR5/3 dry; 7.5-10YR4/3 moist, to 6 inches.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Canopy 10-15%. Herbaceous litter is present in some years and absent in others. Large shrubs with large coppice mounds with high infiltration rates. Subshrubs with small mounds with high infiltration rates. Mounds occupy 10-15% of the surface and are evenly spaced over the area.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Large shrubs > subshrubs > perennial grasses > winter annuals > summer annuals > succulents (note: in El Nino years, annual forbs and grasses are #1 in above ground weight.)
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** 30-70% canopy mortality on trees and shrubs; 90-100% mortality on perennial grasses.
-
14. **Average percent litter cover (%) and depth (in):** Herbaceous litter is not persistent on the site, but may be in El Nino years.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 270 lbs/ac unfavorable precipitation; 395 lbs/ac normal precipitation; 520 lbs/ac favorable precipitation.
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not**

invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Sahara mustard (potential), schismus

17. **Perennial plant reproductive capability:** Not impaired for shrubs; drought impaired for perennial grasses and forbs.
-