

Ecological site F041XA112AZ Sandy Wash 16-20" p.z. woodland

Last updated: 4/09/2021 Accessed: 05/12/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 041X–Madrean Archipelago

AZ 41.1 – Mexican Oak-Pine Forest and Oak Savannah

Elevations range from 4500 to 10,700 feet and precipitation ranges from 16 to 30 inches. Vegetation includes Emory oak, Mexican blue oak, Arizona white oak, one-seed juniper, alligator juniper, sacahuista, California bricklebush, skunkbush sumac, Arizona rosewood, wait-a-bit mimosa, sideoats grama, blue grama, purple grama, wooly bunchgrass, plains lovegrass, squirreltail, and pinyon ricegrass. The soil temperature regime ranges from thermic to mesic and the soil moisture regime ranges from aridic ustic to typic ustic. This unit occurs within the Basin and Range Physiographic Province and is characterized by numerous mountain ranges that rise abruptly from broad, plain-like valleys and basins. Igneous and metamorphic rock classes dominate the mountain ranges and sediments filling the basins represent combinations of fluvial, lacustrine, colluvial and alluvial deposits.

Ecological site concept

Sandy Wash, woodland, 16-20" p.z., is a gallery woodland dominated by oak trees (Emory and Arizona white oak). The sub-surface water table is seasonal; however, plants benefit from additional moisture received as rainfall runoff. Soils are sandy.

Associated sites

F041XA113AZ	Sandy Bottom 16-20" p.z. woodland
R041XA102AZ	Shallow Hills 16-20" p.z.
R041XA107AZ	Loamy Slopes 16-20" p.z.
R041XA108AZ	Loamy Upland 16-20" p.z.
R041XA110AZ	Sandy Loam Upland 16-20" p.z.
R041XA111AZ	Volcanic Hills 16-20" p.z.

Similar sites

R041XC316AZ	Sandy Wash	12-16" p.z.

Table 1. Dominant plant species

Tree	(1) Quercus emoryi (2) Quercus arizonica
Shrub	Not specified
Herbaceous	(1) Bouteloua curtipendula(2) Leptochloa dubia

Physiographic features

This site occurs in the middle elevations of the Madrean Basin and Range province in southeastern Arizona. It occurs on floodplains, low stream terraces and alluvial fans. It benefits on a regular basis from extra moisture received as over-bank flooding of stream channels and as runoff from adjacent uplands. It does not benefit from shallow water tables.

Landforms	(1) Flood plain(2) Stream terrace(3) Alluvial fan
Flooding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Flooding frequency	Occasional to frequent
Ponding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Ponding frequency	None to rare
Elevation	1,433–1,615 m
Slope	0–3%
Aspect	Aspect is not a significant factor

Table 2. Representative physiographic features

Climatic features

Precipitation in this zone of the common resource area ranges from 16-20 inches per year with elevations from 4700-5500 feet. Approximately 40% of this moisture comes as gentle rain or snow during the winter-spring (Oct-Apr) season; originates in the north Pacific and Gulf of California and comes as frontal storms with long duration and low intensity. The remaining 60% falls in the summer season (May-Sep); originates in the Gulf of Mexico and are convective, usually brief, intense thunderstorms. Snow is common Dec-Mar, averaging 5-15 inches per year, but rarely lasts more than a week. May and June are the driest months. Humidity is low.

Temperatures are mild. Freezing temperatures are common at night from Oct-May, but daytime temperatures are almost always over 40 F. Below 0 F temperatures can occur Dec-Feb. Daytime summer highs rarely exceed 95 F.

Species like plains lovegrass, bottlebrush squirreltail, Arizona white oak and Emory oak begin growth in late March to April. Warm season grasses begin growth in July or August with receipt of the first summer rains.

Table 3. Representative climatic features

Frost-free period (average)	175 days
Freeze-free period (average)	
Precipitation total (average)	457 mm

Influencing water features

There are no water features associated with this site.

Soil features

These are deep, young soils on gravelly and sandy alluvium of mixed origins. They are well drained and dark colored throughout. Soil surface textures range from very cobbly sandy loam to cobbly sand. Soils are sandy to coarse loamy. Plant-soil moisture relationships are very good due to the extra water received by the site.

This site is NOT CURRENTLY CORRELATED TO A SOIL SERIES IN ANY ARIZONA SOIL SURVEY AREA.

Table 4. Representative soil features

Surface texture	(1) Cobbly sand(2) Very gravelly sandy loam(3) Cobbly loamy sand
Family particle size	(1) Sandy
Drainage class	Somewhat excessively drained to well drained
Permeability class	Moderately rapid to rapid
Soil depth	152 cm
Surface fragment cover <=3"	10–60%
Surface fragment cover >3"	1–15%
Available water capacity (0-101.6cm)	4.57–15.24 cm
Calcium carbonate equivalent (0-101.6cm)	0–5%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–2
Soil reaction (1:1 water) (0-101.6cm)	6.1–7.8
Subsurface fragment volume <=3" (Depth not specified)	10–65%
Subsurface fragment volume >3" (Depth not specified)	1–15%

Ecological dynamics

The plant communities found on an ecological site are naturally variable. Composition and production will vary with yearly conditions, location, aspect, and the natural variability of the soils. The Historical Climax Plant Community represents the natural potential plant communities found on relict or relatively undisturbed sites. Other plant

communities described here represent plant communities that are known to occur when the site is disturbed by factors such as fire, grazing, or drought.

Production data provided in this site description is standardized to air dry weight at the end of the summer growing season. The plant communities described in this site description are based on near normal rainfall years.

NRCS uses a Similarity Index to compare existing plant communities to the plant communities described here. Similarity index is determined by comparing the production and composition of a plant community to the production and composition of a plant community described in this site description. To determine Similarity index, compare the production (air dry weight) of each species to that shown in the plant community description. For each species, count no more than the maximum amount shown for the species, and for each group, count no more than the maximum amount shown for the resulting total by the total normal year production shown in the plant community description. If the rainfall has been significantly above or below normal, use the total production shown for above or below normal years. If field data is not collected at the end of the summer growing season, then the field data must be corrected to the end of the year production before comparing it to the site description. The growth curve can be used as a guide for estimating production at the end of the summer growing season.

State and transition model



MLRA 41-1 (16-20"), Sandy Wash, QUAR, QUEM

Figure 6. State and Transition, Sandy Wash ,QUAR,QUEM 16-20"

State 1 Historic Climax Plant Community

Community 1.1 Historic Climax Plant Community



Figure 7. Sandy Wash QUAR, QUEM 16-20" pz. HCPC

This site has a mixed plant community with an over-story canopy of Arizona white oak and Emory oak, and lesser amounts of walnut, ash, desert willow and other trees. The under-story is dominated by perennial grasses and forbs. Evergreen live-oaks change their leaves in April and May. The aspect is evergreen woodland. Channel and stream bank erosion and sedimentation are natural features of this site. Stream flow is intermittent and surface flow occurs sporadically during the winter months only. Springtime stream flow is not sufficient enough for the broadleaf deciduous riparian species. Oak species present on the site reproduce from acorns which germinate in the summer immediately after being produced. These species also sprout after cutting, flood injury, or fire. Tree canopy ranges from 20 -50%. Number of stems ranges from 200-400 per acre. Under-story composition and production varies with percent canopy cover. Trees reach maximum size on this site. Disturbance of the dominant native tree cover on the site can lead to increases of species such as mesquite and juniper. Disturbance of the under-story can lead to increases in species like southwestern rabbitbrush and wait-a-bit mimosa. Management of the under-story should be aimed at leaving enough cover to protect the soils from erosion and to trap sediment in the summer flooding season.

Forest overstory. The overstory part of the plant community is dominated by Arizona white oak and Emory oak. There may be lesser amounts of grey oak and alligator juniper. Deciduous tree species are present in minor amounts. They include netleaf hackberry, Arizona black walnut, Arizona ash, desert willow and an occasional Arizona sycamore.

Tree canopy ranges from 20 to 50 percent. Trees reach maximum size (40 feet) on the site.

"Forest Overstory Composition" as shown in the following tables is in percent canopy cover and not percent frequency.

The "Plant Community Annual Production" table shows production for 50% canopy as low, 35% canopy as the representative value and 20% canopy as the high value.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Tree	1681	2802	4483
Grass/Grasslike	504	1345	2354
Forb	39	112	224
Shrub/Vine	22	112	168
Total	2246	4371	7229

Table 6. Soil surface cover

Tree basal cover	1%
Shrub/vine/liana basal cover	0%
Grass/grasslike basal cover	3-10%
Forb basal cover	0-5%
Non-vascular plants	0%
Biological crusts	0%
Litter	50-80%
Surface fragments >0.25" and <=3"	10-60%
Surface fragments >0.25" and <=3" Surface fragments >3"	10-60% 1-15%
Surface fragments >0.25" and <=3" Surface fragments >3" Bedrock	10-60% 1-15% 0%
Surface fragments >0.25" and <=3" Surface fragments >3" Bedrock Water	10-60% 1-15% 0% 0%

Table 7. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	-	_	0-1%	0-2%
>0.15 <= 0.3	-	0-1%	1-5%	1-5%
>0.3 <= 0.6	-	0-2%	5-20%	1-10%
>0.6 <= 1.4	-	0-5%	15-30%	0-10%
>1.4 <= 4	0-2%	0-5%	0-5%	0-5%
>4 <= 12	19-45%	_	_	_
>12 <= 24	1-10%	_	-	_
>24 <= 37	-	_	-	_
>37	-	_	-	_

Figure 9. Plant community growth curve (percent production by month). AZ4111, 41.1 16-30. Growth begins in the spring, semi-dormancy occurs during the June drought, most growth occurs during the summer rainy season..

Jan	Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	5	10	0	15	45	20	5	0	0

State 2 Exotic grasses in under-story

Community 2.1

Exotic grasses in under-story



Figure 10. Sandy Wash, QUAR, QUEM 16-20" pz. Lehmann

This state occurs where exotic grasses like Lehmann, Boer, weeping lovegrass, bermuda, Johnson grass, and / or yellow bluestem has invaded from adjacent or up-stream areas with a seed source. As the non-native grasses increase to dominate the under-story, native perennial grasses and forbs are reduced to minor amounts. Fires will act to increase the dominance of Lehmann lovegrass in the under-story.

State 3 Shrub invaded

Community 3.1 Shrub invaded



Figure 11. Sandy Wash, QUAR, QUEM 16-20" pz. juniper

This state occurs where mesquite and / or juniper species invade and increase in the absence of fire for long periods of time. These species can dominate the openings in the tree canopy with cover ranging from 5 to 30%. Fires can still burn after wet summers but these shrubs are well established and will sprout and quickly dominate the openings again.

State 4 Eroded channel

Community 4.1 Eroded channel

This state occurs where watersheds are destabilized buy continuous grazing and / or the interaction of climate and severe fire. Large amounts of runoff and sediment are delivered to this site and cause gully and stream bank erosion. Channel erosion can deepen floodways and reduce flooding of the stream terraces. Sediment can clog

channels causing new channels to cut through terraces; up-rooting trees and removing cover and woody debris.

Additional community tables

Animal community

The under-story plant community is suitable for grazing by all classes of livestock at any season. Use should be limited to 50% or less to leave enough under-story material to protect soils from erosion and to trap sediment during the spring and summer flooding seasons. Oak seedlings establish in late summer and fall from the preceding acorn crop. In years with high germination, deferment from grazing for the following winter and spring, followed by grazing in the summer will help establishment by selective grazing of grass to reduce competition with oak seedlings. Seasonal carrying capacities should be evaluated based upon on-site investigations as under-story varies greatly with percent of canopy cover and production can vary greatly from year to year depending on seasonality and frequency of flooding.

This is a very productive and widely used wildlife habitat. Excellent diversity and food, cover, edge with adjacent grasslands and seasonally available water make this site home to a great variety of wildlife species. Acorn production can be very high. Mature Emory oaks can produce up to 15,000 acorns per tree in a season and mature Arizona white oaks can produce double that amount.

Hydrological functions

These sandy wash bottoms are extremely important for ground water recharge in upper basins along the mountain front. Deep, very coarse textured soils allow flood water and winter stream-flow to percolate to great depths.

Recreational uses

Camping, hunting, picnicking, bird watching, horseback riding, hiking, photography

Wood products

Net annual growth of oaks on this site averages about 1% of total volume. Heart-rot sets in on stems at 20+ years of age. If harvested on a 25-30 year rotation, production of coppice volume will be much higher (3% total volume) than yearly production of old age stands (due to lack of heart-rot). CMAI for coppice rotation on this site ranges from 5-15 ft3/ac/yr. These species take a long time to reach sexual maturity. Emory oak does not bear acorns until 40 yrs old. Arizona white oak does not bear acorns until 80 years old. Productivity declines rapidly after 160 years of age. The lifespan of these species are from 200-300 years. Dead limbs in mature stands range from 10-15% of total volume. Trees with 12-inch diameter stems are probably between 60-80 years old. Emory oak grows rapidly from 0-10 years, moderately 10-20 years and slowly 20 years on.

Equipment is limited by stony, cobbly soils, flooding that is common, and a high water table in winter.

Erosion potential is severe for water erosion during flooding.

Compaction potential and workability for roads is poor due to coarse textured, cobbly, stony soils. Rutting potential is low. Revegetation potential is good due to excellent climate.

Prescriptions for harvest cutting should be designed to allow for harvest of wood products and to preserve other values like wildlife, scenic, forage, and hydrologic. A 25-30 year rotation on coppice sprouts will maximize wood harvest. Initial volume reduction can be from 30-50% with a second harvest of 30-50% at 10-15 years and a third harvest on the older coppice sprouts which will average 20 ft with 8-10 inch diameters.

Thinning areas to 20% canopy will maximize forage production for livestock. Thinning to leave 10 feet between tree canopies will maximize wood production.

Prescribed burning in the cool season can reduce ladder fuels and freshen coarse forage species like sacaton and deergrass. Mechanical tree removal is severely limited due to wet soils caused by winter flooding. Control mistletoe to limit tree damage and loss. Coppice treatment at appropriate intervals to avoid loss of production to heart-rot

fungus. Replanting is suitable for oak (summer and fall) and walnut (spring). Seed mortality has slight limitations due to coarse textured soils. Natural regeneration is good as long as natural flood regime is unaltered. Oak and walnut seedlings need protection from fire for 2-3 years to establish. Plant competition for tree seedlings is severe from grass and other herbaceous species. Prescribed grazing can reduce competition. Wind-throw hazard is slight.

Other products

Acorns (bellotas) and walnuts for food. Medicinal herbs like yerba de pasmo, herbaceous sage and Hopi tea. Herbs for seasoning like wild oregano and terragon.

Table 8. Representative site productivity

Common Name	Symbol	Site Index Low	Site Index High	CMAI Low	CMAI High	Age Of CMAI	Site Index Curve Code	Site Index Curve Basis	Citation
Arizona white oak	QUAR	30	40	5	15	-	-	-	
Emory oak	QUEM	30	40	5	15	_	-	-	

Inventory data references

Range 417s include 1 in good condition. Wood 5s include 1 at 35% canopy cover.

Type locality

Location 1: Cochise County, AZ					
Township/Range/Section	T21s R19e S33				
General legal description	SW 1/4 of section - Ft. Huachuca at the mouth of Blacktail Canyon				
Location 2: Cochise Coun	ity, AZ				
Township/Range/Section	TT17S R28E S15				
General legal description	Willcox - Oak Ranch - Fivemile Creek				
Location 3: Pinal County, AZ					
Township/Range/Section	T21S R18E S30				
General legal description	NE 1/4 of section - Tucson - Jelks Triangle M. Ranch - Post Canyon				

Contributors

Dan Robinett Larry D. Ellicott

Approval

Curtis Talbot, 4/09/2021

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	

Date	05/12/2025
Approved by	Curtis Talbot
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values):
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 17. Perennial plant reproductive capability: