

## Ecological site R042AD005NM Clay Loam Upland, Dry Mixed Prairie

Accessed: 05/11/2025

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

**Figure 1. Mapped extent**

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

### Physiographic features

This site series consists of very deep, moderately slowly permeable soils that formed in calcareous alluvium. These gently sloping soils are on inset fans, alluvial fan and fan remnants. Slope ranges from 2 to 5 percent. Mean annual precipitation is about 13 inches and the mean annual temperature is about 60 degrees F.

**Table 2. Representative physiographic features**

Landforms	(1) Alluvial fan
Flooding frequency	None
Ponding frequency	None
Elevation	4,700–6,000 ft
Slope	2–5%
Aspect	Aspect is not a significant factor

### Climatic features

Average precipitation for this site is approximately 12 to 14 inches. Variations of 5 inches are not uncommon. Approximately 75 percent occurs from May through October with most of the rainfall occurring from July to September. Most of the summer precipitation comes in the form of high intensity, short duration thunderstorms. Rain and snow of low intensity characterize the limited winter precipitation.

Temperatures are mild. Freezing temperatures are common at night from December through April, however, temperatures during the day are frequently above 50 degrees F. Occasionally in December to February brief periods of 0 degrees F. temperatures may be expected. During June to August some days may exceed 100 degrees F.

The mean annual precipitation figures are derived from rain gauge data collected by the BLM (1971 to 1990), and NOAA weather maps utilizing prism model estimation techniques. There are no permanent weather stations within the boundaries of the Land Resource Unit.

**Table 3. Representative climatic features**

Frost-free period (average)	180 days
Freeze-free period (average)	185 days
Precipitation total (average)	14 in

## Influencing water features

This site is not influenced by water from wetland or stream.

## Soil features

Soils are deep or very deep. Surface textures are silt loams. Underlying layers are loam, silt loam, silty clay loam or clay loam with Clay content ranging from 27 to 35 percent and 0 to 5 percent coarse fragments. The soils are well drained and have a moderately slow permeability. Available water holding capacity to a depth of 40 inches is moderate. The site usually receives some extra water from higher, surrounding terrain. If unprotected by adequate vegetative cover soils become susceptible to water erosion.

Minimum and maximum values listed below represent the characteristic soils for this site.

Characteristic soils:

Double

**Table 4. Representative soil features**

Surface texture	(1) Sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained to moderately well drained
Permeability class	Moderately slow
Soil depth	60 in
Surface fragment cover <=3"	0–2%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	7–8 in
Calcium carbonate equivalent (0-40in)	0–15%
Electrical conductivity (0-40in)	0–2 mmhos/cm
Sodium adsorption ratio (0-40in)	0–1
Soil reaction (1:1 water) (0-40in)	7.9–8.4
Subsurface fragment volume <=3" (Depth not specified)	0–5%
Subsurface fragment volume >3" (Depth not specified)	0%

## Ecological dynamics

The Clay Loam Upland site is associated with the Limestone Hills ecological site. The Clay loam Upland site is

positioned down slope from the Limestone hills site from which it receives run-on water. This site can also occur adjacent to the Gravelly, Shallow Sandy, and Limy ecological sites.

The aspect of the site is that of open grassland sparsely dotted with shrubs. Warm season perennial grasses dominate this community. Alkali sacaton and blue grama are the dominant plants both in aspect and composition. Forb production and composition fluctuate from year to year. Some of the more common forbs include grassland croton, bladderpod and desert holly.

Characteristic shrubs include soaptree yucca, winterfat, four-wing saltbush and prickly pear. This site is resistant to state change and shrub invasion due to the extra run-on water it receives. If the hydrology is altered and the extra run-on water is diverted, the site dries and more drought tolerant species eventually dominate. As the site dries plant community dominance shifts from alkali sacaton to one dominated by tobosa or tobosa and burrograss. Drought and overgrazing can expedite this state change by reducing grass cover and organic matter and increasing the amount of bare ground.

## State and transition model




Figure 4. MLRA42, SD4, Clay Loam Upland

## State 1 HCPC

The historic community for this site is characterized by a dominance of alkali sacaton. Blue grama is secondary and is evenly dispersed throughout the community. A fair amount of tobosa appears in patches scattered across the site. Other common grass species include sand muhly, ear muhly and burrograss. Forb composition varies seasonally and yearly, but bladderpod and croton are two of the more consistent forbs. Shrubs are only a minor component, and occur as widely spaced individuals. Soaptree yucca is the most common shrub on this site; others include winterfat, four-wing saltbush, prickly pear and cholla. Grazing induced retrogression can cause an increase in tobosa and a decrease in alkali sacaton and more palatable species such as blue grama and vine mesquite.

Drought or heavy continuous use on alkali sacaton will cause it to thin out and form clumps or tussocks with bare areas in between, thereby decreasing cover and increasing the amount of bare ground. This site however, is resistant to change if the hydrology remains intact and recovers rapidly during years of adequate precipitation.

Community 1.1  
Grassland



Figure 5. Grassland

Alkali sacaton and blue grama are the dominant species. Grass cover is uniformly distributed with few large bare areas. There is little evidence of active rills and no gully formation if plant cover remains intact. Litter movement associated with overland flow is limited to smaller size class litter and short distances. Shrubs are a minor component of the site. The aspect of this site is open grassland sparsely dotted with shrubs. Warm season perennial grasses dominate this site. Alkali sacaton and blue grama are the dominant plants both in aspect and composition. Forb production fluctuates from year to year and season to season.

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	920	1330	1700
Forb	65	56	82
Shrub/Vine	15	14	18
Total	1000	1400	1800

Figure 7. Plant community growth curve (percent production by month).  
NM5805, R042XD005NM Clay Loam Upland HCPC Grassland State.  
R042XD005NM Clay Loam Upland Warm Season Grassland - Average rainfall year.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	3	8	7	18	28	25	6	2	0

State 2  
Dry Grassland State

This state is characterized by a change in hydrology. Historic vegetation is supported by run-on water from higher adjacent sites. When a road, trail, or other natural or man-made disturbances cause an obstruction or diversion of natural flow patterns the hydrology is altered. Without the run-on water the site dries. Alkali sacaton and blue grama decrease and tobosa and burrograss increase. Alkali sacaton cover is no longer uniformly distributed, the plants clump up forming tussocks with interconnected bare areas between plants. Physical crusts form in the bare areas when soil particles dislodged by erosion plug the pores in the soil, reducing infiltration. These physical crusts discourage grass seedling establishment and provide pathways for the redistribution of surface litter and organic matter. The decline of alkali sacaton and blue grama may be accelerated by decreased infiltration, drought or

overgrazing. Diagnosis: Tobosa is the dominant grass. Alkali sacaton cover is reduced and if it continues to decline becomes patchy. Burrograss cover increases and may become co-dominant with tobosa. Large connected bare patches are evident. Production is reduced compared with the sacaton grassland state.

## Community 2.1

### Dry Grassland



Figure 8. Dry Grassland

Tobosa is the dominant grass. Alkali sacaton cover is reduced and if it continues to decline becomes patchy. Burrograss cover increases and may become co-dominant with tobosa. Large connected bare patches are evident. Production is reduced compared with the sacaton grassland state.

## Transition 1a

### State 1 to 2

Key indicators of approach to transition: Reduction in alkali sacaton and blue grama cover and increase in size and frequency of bare patches. Increase in tobosa and or burrograss cover. The formation of roads, gullies or other features (on or off site) that disrupts natural overland flow on site. Formation of physical crusts—indicating loss of organic matter and decrease in soil aggregate stability and reduced infiltration.<sup>1</sup> Evidence of litter movement—indicating loss or redistribution of organic matter.

## Transition 1b

### State 2 to 1

Restoration of natural hydrologic processes is necessary to reverse the transition back to the Sacaton Grassland State. Prescribed grazing will help to restore cover, increase organic matter, and infiltration.

## Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1				560–700	
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	560–700	–
2				210–350	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	210–350	–
3				112–210	
	tobosagrass	PLMU3	<i>Pleuraphis mutica</i>	112–210	–
4				42–70	
	vine mesquite	PAOB	<i>Panicum obtusum</i>	30–70	–
	burrograss	SCBR2	<i>Scleropogon brevifolius</i>	30–60	–
	ear muhly	MUAR	<i>Muhlenbergia arenacea</i>	30–60	–
	sand muhly	MUAR2	<i>Muhlenbergia arenicola</i>	30–60	–
<b>Forb</b>					
5				15–20	
	croton	CROTO	<i>Croton</i>	15–20	–
	bladderpod	LESQU	<i>Lesquerella</i>	15–20	–
6				3–8	
	dwarf desertpeony	ACNA2	<i>Acourtia nana</i>	3–8	–
	broadleaved pepperweed	LELA2	<i>Lepidium latifolium</i>	3–8	–
7				10–14	
	Forb, perennial	2FP	<i>Forb, perennial</i>	10–14	–
8				10–14	
	Forb, annual	2FA	<i>Forb, annual</i>	10–14	–
	herb sophia	DESO2	<i>Descurainia sophia</i>	10–14	–
<b>Shrub/Vine</b>					
9				5–7	
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	5–7	–
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	5–7	–
10				5–7	
	American tarwort	FLCE	<i>Flourensia cernua</i>	5–7	–
	creosote bush	LATR2	<i>Larrea tridentata</i>	5–7	–
	pricklypear	OPUNT	<i>Opuntia</i>	5–7	–
	soaptree yucca	YUEL	<i>Yucca elata</i>	5–7	–

## Animal community

The open grassland aspect of this site attracts pronghorn antelope. Mule deer occasionally use the site in conjunction with adjacent sites. This site also supports black-tailed jackrabbits, meadowlark, mourning dove, and scaled quail and provides nesting, hiding and thermal cover for a variety of small rodents, birds and reptiles and their associated predators.

## Hydrological functions

Most summer rainfall occurs as brief sometimes-heavy thunderstorms. Runoff is medium, and the hazard of water erosion is moderate.

Soil Name Hydrologic Group

Double B

## Recreational uses

This site offers good potential for antelope and predator hunting, wildlife observation and photography. Scenic beauty of this site will especially appeal to those who value wide open prairie grasslands.

## Wood products

This site has no significant value for wood products.

## Other products

Grazing: This site is suitable for grazing by all kinds and classes of livestock during all seasons of the year. Currently the majority of the livestock use on this site is with mother cows in cowcalf operations. Historic use has been sheep and cattle. As the site deteriorates there will be an increase in bare ground leaving the exposed soil susceptible to wind and water erosion. This site responds best to a system of management that rotates the season of use. This site is suitable for artificial revegetation depending on slope and amount of extra run-on water.

Initial starting stocking rates will be determined with the landowner or decision-maker. They will be based on past use histories and type and condition of the vegetation. Calculations used to determine initial starting stocking rate will also be based on forage preference ratings.

## Inventory data references

Supporting information includes limited clipping data, soil survey investigations, aerial photographs, and personal observations.

## Other references

1. U.S. Department of Agriculture, Natural Resources Conservation Service. 2001. Soil Quality Information Sheet. Rangeland Soil Quality—Organic Matter, Aggregate Stability, Infiltration. Rangeland Sheets 6,3,5 [Online]. Available:  
<http://www.statlab.iastate.edu/survey/SQL/range.html>

## Contributors

David Trujillo  
Jason S. Martin

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	

Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:



Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 

14. **Average percent litter cover (%) and depth ( in):**
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
- 

17. **Perennial plant reproductive capability:**
-