

Ecological site R042BB011NM

Deep Sand, Desert Shrub

Accessed: 05/11/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Associated sites

R042BB012NM	Sandy, Desert Shrub This site often integrates with Sandy Sites.
R042BB024NM	Gravelly Sand, Desert Shrub This site often integrates with Gravelly Sand sites.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site occurs on level to gently sloping old eolian and alluvial deposits. Slopes range from 1 to 9 percent and average about 3 percent. Elevations range from about 3,800 to 5,000 feet.

Table 2. Representative physiographic features

Landforms	(1) Basin floor (2) Sand sheet (3) Fan remnant
Flooding duration	Extremely brief (0.1 to 4 hours) to brief (2 to 7 days)
Flooding frequency	Very rare to rare
Ponding frequency	None
Elevation	1,158–1,524 m
Slope	1–9%
Aspect	Aspect is not a significant factor

Climatic features

Annual average precipitation ranges from 7.35 to 11.90 inches. Wide fluctuations from year to year are common, ranging from a low of about 2 inches to a high of over 20 inches. At least one-half of the annual precipitation comes in the form of rainfall during July, August, and September. Precipitation in the form of snow or sleet averages less

than 4 inches annually. The average annual air temperature is about 60 degree F. Summer maximums can exceed 100 degrees F. and winter minimums can go below zero. The average frost-free season exceeds 200 days and extends from April 1 to November 1. Both the temperature regime and rainfall distribution favor warm-season perennial plants on this site. Spring moisture conditions are only occasionally adequate to cause significant growth during this period of year. High winds from the west and southwest are common from March to June, which further tends to create poor soil moisture conditions in the springtime.

Climate data was obtained from
<http://www.wrcc.dri.edu/summary/climsmnm.html>
 web site

Table 3. Representative climatic features

Frost-free period (average)	205 days
Freeze-free period (average)	227 days
Precipitation total (average)	305 mm

Influencing water features

This site is not influenced by water from wetland or stream.

Soil features

The soils are moderately deep to deep. The surface layer is typically a sand or loamy sand overlying fine sandy loams, loamy sands to sands. They may have accumulations of calcium carbonate below 20 inches. The soils are frequently calcareous throughout. They are well drained and moderately rapid to rapidly permeable with moderate water-holding capacity. Because of the sandy surface and inability to hold a large amount of water, if unprotected, wind blowing becomes severe.

Table 4. Representative soil features

Surface texture	(1) Loamy fine sand (2) Loamy sand (3) Sand
Family particle size	(1) Sandy
Drainage class	Well drained to excessively drained
Permeability class	Moderately slow to rapid
Soil depth	61–152 cm
Surface fragment cover <=3"	0–5%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	5.08–12.7 cm
Electrical conductivity (0-101.6cm)	0–8 mmhos/cm
Soil reaction (1:1 water) (0-101.6cm)	7.4–8.4
Subsurface fragment volume <=3" (Depth not specified)	0–5%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics


Overview:

This site often intergrades with either the Sandy or Gravelly sand sites. Deep sand sites located on hills, known as Sandhills sites in Texas, are present in SD-2 and are included here. The soil profiles of dunes in the mesquite-dominated state in the Sandy site are similar to those of Deep sand sites, and this state and site have been equated. It is unknown, however, if mesquite coppice dunes sites could support Deep sand plant communities. They may not, due perhaps to differences in landscape position or the size of the deep sand soil patches, which are relatively small in Sandy sites. Until it is demonstrated that Sandy-derived coppice dunes can be rehabilitated as Deep sand sites (which has not yet been attempted so far as the author is aware) it is best to treat these as separate classifications. The historic plant community type of the Deep sand site is dominated by dropseeds (*Sporobolus flexuosus*, *S. contractus*, *S. cryptandrus*), especially the giant dropseed (*S. giganteus*). Black grama (*Bouteloua eriopoda*) and bush muhly (*Muhlenbergia porteri*) are also important grasses. Under heavy grazing pressure, the more palatable black grama and bush muhly decline, and eventually dropseeds decline. Shrubs that are common in the site, including sand sage (*Artemisia filifolia*) and broom dalea (*Psoralea scoparius*), increase in representation and cover with grazing or perhaps due to climate change. Honey mesquite (*Prosopis glandulosa*) may invade or increase. Shrub control and grazing management results in varying degrees of restoration or grass cover in some cases. In other cases, grass cover is highly reduced and broom dalea and/or sand sage persist as dominants or mesquite dunes may persist. It is not known what soil or climatic changes underlie these persistent vegetation changes.

No systematic studies of communities, states or transitions have been performed in Deep sand sites in SD-2. Because of the unique plant communities and soils of these sites, studies of soil variation among sites exhibiting different states would provide a useful contribution.

State and transition model

State-Transition model: MLRA 42, SD-2, sandy subgroup: Deep sand



- 1a. Persistent reduction of grasses, extinction of seed pool, competitive dominance by shrubs
- 1b. Shrub removals and possibly seed additions
- 2a. Soil degradation?, climate change?, mesquite seed introduction.
- 2b. Shrub removal and soil amendments?
- 3a. Mesquite invasion. 3b. Shrub removal and prevention of seed additions

State 1
Historic Climax Plant Community

Community 1.1
Historic Climax Plant Community

MLRA 42; SD-2; Deep sand

Dropseed-dominated



- Giant dropseed, spike dropseed, soaptree yucca
- Normally has patches of open ground due to dominance of bunchgrasses, abundant litter
- Wind-driven sand trapped by grasses
- Copia sand, soils, El Paso, TX

Broom dalea-sand sage, broom dalea/dropseeds



- Giant dropseed patch in a matrix of broom dalea, saltbush, and other shrubs
- More open ground, less litter
- Sand blowouts and during apparent
- Copia sand, Fort Bliss, MacGregor Range Camp, TX

Broom dalea-sand sage, broom-dalea/saltbush



- Broom dalea dominant, followed by saltbush and rosemarymint
- Unclear if this used to have grass
- Sand blowouts and during apparent
- Copia sand, Fort Bliss, MacGregor Range Camp, TX

Mesquite-dominated



- Mesquite dominant left, Bluepoint loamy sand, Dona Ana Co., NM
- Little-leaf sumac dominant right, sand deposition over limestone (= sandhills type). Fort Bliss, MacGregor Range Camp, TX
- Extensive areas of bare ground, erosion apparent
- Unclear why mesquite invades in some situations

Figure 4. MLRA 42; SD-2; Deep sand

State Containing Historic Climax Plant Community: Dropseed-dominated grassland: The historic community of this site is dominated by dropseeds (dropseed/other grasses community) and features a significant cover of black grama and bush muhly. Sand sage, broom dalea, yucca (*Yucca elata*), fourwing saltbush (*Atriplex canescens*) are also common components. Dropseed abundance may fluctuate considerably with drought-wet cycles. Grazing-induced retrogression leads initially to a loss of black grama and bush muhly (dropseed/shrubs community) and eventually to a reduction in dropseeds (Shrubs/dropseed community). Honey mesquite may encroach or invade under some circumstances, although there are deep sand communities where this invasion seems to be limited. It is not known if the nutrient redistribution processes and loss of inter-shrub soil fertility observed in Sandy sites (see the Sandy model) is important in this site. It is believed that in the absence of grazing, invasion and growth of saltbush (a highly palatable shrub) may eventually lead to mesquite mortality (saltbush/shrubs community) and a return to dropseed dominance (if not a restoration of the dropseed community). There are no data to suggest a mechanism for this pattern, nor are there any rigorous documentations of the pattern. There are, however, areas on deep sand soils within the city of Las Cruces, NM, that are ungrazed and are now dominated by saltbush and large mesquite plants. Such locations may be the result of protection from grazing and saltbush proliferation. If the saltbush-succession process does occur, then mesquite-invaded communities can be considered to occur within the dropseed dominated state. Diagnosis: Dropseed cover is dominant or substantial, black grama and bush muhly usually present and may dominate in some cases. Transition to broom dalea-sand sage state (1a): The causes of the transition from the broom dalea/sand sage state are unknown. Given the wide amplitude of conditions across which dropseed communities are believed to be recoverable, it seems possible that shrubs may exert a strong competitive influence over the system once they establish in high densities. Changes in the seasonality of precipitation may be important. Key indicators of approach to transition: Decreases in dropseeds may be a necessary prerequisite for this transition but it is not sufficient to indicate it. Transition to mesquite dune state (2a): The causes of the transition to domination by mesquite are unknown. Heavy grazing or destruction of plants by trampling or vehicles with consequent erosion from around mesquite and loss of soil fertility in shrub interspaces is one possible mechanism. Autogenic (mesquite-caused) degradation of soils may also play a role. Key indicators of approach to transition Decreases in dropseeds and other shrubs may be a necessary prerequisite for this transition but it is not sufficient to indicate it. Changes in the seasonality of precipitation may be important.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	137	305	471
Shrub/Vine	36	78	121
Forb	24	52	81
Total	197	435	673

Table 6. Soil surface cover

Tree basal cover	0%
Shrub/vine/liana basal cover	0%
Grass/grasslike basal cover	15%
Forb basal cover	0%
Non-vascular plants	0%
Biological crusts	0%
Litter	9%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	75%

**Figure 6. Plant community growth curve (percent production by month).
NM2503, R042XB011NM-Deep Sand Warm Season Plant HCPC. SD-2 Deep
Sand HCPC Warm Season Plant Community.**

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	5	10	10	25	30	15	5	0	0

State 2

Broom dalea-Sandsage State

Community 2.1

Broom dalea-Sandsage State

Additional States: Broom-dalea-sand sage: Dropseed abundance is normally variable across years as these grasses are drought sensitive, but if seeds are depleted, or due to the persistent effects of soil degradation or to the effects of herbivory by small animals (see Sandy model), the average abundance of dropseeds at a site has declined. This may be coincident with abundant cover of broom dalea and sand sage, as well as other shrubs including mesquite and little-leaf sumac (*Rhus microphylla*). In some areas, broom dalea and other shrubs are extremely dominant and this appears to be persistent. For example, these communities have been observed on the Fort Bliss, which has been ungrazed for over 50 years. The causes of broom dalea or sand sage dominance are not understood. A persistent broom dalea state may be associated with a petrocalcic horizon underlying the soil at a depth of 100-150 cm (David Trujillo, NRCS, personal communication). Diagnosis: Dropseed cover is sparse and patchy to absent. Bare ground cover may be low due to the high cover of shrubs. Transition to dropseed-dominated state (1b): Seeding and/or shrub removal with herbicides. In many cases, dropseed cover recovers only partially and temporarily, and shrubs may recover. Transition to mesquite dune state (3a): Causes are unknown. Mesquite invasion of the broom dalea-sand sage state is possible.

State 3

Mesquite dune State

Community 3.1

Mesquite dune State



Figure 7. mesquite dune state

Mesquite dune: In many areas, only a few species of shrubs may dominate, shrub and grass cover is sparse, and mesquite may form coppice dunes. Where gravel content is high (e.g., >5%), creosotebush (*Larrea tridentata*) may also be present. In some cases, mesquite may be extremely dominant and few other shrubs are present. On sandhills, little-leaf sumac may attain extreme dominance instead of mesquite. Dynamics within this state may occur and be attributable to climate or grazing-associated disturbances (e.g. trampling of shrubs) or to off-road vehicle activity. In general, wind and water erosion rates appear to be high in this state due to the low plant cover, and this likely maintains very low soil fertility. The conditions under which mesquite attains dominance are not understood. Diagnosis: Dropseed cover sparse to absent. Bare ground patches are large, forming blowouts and “streets” (i.e. elongated patches in the direction of prevailing winds). Evidence of interdune erosion (pedestalling, blowouts) are present. Transition to dropseed-dominated state (2b): Possibly with seeding and/or shrub removal with herbicides. Soil stabilization and fertility amendments would probably be needed; this has not been attempted. Transition to broom dalea-sand sage state (3b): Strategy unknown. Information sources and theoretical background: Communities, states, and transitions are based upon information in the ecological site description and observations by Jim Powell, NRCS (retired)

Additional community tables

Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Warm Season			87–109	
	spike dropseed	SPCO4	<i>Sporobolus contractus</i>	87–109	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	87–109	–
	mesa dropseed	SPFL2	<i>Sporobolus flexuosus</i>	87–109	–
2	Warm Season			130–152	
	giant dropseed	SPGI	<i>Sporobolus giganteus</i>	130–152	–
3	Warm Season			21–44	
	black grama	BOER4	<i>Bouteloua eriopoda</i>	21–44	–
4	Warm Season			4–21	
	bush muhly	MUPO2	<i>Muhlenbergia porteri</i>	4–21	–
5	Warm Season			4–21	
	plains bristlegass	SEVU2	<i>Setaria vulpiseta</i>	4–21	–
6	Warm Season			4–13	
	threeawn	ARIST	<i>Aristida</i>	4–13	–

	low woollygrass	DAPU7	<i>Dasyochloa pulchella</i>	4–13	–
7	Warm Season			4–13	
	Graminoid (grass or grass-like)	2GRAM	<i>Graminoid (grass or grass-like)</i>	4–13	–
Shrub/Vine					
8	Shrub			21–44	
	sand sagebrush	ARFI2	<i>Artemisia filifolia</i>	21–44	–
	broom dalea	PSSC6	<i>Psoralea scoparius</i>	21–44	–
	yucca	YUCCA	<i>Yucca</i>	21–44	–
9	Shrub			4–21	
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	4–21	–
	littleleaf sumac	RHMI3	<i>Rhus microphylla</i>	4–21	–
10	Shrub			4–21	
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	4–21	–
11	Shrub			4–21	
	mormon tea	EPVI	<i>Ephedra viridis</i>	4–21	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	4–21	–
Forb					
12	Forb			4–13	
	dwarf desertpeony	ACNA2	<i>Acourtia nana</i>	4–13	–
	woolly plantain	PLPA2	<i>Plantago patagonica</i>	4–13	–
13	Forb			4–13	
	herb sophia	DESO2	<i>Descurainia sophia</i>	4–13	–
	touristplant	DIWI2	<i>Dimorphocarpa wislizeni</i>	4–13	–
	redstem stork's bill	ERCI6	<i>Erodium cicutarium</i>	4–13	–
	buckwheat	ERIOG	<i>Eriogonum</i>	4–13	–
	spurge	EUPHO	<i>Euphorbia</i>	4–13	–
	bladderpod	LESQU	<i>Lesquerella</i>	4–13	–
14	Forb			13–21	
	Indian blanket	GAPU	<i>Gaillardia pulchella</i>	13–21	–
	woolly paperflower	PSTA	<i>Psilostrophe tagetina</i>	13–21	–
	globemallow	SPHAE	<i>Sphaeralcea</i>	13–21	–
15	Annual Forbs			4–21	
	Forb, annual	2FA	<i>Forb, annual</i>	4–21	–
16	Perennial Forbs			4–21	
	Forb, perennial	2FP	<i>Forb, perennial</i>	4–21	–

Animal community

This site provides habitat which supports a resident animal community that is characterized by pronghorn antelope, badger, kit fox, spotted ground squirrel, desert pocket mouse, Ord kangaroo rat, southern plains woodrat, scaled quail, pyrrhuloxia, roadrunner, burrowing owl, desert horned lizard, and Couch's spadefoot toad.

Where large mesquite and yucca are present on this site, white necked raven and mourning dove nest. When site deterioration produces a dune-interdune aspect with mesquite invasion,

animal population densities shift in favor of burrowing mammals, their predators and hru bdependent birds.

Hydrological functions

The runoff curve numbers are determined by field investigations using hydraulic cover conditions and hydrologic soil groups.

Hydrologic Interpretations

Soil Series-----Hydrologic Group

Bluepoint-----A

Pintura-----A

Yturbide-----A

Recreational uses

Suitability for camping and picnicking is fair, and hunting is fair for pronghorn antelope, quail, dove, small game, and waterfowl where seasonal open water occurs. Photography and bird watching for numerous birds, raptors and others can be fair to good, especially during migration seasons. Most small animals of the site are nocturnal and secretive, seen only at night, early morning or evening. Scenic beauty is greatest during spring and sometimes summer months when flowering of forbs, shrubs, and cacti occurs.

Wood products

The site has no significant value for wood products.

Other products

This site at its potential is suitable for grazing in all seasons of the year. Green forage in the form of forbs and a few early season grasses are occasionally produced in significant amounts in the spring. The dominant production comes in the summer. The site is suitable for use by all classes of livestock. Site deterioration caused by inadequate grazing management is characterized by a decrease or disappearance of black grama and bush muhly, an increase in sand sagebrush and broom delea, mesquite invasion, and eventual dunning or hummocking. The dropseeds may also decrease, coming and going with good and bad years but unable to hold a consistent footing even in normal years. Site recovery can be reasonably rapid following mesquite control coupled with good grazing management. Given adequate opportunity, secondary succession will take place, usually by fourwing saltbush replacing the mesquite, and after several years a gradual re-establishment of the dropseeds and other plants more typical of the natural potential for the site.

Other information

Guide to Suggested Initial Stocking Rate Acres per Animal Unit Month

Similarity Index-----Ac/AUM

100 - 76-----5.0 – 5.9

75 – 51-----5.6 – 7.1

50 – 26-----6.8 – 16.0

25 – 0-----16.0 - +

Other references

Other References:

Data collection for this site was done in conjunction with the progressive soil surveys within the Southern Desertic Basins, Plains and Mountains, Major Land Resource Areas of New Mexico. This site has been mapped and correlated with soils in the following soil surveys. Sierra County Dona Ana County Grant County Hidalgo County Luna County Otero County

Characteristic Soils Are:

Bluepoint loamy fine sand
Pintura loamy fine sand
Yturbide loamy sand
Other Soils included are:
Wink loamy fine sand, more than 5"surface
Hucco loamy fine sand, more than 5"surface
Yturbide gravelly sand

Contributors

Don Sylvester
Dr. Brandon Bestelmeyer

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
