

Ecological site F043AY567ID

Vitrandid Hills and Mountains 30-45" PZ Frigid Western Bitterroot Foothills

Last updated: 10/14/2020

Accessed: 05/11/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 043A–Northern Rocky Mountains

Major Land Resource Area (MLRA): 043A–Northern Rocky Mountains

Description of MLRAs can be found in: United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

Available electronically at: http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2_053624#handbook

LRU notes

Most commonly found in LRU 43A09 (Western Bitterroot Foothills). Also found in adjacent areas of 43A11 (Bitterroot Metasedimentary Zone). Climate parameters were obtained from PRISM and other models for the area. Landscape descriptors are derived from USGS DEM products and their derivatives.

Classification relationships

Relationship to Other Established Classifications:

United States National Vegetation Classification (2008) – A3362 Grand fir – Douglas-fir Central Rocky Mt. Forest and Woodland Alliance.

Washington Natural Heritage Program. Ecosystems of Washington State, A Guide to Identification, Rocchio and Crawford, 2015 –

Northern Rocky Mt. Mesic Montane Mixed Conifer Forest.

Description of Ecoregions of the United States, USFS PN # 1391, 1995 - M333 Northern Rocky Mt. Forest-Steppe-Coniferous Forest-Alpine Meadow Province

Level III and IV Ecoregions of WA, US EPA, June 2010 -15w Western Selkirk Maritime Forest. 15y Selkirk Mountains. 15u Inland Maritime Foothills and Valleys.

This ecological site includes the following USDA Forest Service Plant Associations: ABGR/ACGLD/CLUN, Forest Plant Associations of the Colville Nat. Forest (Williams et. al. 1995) and ABGR/CLUN (Forest Habitat Types of N. Idaho, A Second Approximation, USFS Revised 1991.

Ecological site concept

This ES group is distinguished by an overstory of grand fir and Douglas-fir. Understory shrubs can include *Acer glabrum*, *Linnaea borealis*, *Menziesia ferruginea*, *Physocarpus malvaceus*, *Spiraea betulifolia*, *Symphoricarpos*

occidentalis, and *Vaccinium membranaceum*. Herbaceous layers may be graminoid- or forb-dominated and may include *Bromus vulgaris*, *Calamagrostis rubescens*, *Carex geyeri*, *Clintonia uniflora*, *Coptis occidentalis*, *Cornus canadensis*, *Linnaea borealis*, and *Trautvetteria caroliniensis*. It occurs on foothills, mountainsides, and canyon walls. These soils have developed in highly mixed Mazama tephra deposits and alluvium, or colluvium and residuum from granite or metamorphic rock. The soils are very deep and have adequate available water capacity to a depth of 40 inches. The soils are well drained. This ES group fits into the National Vegetation Standard's Grand Fir - Douglas-fir Central Rocky Mountain Forest & Woodland Alliance and Washington State's Natural Heritage Program's Northern Rocky Mt. Mesic Montane Mixed Conifer Forest.

Table 1. Dominant plant species

| | |
|------------|--|
| Tree | (1) <i>Abies grandis</i> (2) <i>Pseudotsuga menziesii</i> var. <i>glauca</i> |
| Shrub | (1) <i>Rosa gymnocarpa</i> (2) <i>Linnaea borealis</i> ssp. <i>longiflora</i> |
| Herbaceous | (1) <i>Clintonia uniflora</i> (2) <i>Coptis occidentalis</i> |

Physiographic features

Physiographic Features

Landscapes: Foothills, Mountains, Plateau

Landform: hillslopes, mountain slopes, structural bench

Elevation (m): Total range = 490 to 1540 m

(1,605 to 5,050 feet)

Central tendency = 750 to 1065 m

(2,460 to 3,495 feet)

Slope (percent): Total range = 0 to 90 percent

Central tendency = 20 to 45 percent

Aspect: Total range: 85-175-360

Central tendency: 120-175-245

Table 2. Representative physiographic features

| | |
|--------------------|---|
| Landforms | (1) Mountains > Mountain slope (2) Foothills > Hillslope (3) Plateau > Structural bench |
| Flooding frequency | None |
| Ponding frequency | None |
| Elevation | 2,460–3,495 ft |
| Slope | 20–45% |
| Water table depth | 80 in |
| Aspect | SE, S, SW |

Table 3. Representative physiographic features (actual ranges)

| | |
|--------------------|----------------|
| Flooding frequency | None |
| Ponding frequency | None |
| Elevation | 1,605–5,050 ft |
| Slope | 0–90% |
| Water table depth | 80 in |

Climatic features

Climatic Features

Frost-free period (days): Total range = 70 to 125 days
Central tendency = 85 to 110 days

Mean annual precipitation: Total range = 610 to 1295 mm
(24 to 51 inches)
Central tendency = 770 to 1060mm
(30 to 42 inches)

MAAT (C): Total range = 6.0 to 10.2
(43 to 50 F)
Central tendency = 7.1 to 8.3
(45 to 47 F)

Climate Stations: none

Influencing water features

Water Table Depth:
>200 cm
(>80 inches)

Flooding:
Frequency: None
Duration: None

Ponding:
Frequency: None
Duration: None

Soil features

Representative Soil Features

This ecological subsite is associated with several soil series (e.g. Aldermand, Keeler, Bigtalk, Statemeadow, and Kingspeak). The soil components are Vitrandic Eutrudepts, Vitrandic Hapludalfs, and Typic Eutrudepts. These soils have developed in highly mixed Mazama tephra deposits and alluvium, or colluvium and residuum from granite or metamorphic rock. The soils are very deep and have adequate available water capacity to a depth of 40 inches. The soils are well drained.

Table 4. Representative soil features

| | |
|-----------------|--|
| Parent material | (1) Volcanic ash (2) Alluvium (3) Colluvium–granite (4) Colluvium–metamorphic rock (5) Residuum–granite (6) Residuum–metamorphic rock |
| Surface texture | (1) Ashy silt loam (2) Ashy loam (3) Loam |
| Drainage class | Well drained |

| | |
|---|-----------------|
| Permeability class | Moderately slow |
| Depth to restrictive layer | 80 in |
| Surface fragment cover >3" | 0% |
| Available water capacity (0-40in) | 7.5 in |
| Calcium carbonate equivalent (0-60in) | 0% |
| Electrical conductivity (0-60in) | 0 mmhos/cm |
| Soil reaction (1:1 water) (0-60in) | 6 |
| Subsurface fragment volume <=3" (10-60in) | 5% |
| Subsurface fragment volume >3" (10-60in) | 0% |

Table 5. Representative soil features (actual values)

| | |
|---|-----------------------------|
| Drainage class | Well drained |
| Permeability class | Moderately slow to moderate |
| Depth to restrictive layer | 80 in |
| Surface fragment cover >3" | 0% |
| Available water capacity (0-40in) | 5.4–7.9 in |
| Calcium carbonate equivalent (0-60in) | 0% |
| Electrical conductivity (0-60in) | 0 mmhos/cm |
| Soil reaction (1:1 water) (0-60in) | 5–7.3 |
| Subsurface fragment volume <=3" (10-60in) | 0–25% |
| Subsurface fragment volume >3" (10-60in) | 0–35% |

Ecological dynamics

Ecological Dynamics of the Site

This grand fir site is on the moister end of the grand fir series and can have many tree species occurring depending on disturbance and seed source. Douglas-fir is the main seral species and will occur in mature stands with grand fir with fire exclusion. Engelmann spruce can also be present in fire excluded mature stands. With stand replacing or mixed severity disturbance Douglas-fir, western larch, ponderosa pine, western white pine, and lodgepole pine can occur in mixed stands. Quaking aspen, paper birch, and black cottonwood can be present in early to mid-seral stands.

Key understory species include Douglas maple, ninebark, oceanspray, serviceberry, snowberry, twinflower, honeysuckle, Oregon grape, rose, thimbleberry, pachistima, Scouler willow, spirea, pinegrass, brides bonnet, pathfinder, sweetroot, starry solomonplume, western princes pine, and white hawkweed.

Root rot and beetle kill will occur in the mature grand fir – Douglas fir stands. Western white pine once would have been more prevalent in the seral to mature stands, however, blister rust has made it a minor component of current

stands. Low intensity frequent fires will favor more open western larch, Douglas-fir, and ponderosa pine stands. Mixed severity fires will create a mosaic of mixed stands of all tree species. Severe fires may create ceanothus brush fields with tree regeneration slow to reclaim the site.


State and transition model

State and Transition Diagram

Ecological Site

Frigid Udic Loamy Foothills/Mountainsides (Grand Fir Moist Herb)

Abies grandis/*Clintonia uniflora* (Grand Fir / Bride's Bonnet)



State 1

Reference State

This state is dependent on fire severity and time intervals of fire. Many tree species will grow on this moist grand fir site. Douglas-fir, western larch, western white pine, ponderosa pine, lodgepole pine, Engelmann spruce, and grand fir are the major conifer species. Quaking aspen, paper birch and black cottonwood can also be present. This ecological site has good forest productivity for Douglas-fir, western larch, and ponderosa pine. In N. Idaho western white pine would have good productivity. Low severity frequent fires create open stands of larch, Douglas-fir, ponderosa pine, and lodgepole pine. Mixed severity fires create a mosaic of a variety of tree species and different age groups. Tree regeneration after stand replacing fires will be variable depending on seed source. Western larch, western white pine, Douglas-fir, ponderosa pine, and lodgepole pine will be the main species. Western white pine used to play a major role in regeneration, however, the white pine blister rust has reduced its occurrence in regenerating stands. Western larch, Douglas-fir, and ponderosa pine are more prevalent. Reoccurring severe fires in the stand initiation phase can create shrub fields of ceanothus spp., and a variety of other shrubs. Fire exclusion for long periods allow the forest to move to an all aged grand fir, Douglas-fir, and Engelmann spruce stand. Root rot and defoliating insects then becomes are concern in the grand fir and Douglas-fir causing much downed wood and fuel loads.

Characteristics and indicators. Plant List Overstory Trees Douglas-fir (*Pseudotsuga menziesii*) Western Larch (*Larix occidentalis*) Grand fir (*Abies grandis*) Understory trees Grand fir (*Abies grandis*) Understory Shrubs Douglas

maple (*Acer glabrum* Douglasii) Western prince's pine (*Chimaphila umbellata*) Serviceberry (*Amelanchier alnifolia*) Oregon grape (*Berberis aquifolium*) Snowberry (*Symphoricarpos albus*) Twinflower (*Linnaea borealis*) Pachistima (*Pachistima myrsinites*) Ninebark (*Physocarpus malvaceus*) Oceanspray (*Holodiscus discolor*) Shiny leaf spirea (*Spiraea betulifolia*) Big huckleberry (*Vaccinium membranaceum*) (cooler sites) Scouler willow (*Salix scouleriana*) Utah honeysuckle (*Lonicera utahensis*) Mountain ash (*Sorbus scopulina*) (cooler sites) Understory Herbs Pathfinder (*Adenocaulon bicolor*) Sweetroot (*Osmorhiza chilensis*) Starry solomonplume (*Smilacina stellata*) Queencup beadlely (*Clintonia uniflora*) White hawkweed (*Hieracium albiflorum*) Western meadowrue (*Thalictrum occidentale*)

Dominant plant species

- Rocky Mountain Douglas-fir (*Pseudotsuga menziesii* var. *glauca*), tree
- western larch (*Larix occidentalis*), tree
- grand fir (*Abies grandis*), tree
- Rocky Mountain maple (*Acer glabrum*), shrub
- pipsissewa (*Chimaphila umbellata*), shrub
- Saskatoon serviceberry (*Amelanchier alnifolia*), shrub
- hollyleaved barberry (*Mahonia aquifolium*), shrub
- common snowberry (*Symphoricarpos albus*), shrub
- longtube twinflower (*Linnaea borealis* ssp. *longiflora*), shrub
- Oregon boxleaf (*Paxistima myrsinites*), shrub
- mallow ninebark (*Physocarpus malvaceus*), shrub
- oceanspray (*Holodiscus discolor*), shrub
- white spirea (*Spiraea betulifolia*), shrub
- thinleaf huckleberry (*Vaccinium membranaceum*), shrub
- Scouler's willow (*Salix scouleriana*), shrub
- common lomatium (*Lomatium utriculatum*), shrub
- Greene's mountain ash (*Sorbus scopulina*), shrub

Community 1.1

Reference Phase

Mixed severity fires create a mosaic of mixed stands of seral species mixed with grand fir. Regeneration could include all the species depending on exposed mineral soil and seed source. Western larch, ponderosa pine, Douglas-fir, and lodgepole pine would be first to regenerate. Western white pine will also be present, but not as prevalent as before due to blister rust. Grand fir will be present, however will come underneath the seral species in time. Quaking aspen can establish after these mixed fires and be a component in the stand. Age: 80+ years

Dominant plant species

- grand fir (*Abies grandis*), tree
- western larch (*Larix occidentalis*), tree
- Rocky Mountain Douglas-fir (*Pseudotsuga menziesii* var. *glauca*), tree
- hollyleaved barberry (*Mahonia aquifolium*), shrub
- common snowberry (*Symphoricarpos albus*), shrub
- longtube twinflower (*Linnaea borealis* ssp. *longiflora*), shrub
- Oregon boxleaf (*Paxistima myrsinites*), shrub
- mallow ninebark (*Physocarpus malvaceus*), shrub
- oceanspray (*Holodiscus discolor*), shrub
- white spirea (*Spiraea betulifolia*), shrub
- thinleaf huckleberry (*Vaccinium membranaceum*), shrub
- Scouler's willow (*Salix scouleriana*), shrub
- common lomatium (*Lomatium utriculatum*), shrub
- Greene's mountain ash (*Sorbus scopulina*), shrub

Community 1.2

Stand Initiation

Herbs and shrubs reestablish on site. Ceanothus species can dominate if fire caused soil degradation. Tree regeneration can be mixed or dominated by western larch and lodgepole pine. Ponderosa pine and Douglas-fir can

also be present. White pine was a major factor in regeneration before the blister rust reduced its significance. Age: 0 – 20+ years

Community 1.3

Stem Exclusion

Dense stands of mixed seral species or dominated by larch or lodgepole pine. Douglas-fir, ponderosa pine, and western white pine, can be in the stand. Age: 100+ years

Dominant plant species

- western larch (*Larix occidentalis*), tree
- lodgepole pine (*Pinus contorta* var. *latifolia*), tree
- Rocky Mountain Douglas-fir (*Pseudotsuga menziesii* var. *glauca*), tree
- ponderosa pine (*Pinus ponderosa*), tree
- western white pine (*Pinus monticola*), tree

Community 1.4

Frequent Low Intensity Fires

Frequent low severity fires create an open stand of mature Douglas-fir, Western larch, ponderosa pine, and lodgepole pine. This condition is perpetuated with continual ground fires. Age: 30 – 70 years

Dominant plant species

- Rocky Mountain Douglas-fir (*Pseudotsuga menziesii* var. *glauca*), tree
- western larch (*Larix occidentalis*), tree
- lodgepole pine (*Pinus contorta* var. *latifolia*), tree
- ponderosa pine (*Pinus ponderosa*), tree

Pathway 1.1A

Community 1.1 to 1.2

Stand replacing fire back to the herbs and shrub stage. Regeneration dependent on seed source and brush competition.

Pathway 1.2A

Community 1.2 to 1.3

Tree regeneration grows into dense pole stand. Species can be mixed or dominated by larch or lodgepole.

Pathway 1.3B

Community 1.3 to 1.1

Mixed severity fires create patchy mosaic as in reference plant community phase

Pathway 1.3A

Community 1.3 to 1.2

Stand replacing fire.

Pathway 1.3C

Community 1.3 to 1.4

Frequent low severity fires create open pole stands.

Pathway 1.4B

Community 1.4 to 1.1

Fire interval lengthens, and tree regeneration becomes established. Mixed severity fires create patchy mosaic of seral species and grand fir.

State 2

Fire Exclusion

An all aged stand of grand fir and Douglas-fir with Engelmann spruce mixed in occurs when fire interval is extended 100 years or more. Grand fir will dominate the regeneration in the understory. In the drier areas of this site root rot and defoliating insects can be a problem. Snags and downed wood will be common in these areas. This state will be susceptible to stand replacing fires.

Dominant plant species

- grand fir (*Abies grandis*), tree
- Rocky Mountain Douglas-fir (*Pseudotsuga menziesii* var. *glauca*), tree
- Engelmann spruce (*Picea engelmannii*), tree

Community 2.1

Fire Exclusion

An all aged stand of grand fir and Douglas-fir with Engelmann spruce mixed in occurs when fire interval is extended 100 years or more. Grand fir will dominate the regeneration in the understory. In the drier areas of this site root rot and defoliating insects can be a problem. Snags and downed wood will be common in these areas. This state will be susceptible to stand replacing fires.

Dominant plant species

- grand fir (*Abies grandis*), tree
- Rocky Mountain Douglas-fir (*Pseudotsuga menziesii* var. *glauca*), tree
- Engelmann spruce (*Picea engelmannii*), tree

State 3

Shrubfields

Successive severe ground fires create shrub fields for long time periods. Ceanothus species, ninebark, oceanspray, snowberry, spirea, Douglas maple, Scouler willow and serviceberry will dominate preventing tree regeneration.

Dominant plant species

- snowbrush ceanothus (*Ceanothus velutinus*), shrub
- redstem ceanothus (*Ceanothus sanguineus*), shrub
- mallow ninebark (*Physocarpus malvaceus*), shrub
- oceanspray (*Holodiscus discolor*), shrub
- common snowberry (*Symphoricarpos albus*), shrub
- white spirea (*Spiraea betulifolia*), shrub
- Rocky Mountain maple (*Acer glabrum*), shrub
- Scouler's willow (*Salix scouleriana*), shrub
- Saskatoon serviceberry (*Amelanchier alnifolia*), shrub

Community 3.1

Shrubfields

Successive severe ground fires create shrub fields for long time periods. Ceanothus species, ninebark, oceanspray, snowberry, spirea, Douglas maple, Scouler willow and serviceberry will dominate preventing tree regeneration.

Dominant plant species

- snowbrush ceanothus (*Ceanothus velutinus*), shrub

- redstem ceanothus (*Ceanothus sanguineus*), shrub
- mallow ninebark (*Physocarpus malvaceus*), shrub
- oceanspray (*Holodiscus discolor*), shrub
- common snowberry (*Symphoricarpos albus*), shrub
- white spirea (*Spiraea betulifolia*), shrub
- Rocky Mountain maple (*Acer glabrum*), shrub
- Scouler's willow (*Salix scouleriana*), shrub
- Saskatoon serviceberry (*Amelanchier alnifolia*), shrub

Transition T1A

State 1 to 2

100+ years of fire exclusion will move the site to a mature grand fir – Douglas-fir site.

Transition T1B

State 1 to 3

Successive severe burns kill regeneration and maintaining sprouting shrubs dominance.

Restoration pathway R2A

State 2 to 1

Overstory management with ground burning will favor seral species.

Restoration pathway R3A

State 3 to 1

Careful site selection and fire control allow tree regeneration to establish

Additional community tables

References

. 1998. NRCS National Forestry Manual.

. 2017. NRCS Soil and Site Index data for NE WA and N. Idaho.

Cooper, S.V., K.E. Neiman, R. Steele, and D.W. Roberts. 1991. Forest Habitat types of Northern Idaho, A Second Approximation.

Finklin, A.I. 1983. Climate of Priest River Experimental Forest, northern Idaho. Gen. Tech. Rep. INT-159. U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station, Ogden, UT. 53.

Smith and Fischer. 1997. Fire Ecology of the Forest Habitat Types of Northern Idaho.

Williams, C.K., B.F. Kelley, B.G. Smith, and T.R. Lillybridge. October, 1995. Forested Plant Associations of the Colville National Forest.

Zack, A. 1997. Biophysical Classification- Habitat Groups and Description of Northern Idaho and Northwestern Montana, Lower Clarkfork and Adjacent Areas..

Approval

Curtis Talbot, 10/14/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

| | |
|---|-------------------|
| Author(s)/participant(s) | |
| Contact for lead author | |
| Date | 05/11/2025 |
| Approved by | Curtis Talbot |
| Approval date | |
| Composition (Indicators 10 and 12) based on | Annual Production |

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
