

Ecological site R043AP805MT Shallow Grassland Group

Last updated: 9/08/2023
Accessed: 05/12/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 043A–Northern Rocky Mountains

This MLRA is located in Montana (43 percent), Idaho (34 percent), and Washington (23 percent). It makes up about 31,435 square miles (81,460 square kilometers). It has no large cities or towns. It has many national forests, including the Okanogan, Colville, Kootenai, Lolo, Flathead, Coeur d'Alene, St. Joe, Clearwater, and Kaniksu National Forests.

This MLRA is in the Northern Rocky Mountains Province of the Rocky Mountain System. It is characterized by rugged, glaciated mountains; thrust- and block-faulted mountains; and hills and valleys. Steep-gradient rivers have cut deep canyons. Natural and manmade lakes are common.

The major Hydrologic Unit Areas (identified by four-digit numbers) that make up this MLRA are: Kootenai-Pend Oreille-Spokane (1701), 67 percent; Upper Columbia (1702), 18 percent; and Lower Snake (1706), 15 percent. Numerous rivers originate in or flow through this area, including, the Sanpoil, Columbia, Pend Oreille, Kootenai, St. Joe, Thompson, and Flathead Rivers.

This area is underlain primarily by stacked slabs of layered sedimentary or metasedimentary bedrock. The bedrock formations range from Precambrian to Cretaceous in age. The rocks consist of shale, sandstone, siltstone, limestone, argillite, quartzite, gneiss, schist, dolomite, basalt, and granite. The formations have been faulted and stacked into a series of imbricate slabs by regional tectonic activity. Pleistocene glaciers carved a rugged landscape that includes sculpted hills and narrow valleys filled with till and outwash. Continental glaciation overrode the landscape in the northern half of the MLRA while glaciation in the southern half was confined to montane settings.

The average annual precipitation is 25 to 60 inches (635 to 1,525 millimeters) in most of this area, but it is as much as 113 inches (2,870 millimeters) in the mountains and is 10 to 15 inches (255 to 380 millimeters) in the western part of the area. Summers are dry. Most of the precipitation during fall, winter, and spring is snow. The average annual temperature is 32 to 51 degrees F (0 to 11 degrees C) in most of the area, decreasing with elevation. In most of the area, the freeze-free period averages 140 days and ranges from 65 to 215 days. It is longest in the low valleys of Washington, and it decreases in length with elevation. Freezing temperatures occur every month of the year on high mountains, and some peaks have a continuous cover of snow and ice.

The dominant soil orders in this MLRA are Andisols, Inceptisols, and Alfisols. Many of the soils are influenced by Mount Mazama ash deposits. The soils in the area have a frigid or cryic soil temperature regime; have an ustic, xeric, or udic soil moisture regime; and dominantly have mixed mineralogy. They are shallow to very deep, are very poorly drained to well drained, and have most of the soil texture classes. The soils at the lower elevations include Udivitrands, Vitrixerands and Haplustalfs. The soils at the higher elevations include Dystrocrypts, Eutrocrypts, Vitricryands, and Haplocryalfs. Cryorthents, Cryepts, and areas of rock outcrop are on ridges and peaks above timberline

This area is in the northern part of the Northern Rocky Mountains. Grand fir, Douglas-fir, western red cedar, western hemlock, western larch, lodgepole pine, subalpine fir, ponderosa pine, whitebark pine, and western white pine are the dominant overstory species, depending on precipitation, temperature, elevation, and landform aspect. The understory vegetation varies, also depending on climatic and landform factors. Some of the major wildlife species in this area are whitetailed deer, mule deer, elk, moose, black bear, grizzly bear, coyote, fox, and grouse. Fish, mostly in the trout and salmon families, are abundant in streams, rivers, and lakes.

More than one-half of this area is federally owned and administered by the U.S. Department of Agriculture, Forest Service. Much of the privately-owned land is controlled by large commercial timber companies. The forested areas are used for wildlife habitat, recreation, watershed, livestock grazing, and timber production. Meadows provide summer grazing for livestock and big game animals. Less than 3 percent of the area is cropland.

Ecological site concept

- Site does not receive additional water
- This site has lower productivity, resistance and resilience to disturbance than the similar site Upland Grassland due to shallow soils.
- Dominant Cover: Grassland;Midstatured bunchgrasses rough fescue and bluebunch wheatgrass generally dominant (bluebunch, rough fescue, Idaho fescue, trace cover of Prairie junegrass, Sandberg’s bluegrass), forbs include Indian blanketflower, boreal bedstraw, prairie smoke, twin arnica, rosy pussytoes, silver lupine, Hood's phlox, stoneseed, western sagewort. Shrubs are a relatively small component. Average production is 1075 dry pounds per acre (1000-1200).
- Soils are
 - o Generally not saline or saline-sodic or limy (limited extent)
 - o Shallow (less than 50cm deep to bedrock, lithic, or paralithic root restrictive layer)
 - o Not ashy or medial textural family
 - o Typically less than 5% stone and boulder cover (<15% max)
- Soil surface texture gravelly or cobbly loam in surface mineral 4”
- Parent material is colluvium over residuum weathered from welded tuff or metasedimentary rock
- Drainage class is well drained; no flooding frequency
- Site Landform: hills, glaciated mountain slopes
- Moisture Regime: xeric
- Temperature Regime: frigid
- Elevation Range: 2900-5000 ft
- Slope: 15-50%

Associated sites

R043AP810MT	Upland Grassland Group These sites are associated in that they both reside in hills at elevation range of 2700 to 5000 feet on moderate slopes of 10 to 30 percent. The reference community for both sites has the perennial bunchgrasses and well drained loamy soils, though R043AP805MT has shallow depth.
-------------	---

Similar sites

R043AP810MT	Upland Grassland Group These sites are similar in that they both reside in hills at elevation range of 2700 to 5000 feet on moderate slopes of 10 to 30 percent. The reference community for both sites has the perennial bunchgrasses and well drained loamy soils, though R043AP805MT has shallow depth.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Pseudoroegneria spicata</i> (2) <i>Festuca campestris</i>

Physiographic features

- Site Landform: hills, glaciated mountain slopes
- Elevation Range: 2900-5000 ft
- Slope: 15-50%

Table 2. Representative physiographic features

Landforms	(1) Mountains > Hill (2) Mountains > Mountain slope
Elevation	884–1,524 m
Slope	15–50%
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

- Moisture Regime: xeric
- Temperature Regime: frigid
- Representative Value (RV) of range of Mean Annual Precipitation: 14-25 inches
- Representative Value (RV) of range of Mean Average Annual Temperature: 43-46 degrees
- Representative Value (RV) of range of Frost Free Days: 70-100 days

Table 3. Representative climatic features

Frost-free period (characteristic range)	2-86 days
Freeze-free period (characteristic range)	58-131 days
Precipitation total (characteristic range)	533-610 mm
Frost-free period (actual range)	1-87 days
Freeze-free period (actual range)	44-132 days
Precipitation total (actual range)	508-711 mm
Frost-free period (average)	49 days
Freeze-free period (average)	99 days
Precipitation total (average)	584 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) POLEBRIDGE 1 N [USC00246618], Essex, MT
- (2) POLEBRIDGE [USC00246615], Essex, MT
- (3) WEST GLACIER [USC00248809], Kalispell, MT
- (4) WHITEFISH [USC00248902], Whitefish, MT
- (5) LIBBY 32 SSE [USC00245020], Libby, MT
- (6) LINDBERGH LAKE [USC00245043], Seeley Lake, MT
- (7) YAAK 9NNE [USC00249187], Troy, MT

Influencing water features

- Site does not receive additional water

Wetland description

DOES NOT APPLY

Soil features

- Soils are
 - o Generally not saline or saline-sodic or limy (limited extent)
 - o Shallow (less than 50cm deep to bedrock, lithic, or paralithic root restrictive layer)
 - o Not ashy or medial textural family
 - o Typically less than 5% stone and boulder cover (<15% max)
- Soil surface texture gravelly or cobbly loam in surface mineral 4"
- Parent material is colluvium over residuum weathered from welded tuff or metasedimentary rock
- Drainage class is well drained; no flooding frequency

Table 4. Representative soil features

Parent material	(1) Colluvium (2) Residuum
Surface texture	(1) Gravelly loam (2) Cobbly loam
Drainage class	Well drained
Soil depth	51 cm

Ecological dynamics

1.1 Midstatured bunchgrasses dominant (bluebunch, rough fescue, Idaho fescue, trace cover of Prairie junegrass, Sandberg's bluegrass), Shrubs are a relatively small component.

1.1a extended drought, improper grazing, climate change, catastrophic fire (limited on this site)

1.2 Midstatured bunchgrasses subdominant to increaser bunchgrasses such as needle-and-thread or Idaho fescue.

Shrubs increasing, clubmoss possible (limited extent), mat forming forbs increasing

1.2a proper grazing management, favorable growing conditions, time

T1A poor post settlement grazing (late 1800's), drought with improper grazing, multiple spring grazing, fire suppression

T1B sodbusting, introduction of tame pasture species and other invasive plants, overgrazing, drought, heavy human disturbance, extreme fire (multiple years or very intense)

T1C poor post settlement grazing (late 1800's), drought with improper grazing, multiple spring grazing and/or long term overgrazing, fire suppression

T3A sodbusting, invasive plants, overgrazing, extended drought, adjacent to construction or disturbance event

2.1 Mixed grass dominated site (needle-and-thread and Idaho fescue), midstatured bunchgrasses existent under shrub canopy, possible conifer encroachment, forbs (scarlet globemallow, hood's phlox, mat forming forbs) and shrubs increase (broom snakeweed, big sagebrush)

2.1a improper grazing management, drought, fire, climate change

2.2 Needle-and-thread and Idaho fescue losing dominance to Sandberg bluegrass and prairie Junegrass.

Decreaser bunchgrasses very rare and limited under shrub canopy. Broom snakeweed and Fringed sagewort beginning to replace shrub component

2.2a proper grazing management, time, Integrated Pest Management, brush management

3.1 Shortgrass State lacks midstatured bunchgrasses, Sandberg bluegrass and Junegrass dominant grasses, increaser shrubs nearly replace larger shrub species. Remaining larger shrub species heavily hedged.

T2A overgrazing, introduction of weeds, drought, heavy human disturbance

R2A fire, range seeding, timely moisture, proper grazing management, IPM

R3B Possibly not feasible, range seeding, time, proper grazing management, IPM

T2B sodbusting, introduction of tame pasture species and other invasive plants, overgrazing, extended drought, adjacent to construction or disturbance event, extreme fire (multiple years or very intense)

4.1 Invaded State may resemble reference however contains noxious or invasive weeds such as cheatgrass or knapweed. Conifer encroachment common.


R3A range seeding, time, proper grazing management, IPM

R4A IPM, timely moisture, grazing management, brush management, range seeding

R4B IPM, range seeding, timely moisture, grazing management, brush management, range seeding

State and transition model

Shallow Grassland R043AP805MT



Animal community

Big game wildlife and livestock use this site extensively. Since this site is composed primarily of deep rooted perennial native bunchgrasses, grazing during non-sensitive time periods that is rotational in nature and allows for adequate rest is needed. The shallow depth of the soils associated with this site may lower the resistance and resilience of this site to heavy grazing.

Hydrological functions

Shifts in plant composition away from reference in which native bunchgrasses dominate will result in less plant water availability of the site in general.

Recreational uses

HIKING, BIKING, PHOTOGRAPHY

Wood products

NO USES

Contributors

jay skovlin
stephanie shoemaker

Approval

Kirt Walstad, 9/08/2023

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators

are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	12/18/2020
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
