

Ecological site R043AY510ID

Poorly Drained Floodplains 23-25" PZ Mesic Columbia-Colville Valleys

Last updated: 10/15/2020
Accessed: 05/11/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 043A–Northern Rocky Mountains

Major Land Resource Area (MLRA): 043A–Northern Rocky Mountains

Description of MLRAs can be found in: United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

LRU notes

Major land resource area (MLRA): 043A-Northern Rocky Mountains

Modal LRU – 43A03 - Columbia-Colville Valleys

This LRU is composed predominantly of glaciated valleys and foothills. The soils tend to be loamy to sandy Haploxerepts and Haploxerolls with mixed or distinct ash surfaces. Recent alluvium, till, outwash and residuum from granitic or metamorphic rock are the dominant parent materials. Soil climate is a mesic or frigid temperature regime and xeric moisture regime with average annual precipitation around 490 mm (19 inches).

Others where occurring: 44A01 Spokane-Rathdrum Outwash Plains
43A04 Selkirk Mountains

Classification relationships

This ES fits into the National Vegetation Standard's and is most closely related to Washington State's Natural Heritage Program's. (Compare to previous Washington range site: R044XY601WA, WET MEADOW 16-24 PZ)

Ecological site concept

Ecological Site Concept:

This ES is found on poorly to somewhat poorly drained, loamy sites on floodplains. A water table is within 75 cm of the surface at some time during the May-Oct period. They have mesic temperatures and a plant community dominated by sedges and grasses.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified

Herbaceous	(1) <i>Carex</i> (2) <i>Calamagrostis stricta</i>
------------	--

Physiographic features

Physiographic Features

This ecological site occurs mainly on floodplains and drainageways. Parent materials are mixed alluvium with a mantle of volcanic ash and loess.

Landscapes: Valleys and scablands

Landforms: depressions, floodplains, drainageways

Elevation:

Total range = 390 to 1040 m

(1,285 to 3,405 feet)

Central tendency = 560 to 750 m

(1,835 to 2,465 feet)

Slope (percent):

Total range = 0 to 3 percent

Central tendency = 1 to 2 percent

Water Table Depth:

0 to 92 cm; median = 31 cm

(0 to 36 inches; median = 12inches)

Flooding:

Frequency: occasional to frequent

Duration: brief to long

Ponding:

Frequency: None to frequent

Duration: None to long

Aspect: NA

Table 2. Representative physiographic features

Landforms	(1) Valley > Flood plain (2) Valley > Depression (3) Scabland > Drainageway
Flooding duration	Not specified
Flooding frequency	Not specified
Ponding duration	Not specified
Ponding frequency	Not specified
Elevation	1,835–2,465 ft
Slope	1–2%
Water table depth	12 in
Aspect	Aspect is not a significant factor

Table 3. Representative physiographic features (actual ranges)

Flooding duration	Brief (2 to 7 days) to long (7 to 30 days)
Flooding frequency	Occasional to frequent

Ponding duration	Long (7 to 30 days)
Ponding frequency	None to frequent
Elevation	1,285–3,405 ft
Slope	0–3%
Water table depth	0–36 in

Climatic features

Climatic Features

During the spring and summer, a circulation of air around a high-pressure center brings a prevailing westerly and northwesterly flow of comparatively dry, cool and stable air into the region. As the air moves inland, it becomes warmer and drier which results in a dry season beginning in the late spring and reaching a peak in mid-summer. In the fall and winter, a circulation of air around two pressure centers over the ocean brings a prevailing southwesterly and westerly flow of air into the Pacific Northwest. This air from over the ocean is moist and near the temperature of the water. Condensation occurs as the air moves inland over the cooler land and rises along the windward slopes of the mountains or highlands. This results in a wet season beginning in October, reaching a peak in winter, then gradually decreasing in the spring.

The elevation within the LRU varies from approximately 1,000 feet in the lower river valleys to about 4,900 feet on low mountains and foothills. The annual precipitation increases from 9 inches in the valleys to over 38 inches in the higher terrain. Winter season snowfall averages about 43 inches. Both rainfall and snowfall increase with elevation. Snow can be expected after the first of November and to remain on the ground from the first of December until March or April.

In January, the average maximum temperature is near 29° F and the minimum temperature is 14° F. Minimum temperatures from -5° to -20°F are recorded almost every winter and temperatures ranging to -25° F have been recorded. In July, the average maximum temperature is 81° and the minimum temperature 47° F. Maximum temperatures reach 100° F on a few afternoons each summer and temperatures as high as 108° F have been recorded. The average date of the last freezing temperatures can be expected by early-June and after early-September in the warmer areas.

(Compiled from WRCC: Climate of Washington and available station data)

Frost-free period (days):

Total range = 100 to 140 days

Central tendency = 115 to 125 days

Mean annual precipitation (cm):

Total range = 360 to 855 mm

(14 to 34 inches)

Central tendency = 455 to 615 mm

(18 to 24 inches)

MAAT (C)

Total range = 5.8 to 9.9

(42 to 50 F)

Central tendency = 7.6 to 8.7

(46 to 48 F)

Climate stations: none

Influencing water features

Water Table Depth:

0 to 92 cm; median = 31 cm

(0 to 36 inches; median = 12inches)

Flooding:

Frequency: occasional to frequent

Duration: brief to long

Ponding:

Frequency: None to frequent

Duration: None to long

Soil features

Representative Soil Features

This ecological site is associated with the Cocolalla, Konner, Peone, and Wetthey components. The soils are Aquandic Endoaquepts, Aquandic Endoaquolls, Aquic Xerofluvents, Cumulic Endoaquolls. These soils have developed in mixed alluvium. Most sites have a surface influenced by volcanic ash and loess. Volcanic ash layers are mixed at lower elevations and form distinct, moderately thick layers at higher elevations.

Parent Materials:

Kind: volcanic ash, loess

Origin: mixed

Kind: alluvium

Origin: mixed

Surface Texture: (<2mm fraction)

(1) Ashy-Silt Loam

(2) Silty Clay Loam

(3) Loamy Sand

Surface Fragments

Table 4. Representative soil features

Parent material	(1) Volcanic ash (2) Loess (3) Alluvium
Surface texture	(1) Ashy silt loam (2) Silty clay loam (3) Loamy sand
Drainage class	Poorly drained
Permeability class	Moderate
Depth to restrictive layer	0 in
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	7.8 in
Calcium carbonate equivalent (0-60in)	0%
Electrical conductivity (0-60in)	0 mmhos/cm
Sodium adsorption ratio (0-60in)	0
Soil reaction (1:1 water) (0-60in)	6.7

Table 5. Representative soil features (actual values)

Drainage class	Somewhat poorly drained to poorly drained
Permeability class	Slow to very rapid
Depth to restrictive layer	0 in
Surface fragment cover <=3"	0–6%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	3–8.7 in
Calcium carbonate equivalent (0-60in)	0%
Electrical conductivity (0-60in)	0–2 mmhos/cm
Sodium adsorption ratio (0-60in)	0
Soil reaction (1:1 water) (0-60in)	5.6–7.3

Ecological dynamics

State and transition model

1 Reference State The reference community is a stabilized, persistent sedge dominated herbaceous community with mineral soils that withstands occasional flooding.

1.1 This plant community contains an abundance of sedges, with bluejoint reedgrass and shrubs including willow species on the periphery. This site is stable and typically can withstand occasional flooding.

1.1a Site experiences flooding that exceeds rooting strength of plant community or heavy grazing that reduces vegetation or fire or ice jam or beaver activity or debris flows or slumps that deposit soil and kills vegetation that returns understory community to the pioneering herbaceous community. In periods of extreme drought fire can occur. Fire return interval for surface fires is 50 years, severe fires may be 100 years. All of these disturbances return the site to the mineral soil dominated by pioneering herbaceous species phase.

1.2a Site becomes more stable over time, deeper rooted plants increase such as sedges and bluejoint reedgrass.

1.2 Pioneering herbaceous species establish on mineral soil deposited after flooding.

State 2 Increased nutrient load to system

T1a Increased nutrient load to system causes changes in plant composition and/or production. Site vulnerable to reed canarygrass invasion and dominance and site conversion, if present on site.

R1 Cessation of nutrient load to system, eradication of reed canarygrass stands and restoration efforts needed to restore community.

State 3 Fragmentation of site resulting in patches of non-connected patches of shrubs with altered hydrology, nutrient flow and vegetation propagules and wildlife dispersal altered.

T1b Fragmentation of the intact community and its hydrology and nutrient flows to numerous disconnected patches due to development, dams, extreme grazing practices or ungulate or recreation use.

R2 Improved grazing practices, altered ungulate use, dam and development removal and seeding of shrub species and other restoration practices.

State 4 Dry riparian site: Site is typically dominated by native grasses such as an understory of common snowberry, wood rose, black hawthorn, redosier dogwood, western meadowrue, false Solomons-seal, spreading sweetroot, largeleaf avens, sweetscented bedstraw, and blue wildrye. Site loses hydrology due to downcutting of stream or ditching/drainage or site, watershed alterations to hydrology that leads to site drying, extreme prolonged severe drought may cause as well.

T1c Human land uses that lower the water table including dams, roads, channeling, ditching or draining impact the site. Improper grazing can create trampling, hummocking, pugging and compaction of the soil resulting in potential loss of water at the site and accelerated stream downcutting. Improper grazing practices can also lead to changes in the nutrient cycling of the site. Drying of system as a result of loss of hydrology and increase in drier shrub species encroaching.

R3 Return of natural hydrology of site with restoration of dams, roads, other human development and/or prescribed grazing practices or reduced ungulate use or cessation of watershed factors that impact hydrological function of the site.

T3 T4 T5 T1e Introduction and dominance of non-native species and invasive species. Sites are invaded by noxious weeds or introduced pasture grasses. Pasture grasses may be planted or a result of invasion from neighboring sites. Improper grazing may be a trigger for invasion however flooding may transport seeds to freshly deposited alluvium.

State 5 Invaded State: This includes many non-native species that have come to dominate riparian areas such as cheatgrass (Poa pratensis), timothy (Phleum pratensis) and some native increaser species may include: Juncus arcticus, Iris missouriensis, Argemone anserina, and Dasiphora floribunda. Often sites are a combination both pasture grasses and invading weeds.

State 6 Overgrazed State: This state develops due to non-prescription, extreme grazing practices in which overgrazing of palatable species exists and increaser species gain dominance in the vegetation community.

T1d Overgrazing causes the plant community to change to increaser species.

R4 Sustainable grazing practices employed, adequate restoration of vegetation community with removal of weedy species, seeding of native palatable species.

T5a The overgrazed state transitions to the weed state with the establishment and dominance of noxious, weedy species.

Figure 1. Narrative

Approval

Curtis Talbot, 10/15/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/11/2025
Approved by	Curtis Talbot
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
