

Ecological site R046XS114MT Shallow (Sw) RRU 46-S 13-19 PZ

Last updated: 7/19/2023 Accessed: 05/12/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	G. Petersen
Contact for lead author	grant.petersen@usda.gov
Date	03/01/2020
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1.	Number and extent of rills: Rills are, generally, not present in the reference condition. If present, they will be short and
	inconspicuous on steep slopes exceeding 20%.

- 2. **Presence of water flow patterns:** Water flow patterns are, generally, not present in the reference condition but may be present on the steep, south facing slopes when runoff exceeds infiltration. These patterns will be short and infrequent across gentle slopes and increasingly more common on steeper slopes.
- 3. **Number and height of erosional pedestals or terracettes:** Pedestals are typically not evident in the reference condition. If present, they will be on slopes greater than 25 percent and associated with waterflow patterns.
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): Bare ground is 5-10%; however steeper, southerly aspects may express slightly higher bare ground of 10-15%.
- 5. Number of gullies and erosion associated with gullies: Gullies are not present in the reference condition.
- 6. Extent of wind scoured, blowouts and/or depositional areas: Wind scoured, or depositional areas are not evident in

	the reference condition.
7.	Amount of litter movement (describe size and distance expected to travel): Litter movement is infrequent across gentle slopes and increasingly more common on steeper slopes.
8.	Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): The average soil stability rating is 4-6 under plant canopies and plant interspaces. The A horizon is 2-4" inches thick.
9.	Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Soil Structure at the surface is typically moderate medium granular to weak fine subangular blocky. A Horizon should be 2-4 inches thick with color, when wet, typically ranging in Value of 5 or less and Chroma of 4 or less.
	Local geology may affect color, it is important to reference the Official Series Description (OSD) for characteristic range. https://soilseries.sc.egov.usda.gov/osdname.aspx
10.	Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Infiltration of the Shallow ecological site is moderate to moderately rapid. The site is well drained. An even distribution of mid stature grasses, 65-75% of site production, cool season rhizomatous grasses (10-15%) with shortgrass (15-20%), forbs (5-10), shrubs (10-15%) and trees/tall shrubs (0-1%)
11.	Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): A compaction layer is not present in the reference condition.
12.	Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to): Dominant: Mid-statured, cool season, perennial bunchgrasses (Primarily bluebunch wheatgrass, Columbia needlegrass, spike fescue)
	Sub-dominant: Shortgrass grasses & Grasslikes (Idaho fescue, needle and thread) = Forbs ≥ Shrubs >> trees/tall shrub: Other: Additional:
	Additional.
13.	Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Mortality in herbaceous species is not evident. Species with bunch growth forms may have some natural mortality in centers is 3% or less.
14.	Average percent litter cover (%) and depth (in): Total litter cover ranges from 40-60%.

- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): Average annual production is 1400. Low: 1225 High 1550. Production varies based on effective precipitation and natural variability of soil properties for this ecological site.
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Potential invasive (including noxious) species (native and non-native). Invasive species on this ecological site include (but not limited to) smooth brome, sulphur cinquefoil, dandelion, annual brome spp., spotted knapweed, salsify, leafy spurge, and ventenata,

Native species such as Rocky Mountain juniper, ponderosa pine, limber pine, Douglas fir, creeping juniper, lupine, broom snakeweed, Sandberg bluegrass, etc. when their populations are significant enough to affect ecological function, indicate site condition departure.

17. **Perennial plant reproductive capability:** In the reference condition, all plants are vigorous enough for reproduction either by seed or rhizomes in order to balance natural mortality with species recruitment.