

Ecological site F047XC541UT

High Mountain Very Steep Stony Loam (Douglas-fir)

Last updated: 2/05/2025
Accessed: 05/13/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 047X–Wasatch and Uinta Mountains

MLRA 47 occurs in Utah (86 percent), Wyoming (8 percent), Colorado (4 percent), and Idaho (2 percent). It encompasses approximately 23,825 square miles (61,740 square kilometers). The northern half of this area is in the Middle Rocky Mountains Province of the Rocky Mountain System. The southern half is in the High Plateaus of the Utah Section of the Colorado Plateaus Province of the Intermontane Plateaus. Parts of the western edge of this MLRA are in the Great Basin Section of the Basin and Range Province of the Intermontane Plateaus. The MLRA includes the Wasatch Mountains, which trend north and south, and the Uinta Mountains, which trend east and west. The steeply sloping, precipitous Wasatch Mountains have narrow crests and deep valleys. Active faulting and erosion are a dominant force in controlling the geomorphology of the area. The Uinta Mountains have a broad, gently arching, elongated shape. Structurally, they consist of a broadly folded anticline that has an erosion-resistant quartzite core. The Wasatch and Uinta Mountains have an elevation of 4,900 to about 13,500 feet (1,495 to 4,115 meters).

The mountains in this area are primarily fault blocks that have been tilted up. Alluvial fans at the base of the mountains are recharge zones for the basin fill aquifers. An ancient shoreline of historic Bonneville Lake is evident on the footslopes along the western edge of the area. Rocks exposed in the mountains are mostly Mesozoic and Paleozoic sediments, but Precambrian rocks are exposed in the Uinta Mountains. The Uinta Mountains are one of the few ranges in the United States that are oriented west to east. The southern Wasatch Mountains consist of Tertiary volcanic rocks occurring as extrusive lava and intrusive crystalline rocks.

The average precipitation is from 8 to 16 inches (203 to 406 mm) in the valleys and can range up to 73 inches (1854 mm) in the mountains. In the northern and western portions of the MLRA, peak precipitation occurs in the winter months. The southern and eastern portions have a greater incidence of high-intensity summer thunderstorms; hence, a significant amount of precipitation occurs during the summer months. The average annual temperature is 30 to 50 degrees Fahrenheit (-1 to 15 C). The freeze-free period averages 140 days and ranges from 60 to 220 days, generally decreasing in length with elevation.

The dominant soil orders in this MLRA are Aridisols, Entisols, Inceptisols, and Mollisols. The lower elevations are dominated by a frigid temperature regime, while the higher elevations experience cryic temperature regimes. Mesic temperature regimes come in on the lower elevations and south facing slopes in the southern portion of this MLRA. The soil moisture regime is typically xeric in the northern part of the MLRA, but grades to ustic in the extreme eastern and southern parts. The minerology is generally mixed and the soils are very shallow to very deep, generally well drained, and loamy or loamy-skeletal.

LRU notes

E47C is the Uinta Mountains portion of MLRA 47 that run east and west which includes the Uinta Wilderness and The Flaming Gorge National Recreation Area and towns such as Evanston, Wyoming, Hanna and Tabiona, Utah. Structurally these mountains consist of a broadly folded anticline that has an erosion resistance quartzite core. The Duchesne River and many other tributaries to the Green River run through this range, as well as the headwaters of the Bear River. The lower elevations are dominated by a frigid temperature regime, while the higher elevations experience cryic temperature regimes. The soil moisture regime is typically ustic. The minerology is generally mixed

and the soils are very shallow to very deep, generally well drained, and loamy, sandy or sandy-skeletal.

Ecological site concept

The soils of this site formed mostly in slope alluvium and colluvium derived from sandstone and quartzite or sandstone and shale. Surface soils are extremely cobbly loam to extremely bouldery loam in texture. Rock fragments may be present on the soil surface and throughout the profile and make up more than 50 percent of the soil volume. These soils are deep to very deep, well-drained, and have moderately permeability. pH is neutral to moderately alkaline. Available water-holding capacity ranges from 2.0 to 3.8 inches of water in the upper 60 inches of soil. The soil moisture regime is mostly ustic and the soil temperature regime is cryic. Precipitation ranges from 24 to 34 inches annually.

Associated sites

F047XC508UT	High Mountain Loam (quaking aspen) Sites often occur adjacent to each other.
-------------	----------------------------------------------------------------------------------------

Similar sites

F047XC512UT	High Mountain Stony Loam (Douglas-fir) Similar floristic characteristics, however this site is not as steep.
-------------	------------------------------------------------------------------------------------------------------------------------

Table 1. Dominant plant species

Tree	(1) <i>Pseudotsuga menziesii</i>
Shrub	(1) <i>Mahonia repens</i>
Herbaceous	(1) <i>Carex geyeri</i>

Physiographic features

This site can be found on mountain slopes on very steep slopes between 50 and 90 percent. It can occur at elevations between 6,000 to 9,800 feet. Flooding and ponding do not occur on this site.

Table 2. Representative physiographic features

Landforms	(1) Mountain slope (2) Mountain
Runoff class	High
Flooding frequency	None
Ponding frequency	None
Elevation	1,829–2,987 m
Slope	50–90%
Ponding depth	Not specified
Water table depth	Not specified
Aspect	Aspect is not a significant factor

Climatic features

The climate of this site characterized by cold, snowy winters and cool summers. The average annual precipitation ranges from 24 to 34 inches. March thru May and August, are typically the wettest months with June and July being the driest. The most reliable sources of moisture for plant growth are the snow that accumulates over the winter, and spring rains. Summer thunderstorms are intermittent and sporadic in nature, and thus, are less reliable sources of moisture to support vegetative growth on this site.

Table 3. Representative climatic features

Frost-free period (characteristic range)	
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	610-864 mm
Frost-free period (average)	70 days
Freeze-free period (average)	
Precipitation total (average)	

Influencing water features

Due to its landscape position, this site is not influenced by streams or wetlands.

Wetland description

N/A

Soil features

The soils on this site were formed in slope alluvium and colluvium derived from sandstone and quartzite and shale. The soil is well drained with moderate permeability in the upper 10 inches of soil. The soil is deep with greater than 60 inches to bedrock or a restrictive layer. The soil texture at the surface is extremely boulder loam or extremely cobbly loam. Surface gravels are 1 to 18 percent cover and surface rocks larger than 3 inches are 0 to 48 percent cover. Subsurface gravels are 18 to 26 percent volume and rocks over 3 inches in diameter are 31 to 44 percent volume. Available water capacity is between 2.0 and 3.8 in the upper 40 inches of soil. The soil pH is between 6.6 and 8.4. The soil temperature regime is frigid or cryic.

Soils associated with this site:

Uintah Area (UT047): Dokie (72, 68), Cryochrepts (197), Bigtom (187)

Table 4. Representative soil features

Parent material	(1) Slope alluvium–sandstone and shale (2) Colluvium–sandstone and shale (3) Slope alluvium–quartzite (4) Colluvium–quartzite
Surface texture	(1) Extremely bouldery loam (2) Extremely cobbly loam
Family particle size	(1) Loamy-skeletal
Drainage class	Well drained
Permeability class	Moderate
Depth to restrictive layer	152 cm
Soil depth	152 cm
Surface fragment cover ≤3"	1–18%
Surface fragment cover >3"	0–48%
Available water capacity (0-101.6cm)	5.08–9.65 cm
Calcium carbonate equivalent (0-101.6cm)	0–10%
Electrical conductivity (0-101.6cm)	0 mmhos/cm

Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	6.6–8.4
Subsurface fragment volume <=3" (Depth not specified)	18–26%
Subsurface fragment volume >3" (Depth not specified)	31–44%

Ecological dynamics

Ecological Dynamics of the Site

It is impossible to determine in any quantitative detail the Historic Climax Plant Community (HCPC) for this ecological site because of the lack of direct historical documentation preceding all human influence. In some areas, the earliest reports of dominant plants include the cadastral survey conducted by the General Land Office, which began in the late 19th century for this area. However, up to the 1870s the Shoshone Indians, prevalent in northern Utah and neighboring states, grazed horses and set fires to alter the vegetation for their needs. In the 1860s, Europeans brought cattle and horses to the area, grazing large numbers of them on unfenced parcels year-long). Itinerant and local sheep flocks followed, largely replacing cattle as the browse component increased.

Below is a State and Transition Model diagram that illustrates the “phases” (common plant communities), and “states” (aggregations of those plant communities) that can occur on the site. Differences between phases and states depend primarily upon observations of a range of disturbance histories in areas where this ESD is represented. These situations include grazing gradients to water sources, fence-line contrasts, patches with differing dates of fire, herbicide treatment, tillage, and kinds and times of timber harvest, etc. Reference State 1 illustrates the common plant communities that probably existed just prior to European settlement.

The major successional pathways within states, (“community pathways”) are indicated by arrows between phases. “Transitions” are indicated by arrows between states. The drivers of these changes are indicated in codes decipherable by referring to the legend at the bottom of the page and by reading the detailed narratives that follow the diagram. The transition between Reference State 1 and State 2 is considered irreversible because of the naturalization of exotic species of both flora and fauna, possible extinction of native species, and climate change. There may have also been accelerated soil erosion.

When available, monitoring data (of various types) were employed to validate more subjective inferences made in this diagram. See the complete files in the office of the State Range Conservationist for more details.

The plant communities shown in this State and Transition Model may not represent every possibility, but are probably the most prevalent and recurring plant communities. As more monitoring data are collected, some phases or states may be revised, removed, and/or new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities.” According to the USDA NRCS National Range & Pasture Handbook , Desired Plant Communities (DPC’s) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including descriptions of a plant community is to capture the current knowledge at the time of this revision.

Community 1.1

Reference State

The Reference State is a description of this ecological site just prior to Euro-American settlement but long after the arrival of Native Americans. The description of the Reference State was determined by NRCS Soil Survey Type Site Location information and familiarity with rangeland relict areas where they exist. At the time of European colonization, what would have been observed on these sites depended primarily on the time since the last wildfire occurred. If the site had not experienced fire for about 100 years, Douglas-fir (*Pseudotsuga menziesii*) would have been the dominant species occupying the site with a sparse understory (1.1) due to tree competition, overstory shading, and duff accumulation. Douglas-fir replaces itself without wildfire and would have been the climax dominant. Wildfire (1.1a) would have replaced these stands with a rich herb-dominated vegetation (1.2). In the absence of any major disturbance (1.2a, 1.3a, 1.4a, 1.5a), the re-sprouting vegetation including quaking aspen (*Populus tremuloides*), bigtooth maple (*Acer grandidentatum*), and/or Gambel oak (*Quercus gambelii*) would have

reclaimed the site (1.3), followed by the increasing presence of aspen (*Populus tremuloides*), first as saplings (1.4), and later as mature aspen with Douglas-fir seedlings (1.5), ultimately to where aspen would have been outcompeted by Douglas-fir returning to the climax vegetation (1.1). A more complete list of species by lifeform for the Reference State is available in accompanying tables in the "Plant Community Composition by Weight and Percentage" section of this document. Wildfire (1.1a, 1.5b) would have been the primary disturbance factor prior to colonization.

Community Phase 1.1: densely canopied Douglas-fir/ sparse understory

This plant community (1.1) would have been characterized by a stand of mature Douglas-fir with a sparse understory of Geyer's sedge (*Carex geyeri*), bluegrass (*Poa* spp.), chokecherry (*Prunus virginiana*), Fendler's meadow-rue (*Thalictrum fendleri*), and creeping barberry (*Mahonia repens*).

Community Pathway 1.1a:

Wildfire would have removed the trees, allowing shade-intolerant herbs to flourish briefly.

Community Phase 1.2: herb-dominated

This plant community would have developed within the first 5 years since the last fire, with Geyer's sedge, bluegrass, slender wheatgrass (*Elymus trachycaulus*), heartleaf arnica (*Arnica cordifolia*) and feathery false lily of the valley (*Maianthemum racemosum* ssp. *racemosum*) being the dominant understory species.

Community Pathway 1.2a:

After about 5 years, shrubs would begin to establish in the site.

Community Phase 1.3: re-sprouting aspen, maple, oak/ herbaceous understory

For a period of about 5 to 60 years post fire, a mixture of re-sprouting shrubs would have been dominant on the site, including, quaking aspen, bigtooth maple, and Gambel oak. Other prominent shrubs would have been mountain snowberry (*Symphoricarpos oreophilus*), chokecherry (*Prunus virginiana*), Saskatoon serviceberry (*Amelanchier alnifolia*), and Oregon boxleaf (*Paxistima myrsinites*), among others. The herbaceous understory would also have been fairly intact.

Community Pathway 1.3a:

About 60 years after fire, aspen would become established in the site.

Community Phase 1.4: immature aspen

This plant community would have been dominated by a stand of immature aspen, which is a seral species while Douglas-fir begins to establish itself under other nurse species. A stand of immature aspen would have existed approximately 60 to 80 years since last fire.

Community Pathway 1.4a:

Aspen would have continued to mature while Douglas-fir would have become well established in the understory.

Community Phase 1.5: mature aspen/ Douglas-fir

A stand of mature aspen intermixed with Douglas-fir saplings would have been encountered approximately 80 to 100 years post fire.

Community Pathway 1.5a:

After about 100 years following the last fire, Douglas-fir would become mature, shading out aspen and the shade-intolerant shrub and herb species in the understory.

Community Pathway 1.5b:

Wildfire would have removed the trees, allowing shade-intolerant herbs to flourish briefly.

Transition T1a: from State 1 to State 2 (Reference State to Secondary Forest/ Introduced State)

The simultaneous introduction of exotic species, both plants and animals, and possible extinctions of native flora and fauna, along with climate change, has caused State 1 to transition to State 2. Europeans further altered this vegetation largely through logging, livestock grazing, trapping of fur beaver and changing the fire regime. Continued impacts could prevent the recovery toward potential conifer dominance (State 2, various phases). The reversal of these changes (i.e. a return pathway) back to State 1 is not practical.

a. Nature of Forest Community

The overstory tree canopy cover is 30 to 40 percent. Common understory plants are Geyer sedge, wheeler bluegrass, heartleaf arnica, chokecherry, canyon maple, and creeping Oregon grape. Understory composition by air-dry weight is about 25 percent perennial grasses and grass-like plants, 10 percent forbs, and 65 percent shrubs. Understory production ranges from 500 pounds per acre in favorable years to about 100 pounds per acre in unfavorable years. Understory production includes the total annual production of all species within 4.5 feet of the ground surface.

State 2

Secondary Forest/ Introduced State

Community 2.2

Secondary Forest/ Introduced State

State 2 is similar to State 1 in form and function, with the exception of the presence of non-native plants and animals, possible extinctions of native species, a different climate, and a secondary stand of trees. State 2 is a description of the ecological site shortly following Euro-American settlement. This state can be regarded as the current potential. With the least amount of disturbance or manipulation of fire regime (Alexander 1985; 1988), a mature stand of Douglas-fir with a sparse understory component is expected at this site (2.1). As with the Reference State, time since last wildfire remains the key factor in determining what vegetation would be encountered here. Wildfire, particularly crown fires, or complete harvesting of the forest (2.1b, 2.5b) will replace these stands with a rich herb-dominated vegetation. (2.2). In the absence of any major disturbance (2.2a, 2.3a, 2.4a, 2.5a), the vegetation will progress into more of a shrub-herb co-dominance (2.3), followed by the increasing presence of aspen first as saplings (2.4), and later as mature aspen with Douglas-fir (2.5), ultimately to where Douglas-fir will outcompete aspen returning to the climax vegetation (2.1). Because soils on this site are rocky they are considered "self-armoring," thus making the site resistant to the effects of erosion. Heavy livestock grazing and tree harvests will reduce the stability of this site, while a reduction in livestock use and tree harvests will maintain stability.

Community Phase 2.1: densely canopied Douglas-fir/ sparse understory

This plant community (2.1) is characterized by a stand of mature Douglas-fir with a sparse understory of Geyer's sedge, and bluegrass may be present.

Community Pathway 2.1a:

A stand-replacing wildfire or intensive logging will set the vegetation back to an early seral herb-dominated phase. Logging opens up the forest canopy allowing more understory for 20 to 30 years.

Community Pathway 2.1b:

The removal of only the mature Douglas-fir will allow some aspen to return and the immature Douglas-fir to continue growing.

Community Phase 2.2: herb-dominated

This plant community will develop within the first 5 years since the last fire. Dominant grasses are Geyer's sedge, bluegrass, and slender wheatgrass with heartleaf arnica and feathery false lily of the valley as common forbs. A small component of introduced species may also be present.

Community Pathway 2.2a:

The combination of heavy season long livestock grazing and fire exclusion will accelerate woody plant (shrub) establishment and diminish the herbaceous understory.

Community Phase 2.3: re-sprouting aspen, maple, oak/ herbaceous understory

A plant community co-dominated by shrubs and herbs will develop approximately 5 to 60 years post fire. A small component of introduced species may be present.

Community Pathway 2.3a:

The combination of heavy season long livestock grazing and fire exclusion will accelerate woody plant establishment and diminish the herbaceous understory.

Community Pathway 2.3b:

Heavy browsing will lead to Phase 2.1, a densely canopied Douglas-fir community with a sparse understory.

Community Phase 2.4: immature aspen

Aspen will establish in the site 60 to 80 years after the last fire or complete tree removal.

Community Pathway 2.4a:

The combination of heavy season long livestock grazing and fire exclusion will accelerate woody plant establishment and diminish the understory.

Community Phase 2.5: mature aspen/ Douglas-fir

A stand of mature aspen intermixed with Douglas-fir will develop approximately 80 to 100 years following fire or complete tree removal.

Community Pathway 2.5a:

The combination of heavy season long livestock grazing and fire exclusion will accelerate woody plant establishment and diminish the understory.

Community Pathway 2.5b:

A stand-replacing wildfire or intensive logging will set the vegetation back to an early seral herb-dominated phase. Logging opens up the forest canopy allowing more understory for 20 to 30 years.

Community Pathway 2.5c:

The removal of mature aspen will leave a stand of immature aspen, possibly with a few Douglas-fir in the understory.

Transition T2a: from State 2 to State 3 (Secondary Forest/ Introduced State to Tertiary Forest/ Degraded State)

The Secondary Forest/ Introduced State will transition to the Tertiary Forest/ Degraded State following a second cycle of timber harvest or stand replacing wildfire, and further impacts from heavy continuous season-long grazing. Logging opens up the forest canopy allowing shade tolerant understory for 20 to 30 years. Secondary and tertiary disturbances produce an assemblage of vegetation from degraded temporary meadows to further simplified forests. A key indicator of the approach to this transition the presence of dominant trees that have reached the size required for commercial harvest. The trigger is a management decision. The second cycle of cutting typically is a clear cut, where all stems are downed, slash piled and burned.

State 3

Tertiary Forest/ Degraded State

Community 3.3

Tertiary Forest/ Degraded State

State 3 is characterized by tertiary forests that are further degraded in both understory and remnant tree layers. Active fire suppression speeds up the recovery of dominance by woody plants. Less predictable changes in the vegetation of State 3 will ensue due to global climate changes.

Community Phase 3.1: densely canopied Douglas-fir/ sparse understory

This plant community (3.1) is characterized by a stand of mature Douglas-fir. A sparse understory of Geyer's sedge and bluegrass may be present.

Community Pathway 3.1a:

A stand-replacing wildfire or intensive logging will set the vegetation back to an early seral herb-dominated phase. Logging opens up the forest canopy, allowing shade-intolerant grasses, forbs, and shrubs to flourish for 20 to 30 years.

Community Pathway 3.1b:

The removal of only the mature Douglas-fir will allow aspen to return and the immature Douglas-fir to continue growing.

Community Phase 3.2: herb-dominated

This plant community will develop within the first 5 years since the last fire. Dominant grasses are Geyer's sedge,

bluegrass, and slender wheatgrass. A small component of introduced species may be present.

Community Pathway 3.2a:

After about 5 years, shrubs will begin to establish in the site.

Community Phase 3.3: re-sprouting aspen, maple, oak/ herbaceous understory

A plant community co-dominated by shrubs and herbs will develop approximately 5 to 60 years after fire or complete tree removal. A small component of introduced species may be present.

Community Pathway 3.3a:

Aspen will become established at the site after 60-80 years following the last wildfire or complete tree removal.

Community Phase 3.4: immature aspen

Immature aspen dominate the stand 60 to 80 years following the last fire or complete tree removal.

Community Pathway 3.4a:

Aspen matures and immature Douglas-fir become well established in the understory 80 years after the last fire or complete tree removal.

Community Phase 3.5: mature aspen/ Douglas-fir

A stand of mature aspen intermixed with Douglas-fir will develop approximately 80 to 100 years after the last fire or complete tree removal.

Community Pathway 3.5a:

After about 100 years following the last fire, Douglas-fir will become mature, shading out aspen and the shade-intolerant shrub and herb species in the understory.

Community Pathway 3.5b:


A stand-replacing wildfire or intensive logging will set the vegetation back to an early seral herb-dominated phase clear cut. Logging opens up the forest canopy allowing more understory for 20 to 30 years.

Community Pathway 3.5c:

The removal of mature aspen will leave a stand of immature aspen, possibly with a few Douglas-fir in the understory.


State and transition model

3. Tertiary Forest/ Degraded State




T2a
(Log or WF; HCSLG)

2. Secondary Forest/ Introduced State



T1a
(HC;
Log or WF; HCSLG)

1. Reference State



HB Heavy Browsing
HC Historic Change
HCSLG Heavy Continuous Season Long Grazing
Log Logging

NS Natural Succession
STH Selective Timber Harvest
WF Wildfire

State 1
Reference State

Community 1.1
Reference State

a. Nature of Forest Community The overstory tree canopy cover is 30 to 40 percent. Common understory plants are Geyer sedge, wheeler bluegrass, heartleaf arnica, chokecherry, canyon maple and creeping Oregon grape. Understory composition by air-dry weight is about 25 percent perennial grasses and grass-like plants, 10 percent forbs, and 65 percent shrubs. Understory production ranges from 500 pounds per acre in favorable years to about 100 pounds per acre in unfavorable years. Understory production includes the total annual production of all species within 4.5 feet of the ground surface.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Shrub/Vine	73	219	364
Grass/Grasslike	28	84	140
Forb	11	28	56
Total	112	331	560

Additional community tables

Animal community

a. Livestock Grazing

This site is suited to cattle and sheep grazing during the summer and fall. Livestock will often concentrate on this site taking advantage of the shade and shelter offered by the tree overstory. Many areas are not used because of steep slopes or lack of adequate water. Attentive grazing management is required due to steep slopes and erosion hazards. Harvesting trees under a sound management program can open up the tree canopy to allow increased production of understory species desirable for grazing.

Wildlife species seeking food and cover in this forest site include moose, elk, mule deer, bear, porcupine, snowshoe hare, owl, and woodpecker.

Wood products

6. Silvicultural Practices

a. Douglas-fir seedling establishment may be improved by shade cards that will protect the trees from intense heat on southern or western aspects and by the presence of litter if it does not prevent the seed from reaching moist soil and does not absorb light rain showers.

b. Prescription burning program may be used to reduce competition before replanting a harvested site.

c. Douglas-fir will not regenerate in the shade. Seed tree harvests may be used to regenerate a site with or without prescription burning. Shelter wood cut should be avoided for Douglas-fir regeneration.

d. Harvest cut selectively or in small patches (size dependent upon site conditions) to enhance forage production.

1. Precommercial thinning and improvement cutting – removal of poorly formed, diseased, and low vigor trees of little or no value.

2. Commercial thinning – selectively harvest surplus trees to achieve desired spacing. Save large, healthy, full-crowned trees. Do not select only “high grade” trees during thinning.

e. Pest Control – use necessary and approved control for specific pests or diseases.

f. Fire hazard – fire is usually not a problem in mature grazed stands. Install firebreaks or firelines as necessary.

Other information

4. Limitations and Considerations

- a. Potential for sheet and rill erosion is moderate to severe depending on slope.
- b. Moderate to severe equipment limitations on steeper slopes and on sites having extreme surface stoniness.
- c. Proper spacing is the key to a well managed multiple use and multi-product forest.

5. Essential Requirements

- a. Adequately protect from uncontrolled burning.
- b. Protect soils from accelerated erosion.
- c. Apply proper grazing

Table 6. Representative site productivity

Common Name	Symbol	Site Index Low	Site Index High	CMAI Low	CMAI High	Age Of CMAI	Site Index Curve Code	Site Index Curve Basis	Citation
Rocky Mountain Douglas-fir	PSMEG	41	51	30	40	—	—	—	

Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel were also used.

Other references

Mauk, Ronald L., Henderson, Jan A. "Coniferous Forest Habitat Types of Northern Utah," General Technical Report INT 170, July 1984PSME/BERE/CAGE, Page 28

"Silvics of North America" Agriculture Handbook 654, Volume 1, Conifers

Contributors

Garth W. Leishman, David J. Somerville

Approval

Kendra Moseley, 2/05/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	

Date	05/13/2025
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-