

Ecological site R047XY378CO Mountain Stony Loam

Last updated: 2/11/2025
Accessed: 05/13/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 047X–Wasatch and Uinta Mountains

MLRA 47 occurs in Utah (86 percent), Wyoming (8 percent), Colorado (4 percent), and Idaho (2 percent). It encompasses approximately 23,825 square miles (61,740 square kilometers). The northern half of this area is in the Middle Rocky Mountains Province of the Rocky Mountain System. The southern half is in the High Plateaus of the Utah Section of the Colorado Plateaus Province of the Intermontane Plateaus. Parts of the western edge of this MLRA are in the Great Basin Section of the Basin and Range Province of the Intermontane Plateaus. The MLRA includes the Wasatch Mountains, which trend north and south, and the Uinta Mountains, which trend east and west. The steeply sloping, precipitous Wasatch Mountains have narrow crests and deep valleys. Active faulting and erosion are a dominant force in controlling the geomorphology of the area. The Uinta Mountains have a broad, gently arching, elongated shape. Structurally, they consist of a broadly folded anticline that has an erosion-resistant quartzite core. The Wasatch and Uinta Mountains have an elevation of 4,900 to about 13,500 feet (1,495 to 4,115 meters).

The mountains in this area are primarily fault blocks that have been tilted up. Alluvial fans at the base of the mountains are recharge zones for the basin fill aquifers. An ancient shoreline of historic Bonneville Lake is evident on the footslopes along the western edge of the area. Rocks exposed in the mountains are mostly Mesozoic and Paleozoic sediments, but Precambrian rocks are exposed in the Uinta Mountains. The Uinta Mountains are one of the few ranges in the United States that are oriented west to east. The southern Wasatch Mountains consist of Tertiary volcanic rocks occurring as extrusive lava and intrusive crystalline rocks.

The average precipitation is from 8 to 16 inches (203 to 406 mm) in the valleys and can range up to 73 inches (1854 mm) in the mountains. In the northern and western portions of the MLRA, peak precipitation occurs in the winter months. The southern and eastern portions have a greater incidence of high-intensity summer thunderstorms; hence, a significant amount of precipitation occurs during the summer months. The average annual temperature is 30 to 50 degrees Fahrenheit (-1 to 15 C). The freeze-free period averages 140 days and ranges from 60 to 220 days, generally decreasing in length with elevation.

The dominant soil orders in this MLRA are Aridisols, Entisols, Inceptisols, and Mollisols. The lower elevations are dominated by a frigid temperature regime, while the higher elevations experience cryic temperature regimes. Mesic temperature regimes come in on the lower elevations and south facing slopes in the southern portion of this MLRA. The soil moisture regime is typically xeric in the northern part of the MLRA, but grades to ustic in the extreme eastern and southern parts. The minerology is generally mixed and the soils are very shallow to very deep, generally well-drained, and loamy or loamy–skeletal.

LRU notes

E47C is the Uinta Mountains portion of MLRA 47 that run east and west which includes the Uinta Wilderness and The Flaming Gorge National Recreation Area and towns such as Evanston, Wyoming, Hanna and Tabiona, Utah. Structurally these mountains consist of a broadly folded anticline that has an erosion resistance quartzite core. The

Duchesne River and many other tributaries to the Green River run through this range, as well as the headwaters of the Bear River.

Ecological site concept

The soils of this site formed mostly in colluvium from sandstone. Surface soils are very bouldery loam in texture. Rock fragments may be present on the soil surface and throughout the profile and make up more than 35 percent of the soil volume. These soils are deep to very deep, well-drained, and have moderate to moderately rapid permeability. pH is neutral. Available water-holding capacity ranges from 2.4 to 3.4 inches of water in the upper 60 inches of soil. The soil moisture regime is mostly ustic and the soil temperature regime is frigid. Precipitation ranges from 16 to 22 inches annually.

Associated sites

F048AY924CO	Douglas Fir/Gambel Oak This site is a forested site with Douglas Fir and Gambel's oak as dominant plants. This site is found mostly commonly on mountain slopes, hillslopes, scarp slopes on cuestas, and mesas. Soils are moderately deep to very deep (20 to 60+ inches). Soil surface texture is loam, very stony loam, clay loam, stony sandy clay loam or very stony sandy loam. Subsurface textures can be loamy-skeletal or fine-loamy. It is usually ustic udic or typic ustic and frigid. It is a Douglas Fir – Gambel's Oak – Arizona Fescue plant community. The effective precipitation ranges from 20 to 25 inches.
-------------	--

Similar sites

F047XC458UT	Mountain Stony Loam (quaking aspen thicket) Dominant vegetation is Aspen. Surface soils are very cobbly fine sandy loams in texture and loamy-skeletal family particle size. The soils of this site formed mostly in slope alluvium over outwash derived from metamorphic and sedimentary rock on outwash fans.
R047XC456UT	Mountain Stony Loam (antelope bitterbrush) Dominant vegetation is bitterbrush. Surface soils are cobbly loam in texture and loamy-skeletal family particle size. The soils of this site formed in alluvium derived from sandstone and quartzite with some locations derived from dolomitic limestone.
R047XC460UT	Mountain Stony Loam (shrub) Dominant vegetation is Alderleaf Mountainmahogany and Utah serviceberry. Surface soils are extremely stony loam in texture and loamy-skeletal family particle size. The soils on this site were formed in slope alluvium and colluvium from sandstone and quartzite. The subsoils are extremely stony sandy clay loams.
R047XC461UT	Mountain Stony Loam (curl-leaf mountain mahogany) Dominant vegetation is Curlleaf Mountainmahogany. Surface soils are extremely bouldery loam in texture and loamy-skeletal family particle size. The soils on this site were formed in slope alluvium and colluvium derived from sandstone and quartzite.
R047XC462UT	Mountain Stony Loam (mountain big sagebrush) Dominant vegetation is mountain big sagebrush. Surface soils are very cobbly loam, extremely cobbly loam or stony loam in texture and loamy-skeletal family particle size. It formed in colluvium derived mainly from Duchesne River interbedded sandstone and shale, and Uinta mountain sandstone and quartzite.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Cercocarpus ledifolius</i>
Herbaceous	Not specified

Physiographic features

This site occurs on mountainsides with slopes from 25 to 50 percent.

Table 2. Representative physiographic features

Landforms	(1) Mountainside
Runoff class	High
Flooding frequency	None
Ponding frequency	None
Elevation	2,134–2,438 m
Slope	25–50%
Aspect	Aspect is not a significant factor

Climatic features

The climate of this site is characterized by cold snowy winters and cool dry summers. The average precipitation ranges from 16 to 22 inches annually with 55 to 60 percent coming during the plant dormant period (October to March). Much of this precipitation comes as snow that acts as a reservoir for water until the growing season begins. This winter moisture is the most dependable supply of water for plant growth. Lower precipitation and higher evapotranspiration rates during July, August, and September cause a reduction in plant growth for all species and dormancy in many of the grasses and forbs.

Table 3. Representative climatic features

Frost-free period (characteristic range)	
Freeze-free period (characteristic range)	65-85 days
Precipitation total (characteristic range)	406-559 mm

Influencing water features

Due to its landscape position, this site is not influenced by streams or wetlands.

Wetland description

N/A

Soil features

The soils of this site formed mostly in colluvium from sandstone. Surface soils are very bouldery loam in texture. Rock fragments may be present on the soil surface and throughout the profile and make up more than 35 percent of the soil volume. These soils are deep to very deep, well-drained, and have moderate to moderately rapid permeability. pH is neutral. Available water-holding capacity ranges from 2.4 to 3.4 inches of water in the upper 60 inches of soil. The soil moisture regime is mostly ustic and the soil temperature regime is frigid. Precipitation ranges from 16 to 22 inches annually.

Table 4. Representative soil features

Parent material	(1) Colluvium–sandstone
Surface texture	(1) Very bouldery loam
Family particle size	(1) Loamy-skeletal
Drainage class	Well drained
Permeability class	Moderate to moderately rapid
Depth to restrictive layer	152–203 cm
Soil depth	152–203 cm
Surface fragment cover <=3"	15–19%

Surface fragment cover >3"	25–32%
Available water capacity (50.8-101.6cm)	6.1–8.64 cm
Calcium carbonate equivalent (50.8-101.6cm)	0%
Electrical conductivity (50.8-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (50.8-101.6cm)	0
Soil reaction (1:1 water) (50.8-101.6cm)	6.6–7.3
Subsurface fragment volume <=3" (50.8-101.6cm)	15–19%
Subsurface fragment volume >3" (50.8-101.6cm)	32–34%

Ecological dynamics

It is impossible to determine in any quantitative detail the Historic Climax Plant Community (HCPC) for this ecological site because of the lack of direct historical documentation preceding all human influence. In some areas, the earliest reports of dominant plants include the cadastral survey conducted by the General Land Office, which began in the late 19th century for this area. However, up to the 1870s the Shoshone Indians, prevalent in northern Utah and neighboring states, grazed horses and set fires to alter the vegetation for their needs. In the 1860s, Europeans brought cattle and horses to the area, grazing large numbers of them on unfenced parcels year-long. Itinerant and local sheep flocks followed, largely replacing cattle as the browse component increased.

Below is a State and Transition Model diagram that illustrates the “phases” (common plant communities), and “states” (aggregations of those plant communities) that can occur on the site. Differences between phases and states depend primarily upon observations of a range of disturbance histories in areas where this ESD is represented. These situations include grazing gradients to water sources, fence-line contrasts, patches with differing dates of fire, herbicide treatment, tillage, etc. Reference State 1 illustrates the common plant communities that probably existed just prior to European settlement.

The major successional pathways within states, (“community pathways”) are indicated by arrows between phases. “Transitions” are indicated by arrows between states. The drivers of these changes are indicated in codes decipherable by referring to the legend at the bottom of the page and by reading the detailed narratives that follow the diagram. The transition between Reference State 1 and State 2 is considered irreversible because of the naturalization of exotic species of both flora and fauna, possible extinction of native species, and climate change. There may have also been accelerated soil erosion.

When available, monitoring data (of various types) were employed to validate more subjective inferences made in this diagram. See the complete files in the office of the State Range Conservationist for more details.

The plant communities shown in this State and Transition Model may not represent every possibility, but are probably the most prevalent and recurring plant communities. As more monitoring data are collected, some phases or states may be revised, removed, and/or new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities.” According to the USDA NRCS National Range & Pasture Handbook (USDA-NRCS 2003), Desired Plant Communities (DPC’s) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including descriptions of a plant community is to capture the current knowledge at the time of this revision.

Reference State

The Reference State is a description of this ecological site just prior to Euro-American settlement but long after the arrival of Native Americans. The description of the Reference State was determined by NRCS Soil Survey Type Site Location information and familiarity with rangeland relict areas where they exist. The least modified plant community would have been dominated by a low-statured curl-leaf mountain mahogany (*Cercocarpus ledifolius*),

and a mixture of herbaceous species. Other common shrubs may have included antelope bitterbrush (*Purshia tridentata*) and mountain big sagebrush (*Artemisia tridentata* ssp. *vaseyana*). Dominant grasses would have included bluebunch wheatgrass (*Pseudoroegneria spicata*), needle and thread (*Hesperostipa comata*), spike fescue (*Leucopoa kingii*), and muttongrass (*Poa fendleriana*). Forbs would have included spiny phlox (*Phlox hoodii*), western mountain aster (*Symphyotrichum spathulatum*), and arrowleaf balsamroot (*Balsamorhiza sagittata*), among others (1.1). These sites would have had wildfire return intervals of approximately 20 to 50 years.

Community 1.1: scattered low-statured curl-leaf mountain mahogany and associated shrubs/ rich native perennial herbaceous understory

This plant community would have been characterized by a scattering of low-statured curl-leaf mountain mahogany. Antelope bitterbrush and mountain big sagebrush would have also been present with an understory composed of a variety of forbs (spiny phlox, arrowleaf balsamroot, etc.) and grasses (bluebunch wheatgrass, needle-and-thread, etc.).

Transition T1a: from State 1 to State 2 (Reference State to Curl-leaf Mountain Mahogany Non-native State)

The simultaneous introduction of exotic species, both plants and animals, possible extinctions of native flora and fauna, and climate change has caused State 1 to transition to State 2. Reversal of such historic changes (i.e. a return pathway) back to State 1 is not practical.

State 2 Curl-leaf Mountain Mahogany/ Introduced Non-natives State

State 2 is a description of the ecological site just following Euro-American settlement, which has been influenced by the introduction of several non-native plants and animals, possible extinctions of native species, and a different climate. The plant community will be very similar to State 1 with the exception that some introduced species are likely to be present. This plant community is considered the current potential, and is dominated by curl-leaf mountain mahogany with a scattering of antelope bitterbrush and mountain big sagebrush. The native perennial herbaceous understory is still intact, but a small component of non-native species will also be present. The resiliency of this state is maintained by the root-sprouting nature of the dominant shrubs. Reductions in livestock use during the growth season will maintain the stability of the state. Alternatively, continued season-long heavy livestock grazing will reduced the stability of this state.

Community Phase 2.1: scattered low-statured curl-leaf mountain mahogany and associated shrubs/ rich native perennial herbaceous understory

This plant community is characterized by a scattering of low-statured curl-leaf mountain mahogany, antelope bitterbrush, and mountain big sagebrush. The native perennial understory is composed of a variety of forbs (spiny phlox, western mountain aster, arrowleaf balsamroot, etc.) and grasses (bluebunch wheatgrass, spike fescue, needle and thread, etc.).

Transition T2a: from State 2 to State 3 (Curl-leaf Mountain Mahogany/ Introduced Non-natives State to Tall Dense Curl-leaf Mountain Mahogany State)

Heavy continuous livestock grazing took place on these sites between the 1850s and 1950s. The effects of this heavy livestock grazing were further worsened with fire exclusion policies. A rapid increase of deer numbers in the 1950s and 1960s caused high-lining of the trees. Curl-leaf mountain mahogany began to grow taller and larger in dbh (diameter at breast height), while the herbaceous understory species were reduced greatly. Key indicators of the approach to this transition are a loss of perennial grass understory, an increase in the shrub component relative to grasses, and an increase in height and dbh of mahogany, as well as high-lining and hedging of trees. This transition is triggered by sustained heavy grazing (over time), first by livestock, and later by excessive numbers of deer.

State 3 Tall Dense Curl-leaf Mountain Mahogany State

In the absence of fire, and continued heavy impacts from livestock, deer, and elk grazing, the native grasses will markedly decrease while the shrubs, mainly curl-leaf mountain mahogany, will grow taller and more dense as reproduction becomes rarer. The resiliency of this state is maintained root-sprouting nature of the dominant shrubs. Reductions in livestock use during the growth season will maintain the stability of the state. Alternatively, continued season-long use by both domestic and wild ungulates will reduced the stability of this state.

Community 3.1: abundant curl-leaf mountain mahogany/ depauperate understory

This plant community is characterized by a dramatic increase in curl-leaf mountain mahogany and substantial

reduction in the perennial herbaceous component.

Transition T3a: from State 3 to State 4 (Tall Dense Curl-leaf Mountain Mahogany State to Tall Hedged Curl-leaf Mountain Mahogany State)

The region saw a marked increase in mule deer numbers during the 1940s to 1960s. As a preferred forage species for deer, curl-leaf mountain mahogany experienced heavy browsing pressure during that time. Heavy browsing combined with continued lack of fire caused these sites to transition into a near-monoculture of curl-leaf mountain mahogany. A key indicator of the approach to this transition is a loss of young, short mountain mahogany, and an increase in tall, large mountain mahogany trees. Excessive ungulate use will trigger this transition. A restoration pathway can be established by restoring fire to the ecosystem and reducing animal use on these sites.

Restoration Pathway R3a: from State 3 to State 2 (Tall Dense Curl-leaf Mountain Mahogany State to Curl-leaf Mountain Mahogany/ Introduced Non-natives State)

Brush management using prescribed fire, mechanical, or herbicide treatments, or wildfire, in combination with re-seeding efforts should be considered only on gentle to level slopes (these locations however are unusual). Curl-leaf mountain mahogany will not consistently re-sprout after these treatments, and invasive annuals and biennials are very prominent after disturbances. Re-seeding after wildfire must be done promptly or annuals and bi-annuals will takeover.

State 4


Tall Hedged Curl-leaf Mountain Mahogany State

Following a period of intense deer browsing, curl-leaf mountain mahogany will appear hedged. Without fire, mountain mahogany will grow taller and begin to form a monoculture. Introduced annuals and biennials are also quite common in the understory. In this state, the rockiness of the soil makes it self-armoring, thus reducing the effects of erosion. The steep, rocky locations of this site limit proactive management options.

Community 4.1: near monoculture curl-leaf mountain mahogany/ introduced annuals & biennials

This plant community is super-dominated by curl-leaf mountain mahogany with an understory composed mainly of introduced annuals and biennials. The site will remain in this state until wildfire occurs.

State and transition model



BMA	Brush Management – all (chemical, mechanical, fire)	NF	No Fire
HB	Heavy Browsing	RS	Re-seeding
HC	Historic change	WF	Wetfire
HCSLG	Heavy Continuous Season Long Grazing		

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel were also used.

Other references

USDA-NRCS. 2003. National Range and Pasture Handbook. in USDA, editor, USDA-Natural Resources Conservation Service-Grazing Lands Technology Institute. Keywords: [Western US, Federal guidelines, Range pasture management]

Western Regional Climate Center, Western U.S. Climate Historical Summaries. Available at: <http://www.wrcc.dri.edu/summary/Climsmut.html>. Accessed 15 June 2009.

Web Soil Survey, Official Soil Series Descriptions. Available at: <http://soils.usda.gov/technical/classification/osd/index.html>. Accessed 15 June 2009.

Contributors

Dean Stacy
Suzanne Mayne

Approval

Sarah Quistberg, 2/11/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/13/2025
Approved by	Sarah Quistberg
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not

bare ground):

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

-
16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
-

17. **Perennial plant reproductive capability:**
-