

Ecological site R058BY258WY Shallow Clayey (SwCy) 15-17" PZ

Last updated: 12/10/2024
 Accessed: 05/14/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 058B–Northern Rolling High Plains, Southern Part

MLRA 58B is located in northeastern Wyoming (95 percent) and extreme southeastern Montana (5 percent). It is comprised of sedimentary plains, scoria hills, and river valleys. The major rivers include the Powder, Tongue, Belle Fourche, Cheyenne, and North Platte. Tributaries include the Little Powder River, Little Missouri River, Clear Creek, Crazy Woman Creek, and others. This MLRA is traversed by Interstates 25 and 90, and U.S. Highways 14 and 16. The extent of MLRA 58B covers approximately 12.3 million acres. Major land uses include rangeland (approximately 93 percent), cropland, pasture, and hayland (approximately 2 percent), and forest, urban, and miscellaneous uses (approximately 5 percent). Cities include Buffalo, Casper, Sheridan, and Gillette, WY. Land ownership is mostly private. Federal lands include the Thunder Basin National Grassland (U.S. Forest Service) and lands administered by the Bureau of Land Management. Areas of interest in MLRA 58B in Wyoming include Fort Phil Kearny State Historic Site, Glendo State Park, and Lake DeSmet. The elevations in MLRA 58B increase gradually from north to south and range from approximately 2,900 to 5,900 feet. A few buttes are higher than 6,800 feet. The average annual precipitation in this area ranges from 10 to 17 inches per year. Precipitation occurs mostly during the growing season, often during rapidly developing thunderstorms. Mean annual air temperature is 46 degrees Fahrenheit. Summer temperatures may exceed 100 degrees Fahrenheit. Winter temperatures may drop to below zero. Snowfall averages 45 inches per year, but varies from 25 to over 70 inches in some locales.

Classification relationships

USDA Natural Resources Conservation Service (NRCS):

Land Resource Region—G Western Great Plains Range and Irrigation; Major Land Resource Area (MLRA)—58B Northern Rolling High Plains, Southern Part (USDA, 2006)

Relationship to Other Classifications:

USDA Forest Service (FS) Classification Hierarchy:

Province—331 Great Plains-Palouse Dry Steppe; Section—331G-Powder River Basin; Subsections—331Gb Montana Shale Plains, 331Ge Powder River Basin, 331Gf South Powder River Basin-Scoria Hills (Cleland et al, 1997)

Environmental Protection Agency (EPA) Classification Hierarchy:

Level III Ecoregion—43 Northwestern Great Plains; Level IV Ecoregion—43p Scoria Hills, 43q Mesic-Dissected Plains, 43w Powder River Basin (EPA, 2013)

<https://www.epa.gov/eco-research/ecoregions>

Ecological site concept

The Shallow Clayey 15 to 17 inch Precipitation Zone ecological site occurs on nearly level to steeply sloping hills and ridges, on sedimentary plains or uplands. Primary production is from cool-season midgrasses (bunch and rhizomatous), and secondary warm-season mid- and shortgrasses. There is also lesser component of forbs and shrubs, with occasional trees. Soils are shallow to a restrictive layer, usually of bedrock. Soil textures are clay loam, clay, silty clay, or sandy clay with greater than 35% clay in the profile.

Associated sites

R058BY222WY	Loamy (Ly) 15-17" PZ Loamy soils are lower on the landform, similar to Clayey sites, with more interbedded sedimentary parent material.
R058BY204WY	Clayey (Cy) 15-17" PZ Clayey sites are mod deep to very deep soils with higher species diversity and production. Clayey occurs lower in the landform or on lower sloping sites.
R058BY262WY	Shallow Loamy (SwLy) 15-17" PZ Shallow Loamy sites are similar in production with a slight variation in species due to lighter textured soils. They occur on similar positions on interbedded sedimentary rock.

Similar sites

R058BY158WY	Shallow Clayey (SwCy) 10-14" PZ Shallow Clayey 10-14" P.Z. has lower productivity.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata</i> ssp. <i>wyomingensis</i>
Herbaceous	(1) <i>Pseudoroegneria spicata</i> (2) <i>Nassella viridula</i>

Physiographic features

This site occurs on nearly level to steeply sloping hills and ridges, on sedimentary plains or uplands.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Ridge
Runoff class	Medium to high
Flooding frequency	None
Ponding frequency	None
Elevation	1,067–1,463 m
Slope	0–60%
Water table depth	203 cm
Aspect	Aspect is not a significant factor

Climatic features

The average annual precipitation ranges from 10 to 17 inches per year across MLRA 58B. There are two Precipitation Zones (PZs). The 10 to 14 inch precipitation zone is predominant across the MLRA, including portions of Sheridan, Johnson, and Natrona Counties; portions of Campbell and Converse Counties; and smaller portions of Weston and Niobrara Counties. The 15 to 17 inch precipitation zone occurs in northern and eastern portions of the MLRA, including portions of Sheridan, Campbell, and western Crook Counties. Wide fluctuations in precipitation may occur from year to year, and occasional periods of extended drought (longer than one year in duration) can be expected. Two-thirds of the annual precipitation occurs during the growing season from May through September. Mean Annual Air Temperature (MAAT) is 46 degrees Fahrenheit. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may also occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranching operations during late winter and spring. High-intensity afternoon thunderstorms may occur during the summer. Annual wind speeds average about 5 mph. Daytime winds are generally stronger than nighttime winds. Occasional strong storms may bring brief periods of high winds with gusts of more than 75 mph. The average length of the freeze-free period (28 degrees Fahrenheit) is 125 days and generally occurs from May 16 to September 19. The average frost-free period (32 degrees Fahrenheit) is 101 days and generally occurs from June 1 to September 9.

The growth of native cool-season plants begins in late April to early May with peak growth occurring in mid to late June. Native warm-season plants begin growth in late May to early June and continue into August. Regrowth of cool-season plants occurs in September in most years, depending upon moisture.

Note: The climate described here is based on historic climate station data and is averaged to provide an overview of the annual precipitation, temperatures, and growing season. Future climate is beyond the scope of this document. However, research to determine the effects of elevated CO₂ and heating on mixed-grass prairie ecosystems, and how it may relate to future plant communities, is ongoing.

For detailed information, or to find a specific climate station, visit the Western Regional Climate Center (WRCC) website: Western Regional Climate Center, Historical Data, Western U.S. Climate summaries, NOAA Coop Stations, Wyoming (Note: Montana climate stations are also listed under the Wyoming link).
<https://wrcc.dri.edu/summary/Climsmwy.html>

Wind speed averages can be found at the WRCC home page, under the Specialty Climate tab: <https://wrcc.dri.edu/>

The following tables represent area-wide climate data for the 15 to 17 inch precipitation zone:

Table 3. Representative climatic features

Frost-free period (characteristic range)	88-105 days
Freeze-free period (characteristic range)	122-130 days
Precipitation total (characteristic range)	381-406 mm

Frost-free period (actual range)	83-109 days
Freeze-free period (actual range)	119-130 days
Precipitation total (actual range)	381-432 mm
Frost-free period (average)	101 days
Freeze-free period (average)	125 days
Precipitation total (average)	381 mm

Climate stations used

- (1) DOUGLAS 1 SE [USC00482685], Douglas, WY
- (2) BIDDLE 8 SW [USC00240743], Biddle, MT
- (3) DILLINGER [USC00482580], Gillette, WY
- (4) GILLETTE 4SE [USC00483855], Gillette, WY
- (5) LEITER 9N [USC00485506], Clearmont, WY
- (6) SHERIDAN CO AP [USW00024029], Sheridan, WY

Influencing water features

This upland ecological site is not influenced by a water table or run in from adjacent sites. Due to the semi-arid climate in which it occurs, the water budget is normally contained within the soil pedon. Soil moisture is recharged by spring rains, but it rarely exceeds field capacity in the upper 20 inches before being depleted by evapotranspiration. During intense precipitation events, precipitation rates frequently exceed infiltration rates and the site delivers moisture to downslope sites through surface runoff. Moisture loss through evapotranspiration exceeds precipitation for a majority of the growing season. Soil moisture is the primary limiting factor for vegetative production on this ecological site.

Wetland description

N/A

Soil features

The soils on this site are well drained, shallow to bedrock and formed in residuum and slope alluvium weathered from shale. They typically have a very slow to slow permeability class. The available water capacity is typically very low to low. The surface layer of the soils in this site are typically clay loam, but may include silty clay, clay, or silty clay loam. The surface layer ranges from a depth of 1 to 6 inches thick. The subsoil is typically clay, but may include silty clay, clay loam, or silty clay loam. Soils in this site typically have carbonates at the surface; but some soils may be leached up to 6 inches or do not contain carbonates as they formed in residuum weathered from non-calcareous shale. These soils are susceptible to erosion by water and wind. The potential for water erosion accelerates with increasing slope. The soil moisture regime is typically aridic ustic. The soil temperature regime is mesic or frigid.

Major soil series correlated to this ecological site include: Lismas and Samsil.

The attributes listed below represent 0-40 inches in depth or to the first restrictive layer.

Table 4. Representative soil features

Parent material	(1) Residuum—shale (2) Slope alluvium—shale
Surface texture	(1) Clay loam (2) Silty clay loam (3) Silty clay (4) Clay

Family particle size	(1) Clayey
Drainage class	Well drained
Permeability class	Very slow to slow
Soil depth	25–51 cm
Surface fragment cover <=3"	0–5%
Available water capacity (0cm)	5.08–14.22 cm
Calcium carbonate equivalent (0cm)	0–10%
Electrical conductivity (0cm)	0–4 mmhos/cm
Sodium adsorption ratio (0cm)	0–5
Soil reaction (1:1 water) (0cm)	6.1–8.4
Subsurface fragment volume <=3" (Depth not specified)	0–15%


Ecological dynamics

The Reference state is the plant community in which interpretations are primarily based and is used as a reference in order to understand the original potential of the site. The Reference state evolved under the combined influences of climatic conditions, periodic fire activity, grazing by large herbivores, and impacts from small mammals and insects. Changes may occur to the Reference State due to management actions such as continuous season-long or year-long grazing, increased stocking rates, climatic conditions such as drought, and natural events such as multiple fires in close succession. The Reference state is characterized by cool-season rhizomatous midgrass , cool-season bunch midgrasses and warm-season shortgrass. Forbs and shrubs are also present. Trees such as Rocky Mountain juniper and ponderosa pine occur in minor amounts. The Reference state is not necessarily the management goal, as other vegetative states may be considered desired plant communities as long as critical resource concerns are met.

In addition to the Reference state, other plant communities can occur on this site and are usually the result of historic management practices. Grazing practices such as continuous season-long or year-long grazing, heavier stocking rates, or a combination of these factors on this ecological site results in bunchgrasses such as needle and thread decreasing in both frequency and production. Grasses such as blue grama, threadleaf sedge, and sixweeks fescue will increase. Forbs and shrubs such as hairy false goldenaster, tansyaster, broom snakeweed, and fringed sagewort, will also increase. If continued, the plant community will become sod-bound, and all midgrasses can eventually be removed from the plant community. Over the long-term, this continuous use in combination with high stock densities, will result in bare ground developing, and shrubs such as pricklypear, broom snakeweed; and annual forbs such as wooly plantain, field cottonrose, and pepperweed increasing or invading. Other invasives include field brome (also known as Japanese brome) and cheatgrass. There are various transitional stages which may occur on this ecological site. The information presented is representative of a dynamic set of plant communities that illustrate the complex interaction of several ecological processes.

State and transition model

Shallow Clayey 15-17" PZ



State 1 Reference

The Reference state is characterized by three distinct plant communities. The plant communities, and various successional stages between them, represent the natural range of variability within the Reference state.

Dominant plant species

- Wyoming big sagebrush (*Artemisia tridentata ssp. wyomingensis*), shrub
- bluebunch wheatgrass (*Pseudoroegneria spicata*), grass
- western wheatgrass (*Pascopyrum smithii*), grass
- blue grama (*Bouteloua gracilis*), grass

Community 1.1

Wyoming Big Sagebrush, Bluebunch Wheatgrass, and Western Wheatgrass

This is the interpretive plant community for this site. It is well adapted to the Northern Great Plains climate. This community developed with grazing by large herbivores and is suited to grazing by domestic livestock. Historically, fires likely were patchy and randomly distributed. This plant community can be found on areas where grazed plants receive adequate periods of recovery during the growing season. The potential vegetation is about 75 percent grasses and grass-like, 10 to 15 percent forbs, and 10 to 15 percent woody plants. The plant community is predominately cool-season midgrasses, with a smaller component of warm-season shortgrasses. The major grasses and grass-like include western wheatgrass, green needlegrass, and bluebunch wheatgrass. Secondary grasses include thickspike wheatgrass, little bluestem, prairie Junegrass, threadleaf sedge, Cusick's bluegrass, needle and thread, and blue- and hairy grama, squirreltail, and plains muhly. A variety of forbs include American vetch, white- and purple prairieclover, breadroot scurfpea, and prairie coneflower. Primary shrubs are big

sagebrush, winterfat, or birdfoot sagebrush (see the Species Composition List for additional information.) Plant diversity is high. In the Shallow Clayey 15 to 17 inch Precipitation Zone (PZ) ecological site, the total annual production (air-dry weight) is about 1,300 pounds per acre during an average year, but it can range from about 1,000 pounds per acre in unfavorable years to about 1,600 pounds per acre in above-average years. Defoliation levels should be determined as part of a grazing management plan based on objectives. Nutrient and water cycles, and energy flow are functioning properly. Infiltration rates are moderate, and soil erosion is low. Litter is properly distributed where vegetative cover is continuous. Decadence and natural plant mortality are low. This community is resistant to many disturbances except excessive grazing, or development into urban or other uses.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1177	1530	1883
Shrub/Vine	364	473	577
Forb	168	219	269
Tree	28	37	50
Total	1737	2259	2779

**Figure 9. Plant community growth curve (percent production by month).
WY1501, 15-17NP Upland sites.**

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	10	25	40	10	5	5	5	0	0

Community 1.2

Wyoming Big Sagebrush and Western Wheatgrass

This plant community developed with excessive grazing without adequate recovery opportunity during the growing season. Green needlegrass may initially increase or decrease depending on the season of grazing use. Palatable forbs such as white and purple prairieclover, American vetch, and penstemon are present in reduced amounts. white sagebrush (cudweed sagewort), hairy false goldenaster, slimflower scurfpea, large Indian breadroot, scarlet globemallow, broom snakeweed and fringed sagewort, have increased. Big sagebrush canopy has also increased. Natural disturbances such as lack of fire can contribute to this shift. The total annual production (air-dry weight) is about 1,200 pounds per acre during an average year, but it can range from about 900 pounds per acre in unfavorable years to about 1,500 pounds per acre in above-average years. Total aboveground biomass has been reduced. Reduction of rhizomatous wheatgrasses, nitrogen-fixing forbs, and increased warm-season shortgrasses have begun to alter the biotic integrity of this community. Water and nutrient cycles may be impaired. Nearly all plant species typically found in the community 1.1 are present and will respond to changes in grazing management.

**Figure 10. Plant community growth curve (percent production by month).
WY1501, 15-17NP Upland sites.**

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	10	25	40	10	5	5	5	0	0

Community 1.3

Decreased Midgrasses, Decreased Western Wheatgrass, Decreased Green Needlegrass and Bluebunch Wheatgrass, Increased Blue Grama and Threadleaf Sedge

This plant community developed with excessive grazing without adequate recovery during the growing season. Grazing-tolerant species such as blue grama and threadleaf sedge have noticeably increased. Midgrasses such as green needlegrass may initially increase or decrease depending on the season of grazing use. Palatable forbs such as white and purple prairieclover, American vetch, and penstemon are present in reduced amounts. Hairy false goldenaster, slimflower scurfpea, scarlet globemallow, fringed sagewort, and broom snakeweed have increased. Big sagebrush canopy may be reduced by fire. Natural disturbances such as fire and drought can contribute to this shift. The total annual production (air-dry weight) is about 900 pounds per acre during an average year, but it can

range from about 700 pounds per acre in unfavorable years to about 1,100 pounds per acre in above-average years. Total aboveground biomass has been reduced. Reduction of rhizomatous wheatgrasses, nitrogen-fixing forbs, and increased warm-season shortgrasses have begun to alter the biotic integrity of this community. Water and nutrient cycles may be impaired. Nearly all plant species typically found in the Reference Plant Community are present and will respond to changes in grazing management.

Figure 11. Plant community growth curve (percent production by month). WY5803, Northern Rolling High Plains, Southern Part, cool-season/warm-season co-dominant. Cool-season/warm-season co-dominant.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
		3	10	20	28	21	10	5	3		

Pathway 1.1A Community 1.1 to 1.2

Excessive grazing without adequate recovery between grazing events, drought, or lack of fire, can shift this plant community toward community 1.2. Over a period of years, plant species less tolerant to frequent and severe defoliation will begin to decrease, and those more tolerant will begin to increase. Excessive seasonal-use grazing, from year-to-year (i.e. spring grazing only) will result in a loss of cool-season species. Conversely, summer use only will result in a reduction of warm-season species. Big sagebrush canopy may also increase with lack of fire. Biotic integrity and water and nutrient cycles may become impaired because of this community pathway.

Pathway 1.1B Community 1.1 to 1.3

Excessive grazing without adequate recovery between grazing events, drought, or fire can shift this plant community toward the community 1.3. Over a period of years, plant species less tolerant to frequent and severe defoliation will begin to decrease, and those more tolerant will begin to increase. Excessive grazing from year-to-year will result in a reduction or loss of cool-season species. Biotic integrity and water and nutrient cycles may become impaired because of this community pathway.

Pathway 1.2A Community 1.2 to 1.1

Grazing that allows for adequate recovery between grazing events, along with proper stocking rates, will shift community 1.2 back toward community 1.1. Natural disturbances such as return to normal precipitation and/or fire will contribute to this shift.

Pathway 1.3A Community 1.3 to 1.1

Grazing that allows for adequate recovery between grazing events, and proper stocking rates, will shift community 1.3 back toward community 1.1. Natural disturbances such as return to normal precipitation patterns and/or no fire will contribute to this shift.

State 2 Sod Bound

This state is characterized by the Sod Bound state. An ecological threshold has been crossed and a significant amount of production and diversity has been lost when compared to the Reference state. Significant biotic and soil changes have negatively impacted energy flow and the nutrient and hydrologic cycles. This is a very stable state, resistant to change due to the high tolerance of blue grama and/or buffalograss to grazing, the development of a shallow root system (root pan), and subsequent changes in hydrology and nutrient cycling. The loss of other functional/structural groups such as cool-season bunch and rhizomatous grasses, forbs, and shrubs, reduces the biodiversity and productivity of this site.

Dominant plant species

- blue grama (*Bouteloua gracilis*), grass
- threadleaf sedge (*Carex filifolia*), grass

Community 2.1

Blue Grama and Threadleaf Sedge

This plant community develops under long-term frequent and severe defoliation. This typically occurs when the community has been excessively grazed with heavy stocking rates, throughout the growing season over a period of many years. The midgrasses and palatable forbs have been eliminated. The dominant species are blue grama and threadleaf sedge. These species have developed into a sod-bound condition occurring in localized colonies exhibiting a mosaic appearance. Perennial threeawn species such as purple threeawn have increased. Forbs such as scarlet globemallow, wild onion, death camas, and slimflower scurfpea remain. Forbs and shrubs that continue to increase are Cuman ragweed (western ragweed), hairy false goldenaster, fringed sagewort, and pricklypear. Plant diversity is low. Energy flow, water cycle and mineral cycle have been negatively affected. Litter levels are very low and unevenly distributed. In the 15 to 17" PZ, the total annual production (air-dry weight) is about 500 pounds per acre during an average year, but it can range from about 400 pounds per acre in unfavorable years to about 650 pounds per acre in above-average years. This plant community is extremely resistant to change. Many plant species are missing a seed source is not readily available.

Figure 12. Plant community growth curve (percent production by month).
WY5803, Northern Rolling High Plains, Southern Part, cool-season/warm-season co-dominant. Cool-season/warm-season co-dominant.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
		3	10	20	28	21	10	5	3		

State 3

Eroded

The Eroded state develops with long-term excessive grazing or frequent and severe defoliation, without adequate recovery between grazing events. An ecological threshold has been crossed. Erosion and loss of organic matter or carbon reserves are concerns.

Dominant plant species

- plains pricklypear (*Opuntia polyacantha*), shrub
- Fendler's threeawn (*Aristida purpurea* var. *fendleriana*), grass
- field brome (*Bromus arvensis*), grass

Community 3.1

Plains Pricklypear, Purple Threeawn, and Field Brome

This plant community occurs where the rangeland is grazed year-round, at high stock densities. Physical impact such as trampling, soil compaction, and trailing typically contribute to this transition. The plant composition is made of annuals with a few species of perennial forbs and grasses that are very tolerant to frequent and severe defoliation. Grasses include purple threeawn. Annuals such as sixweeks fescue, Russian thistle, and kochia have increased or invaded. The dominant forbs include hairy false goldenaster, curlycup gumweed, field cottonrose, and woolly plantain. Green sagewort, broom snakeweed, and pricklypear are increasing. Annual bromes such as field brome (also known as Japanese brome), and cheatgrass invade. In the 15 to 17" PZ, the total annual production (air-dry weight) is about 450 pounds per acre during an average year, but it can range from about 300 pounds per acre in unfavorable years to about 600 pounds per acre in above-average years. Soil erosion hazard has increased due to the increase of bare ground. Runoff typically is high and infiltration is low. All ecological functions are impaired. Desertification is advanced.

Figure 13. Plant community growth curve (percent production by month).
WY5804, Northern Rolling High Plains, Southern Part upland w/warm-season. 10-14" PZ, with warm-season dominant grasses and forbs.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			5	20	35	30	8	2			

Transition T1A

State 1 to 2

Excessive grazing without adequate recovery between grazing events or frequent and severe defoliation, will shift this plant community across an ecological threshold toward the Sod Bound state. Biotic integrity and hydrologic function will be impaired because of this transition.

Transition T2A

State 2 to 3

Long-term excessive grazing or frequent and severe defoliation without adequate recovery between grazing events or heavy, excessive grazing with overstocking, will cause a shift across an ecological threshold to the Eroded state. Non-native annual bromes begin to invade in this transition.

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Cool-Season Rhizomatous			280–448	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	280–448	10–25
	thickspike wheatgrass	ELLAL	<i>Elymus lanceolatus ssp. lanceolatus</i>	280–448	10–25
2	Cool-Season Bunch Midgrasses			504–807	
	green needlegrass	NAVI4	<i>Nassella viridula</i>	280–448	10–25
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	112–179	5–10
	Cusick's bluegrass	POCU3	<i>Poa cusickii</i>	112–179	1–10
3	Warm-Season Shortgrass			112–179	
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	112–179	5–10
4	Miscellaneous			280–448	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	280–448	5–25
	hairy grama	BOHI2	<i>Bouteloua hirsuta</i>	280–448	5–25
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	56–90	1–5
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	56–90	1–5
	needle and thread	HECO26	<i>Hesperostipa comata</i>	56–90	1–5
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	56–90	1–5
	needleleaf sedge	CADU6	<i>Carex duriuscula</i>	56–90	1–5
	threadleaf sedge	CAFI	<i>Carex filifolia</i>	56–90	1–5
	squirreltail	ELEL5	<i>Elymus elymoides</i>	56–90	1–5
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	56–90	1–5
	plains muhly	MUCU3	<i>Muhlenbergia cuspidata</i>	56–90	1–5
	Grass, perennial	2GP	<i>Grass, perennial</i>	56–90	1–5
	plains reedgrass	CAMO	<i>Calamagrostis montanensis</i>	56–90	1–5
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	56–90	1–5

Forb					
5	Forbs			168–269	
	American vetch	VIAM	<i>Vicia americana</i>	56–90	1–5
	aster	ASTER	<i>Aster</i>	56–90	1–5
	desertparsley	LOMAT	<i>Lomatium</i>	56–90	1–5
	large Indian breadroot	PEES	<i>Pedimelum esculentum</i>	56–90	1–5
	common yarrow	ACMI2	<i>Achillea millefolium</i>	56–90	1–5
	rosy pussytoes	ANRO2	<i>Antennaria rosea</i>	56–90	1–5
	milkvetch	ASTRA	<i>Astragalus</i>	56–90	1–5
	stemless mock goldenweed	STAC	<i>Stenotus acaulis</i>	56–90	1–5
	sulphur-flower buckwheat	ERUM	<i>Eriogonum umbellatum</i>	56–90	1–5
	scarlet beeblossom	GACO5	<i>Gaura coccinea</i>	56–90	1–5
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	56–90	1–5
	white prairie clover	DACA7	<i>Dalea candida</i>	56–90	1–5
	textile onion	ALTE	<i>Allium textile</i>	56–90	1–5
	spiny phlox	PHHO	<i>Phlox hoodii</i>	56–90	1–5
	prairie thermopsis	THRH	<i>Thermopsis rhombifolia</i>	56–90	1–5
	tapertip hawksbeard	CRAC2	<i>Crepis acuminata</i>	56–90	1–5
	Forb, perennial	2FP	<i>Forb, perennial</i>	56–90	1–5
Shrub/Vine					
6	Shrubs			364–577	
	big sagebrush	ARTR2	<i>Artemisia tridentata</i>	73–146	–
	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i>	73–146	–
	birdfoot sagebrush	ARPE6	<i>Artemisia pedatifida</i>	15–73	–
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	15–73	–
	Woods' rose	ROWO	<i>Rosa woodsii</i>	15–73	–
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	15–73	–
Tree					
7	Trees			28–50	
	Rocky Mountain juniper	JUSC2	<i>Juniperus scopulorum</i>	0–73	–
	ponderosa pine	PIPO	<i>Pinus ponderosa</i>	0–73	–

Animal community

Animal Community – Wildlife Interpretations (from 2001 ESD, will be revised in future updates)

Rhizomatous wheatgrass/ Green Needlegrass (Reference): The predominance of grasses in this plant community favors grazers and mixed-feeders, such as bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. However, topographical variations could provide some escape cover. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Many grassland-obligate small mammals would occur here.

Heavy Sagebrush: This plant community can provide important winter foraging for elk, mule deer and antelope, as

sagebrush can approach 15 percent protein and 40 to 60 percent digestibility during that time. This community provides excellent escape and thermal cover for large ungulates, as well as nesting and brood rearing habitat for sage grouse.

Mixed Sagebrush/Grass: The combination of an overstory of sagebrush and an understory of grasses and forbs provide a very diverse plant community for wildlife. The crowns of sagebrush tend to break up hard crusted snow on winter ranges, so mule deer and antelope may use this state for foraging and cover year-round, as would cottontail and jack rabbits. It provides important winter, nesting, brood-rearing, and foraging habitat for sage grouse. Brewer's sparrows' nest in big sagebrush plants and hosts of other nesting birds utilize stands in the 20-30% cover range.

Big sagebrush/Birdfoot sage/Cheatgrass: This plant community can provide important winter foraging for elk, mule deer and antelope, as sagebrush can approach 15 percent protein and 40 to 60 percent digestibility during that time. Grazing during spring to fall may be restricted due to low grass production. This community provides nesting and brood rearing habitat for sage grouse.

Animal Community – Grazing Interpretations Grazing Interpretations (updated in 2019 Provisional revision)

The following table is a guide to stocking rates for the plant communities described in the Shallow Clayey 15 to 17 inch Precipitation Zone ecological site. These are conservative estimates for initial planning. On-site conditions will vary, and stocking rates should be adjusted based on range inventories, animal kind/class, forage availability (adjusted for slope and distance to water), and the type of grazing system (number of pastures, planned moves, etc.), all of which is determined in the conservation planning process.

The following stocking rates are based on the total annual forage production in a normal year multiplied by 25 percent harvest efficiency of preferred and desirable forage species, divided by 912 pounds of ingested air-dry vegetation for an animal unit per month (Natl. Range and Pasture Handbook, 1997). An animal unit month is defined as the amount of forage required by one livestock animal, with or without one calf, for one month, and is shortened to AUM.

Plant Community (PC) Production (total lbs./acre in a normal year) and Stocking Rate (AUM/acre) are listed below:

Example:

1,300 lbs. per acre X 25% Harvest Efficiency = 325 lbs. forage demand for one month. 325 lbs. per acre/912 demand per AUM = 0.36

Reference Plant Community 1000-1600 .35

Heavy Sagebrush 800-1200 .3

Mixed Sagebrush/Grass 600-1200 .3

Big sagebrush/Birdfoot sage/Cheatgrass 300-600 .1

Increased Bare Ground PC - (*) (*)

* Highly variable stocking rates need to be determined on site. Grazing by domestic livestock is one of the major income- producing industries in the area. Rangelands in this area provide year-long forage under prescribed grazing for cattle, sheep, horses and other herbivores. During the dormant period, livestock may need supplementation based on reliable forage analysis.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group C and D. Infiltration ranges from very slow to slow. Runoff potential for this site varies from moderate to high depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75 percent ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where short-grasses form a strong sod and dominate the site. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information).

None expected on slopes less than 15%. On slopes greater than 15%, if rills are present, they will be discontinuous. Gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals and/or terracettes are not expected to occur on this site. Occasionally, bunch grasses may be pedestalled on steeper slopes (>15% slopes) with no exposed roots. Drought and fire should not increase the incidence of pedestals except on the steepest slopes. Fine litter will generally move short distances. Litter debris dams are occasionally present. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1 to 2 percent of the soil surface.

Recreational uses

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

Wood products

No appreciable wood products are present on the site.

Other products

None noted.

Other information

Site Development & Testing Plan

General Data (MLRA and Revision Notes, Hierarchical Classification, Ecological Site Concept, Physiographic, Climate, and Water Features, and Soils Data):

Updated. All "Required" items complete to Provisional level

Community Phase Data (Ecological Dynamics, STM, Transition & Recovery Pathways, Reference Plant Community, Species Composition List, Annual Production Table):

Updated. All "Required" items complete to Provisional level.

Annual Production Table is from the "Previously Approved" ESD (2001).

The Annual Production Table and Species Composition List will be reviewed for future updates at the Approved level.

Each Alternative State/Community

Complete to Provisional level

Supporting Information (Site Interpretations, Assoc. & Similar Sites, Inventory Data References, Agency/State Correlation, References)

Updated. All "Required" items complete to Provisional level.

Wildlife Interpretations: Narrative is from "Previously Approved" ESD (2001). Wildlife species will need to be updated at the next Approved level.

Livestock Interpretations: Plant community names and stocking rates updated.

Hydrology, Recreational Uses, Wood Products, and Other Products carried over from previously "Approved" ESD (2001).

Existing NRI Inventory Data References updated. More field data collection is needed to support this site concept.

Reference Sheet

Rangeland Health Reference Sheet carried over from previously "Approved" ESD (2005).
It will be updated at the next "Approved" level.

"Future work, as described in a project plan, to validate the information in this provisional ecological site description is needed. This will include field activities to collect low and medium intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document." (NI 430_306 ESI and ESD, April 2015)

Inventory data references

Inventory information has been derived from data collection on private and federal lands by the following methods:

- Double Sampling (Determining Vegetation Production and Stocking Rates, WY-ECS-1)
- Rangeland Health (Interpreting Indicators of Rangeland Health, Version 4, 2005)
- Soil Stability (Interpreting Indicators of Rangeland Health, Version 4, 2005)
- Line Point Intercept (Monitoring Manual for Grassland, Shrubland, and Savanna Ecosystems, Volume II, 2005)
- Soil Pedon Descriptions (Field Book for Describing and Sampling Soils, Version 3, 2012)
- SCS-RANGE-417 (Production & Composition Record for Native Grazing Lands)

National Resources Inventory (NRI)

Number of Records: 8

Sample Period: 2005-2016

Counties: Campbell, Crook, Johnson

Additional reconnaissance data collection includes ocular estimates and other inventory data; vegetative clipping data for NRCS program support; field observations from experienced rangeland personnel

Data collection for this ecological site was done in conjunction with the progressive soil surveys within MLRA 58B Northern Rolling High Plains (Southern Part)

Note: Revisions to soil surveys are on-going. For the most recent updates, visit the Web Soil Survey, the official site for soils information: <http://websoilsurvey.nrcs.usda.gov/app/WebSoilSurvey.aspx>

Other references

Anderson, R.C. 2006. Evolution and origin of the central grassland of North America: Climate, fire, and mammalian grazers. *Journal of the Torrey Botanical Society* 133:626–647.

Bragg, T.B. 1995. The physical environment of the Great Plains grasslands. In: A. Joern and K.H. Keeler (eds) *The changing prairie*. Oxford University Press, Oxford, UK, pages 49–81.

Branson, D.H. and G.A. Sword. 2010. An experimental analysis of grasshopper community responses to fire and livestock grazing in a northern mixed-grass prairie. *Environmental Entomology* 39:1441–1446.

Brinson, M.M. 1993. A hydrogeomorphic classification for wetlands. Technical Report WRP–DE–4. U.S. Army Corps of Engineers Waterways Experiment Station, Vicksburg, MS.

Cleland, D., P. Avers, W.H. McNab, M. Jensen, R. Bailey, T. King, and W. Russell. 1997. National hierarchical framework of ecological units. In: *Ecosystem Management: Applications for Sustainable Forest and Wildlife Resources*, Yale University Press.

Coupland, R.T. 1958. The effects of fluctuations in weather upon the grasslands of the Great Plains. *Botanical Review* 24:273–317.

Davis, S.K., R.J. Fisher, S.L. Skinner, T.L. Shaffer, and R.M. Brigham. 2013. Songbird abundance in native and planted grassland varies with type and amount of grassland in the surrounding landscape. *Journal of Wildlife*

- DeLuca, T.H. and P. Lesica. 1996. Long-term harmful effects of crested wheatgrass on Great Plains grassland ecosystems. *Journal of Soil and Water Conservation* 51:408–409.
- Derner, J.D. and R.H. Hart. 2007. Grazing-induced modifications to peak standing crop in northern mixed-grass prairie. *Rangeland Ecology and Management* 60:270–276.
- Derner, J.D. and A.J. Whitman. 2009. Plant interspaces resulting from contrasting grazing management in northern mixed-grass prairie: Implications for ecosystem function. *Rangeland Ecology and Management* 62:83–88.
- Derner, J.D., W.K. Lauenroth, P. Stapp, and D.J. Augustine. 2009. Livestock as ecosystem engineers for grassland bird habitat in the western Great Plains of North America. *Rangeland Ecology and Management* 62:111–118.
- Dillehay, T.D. 1974. Late Quaternary bison population changes on the southern Plains. *Plains Anthropologist* 19:180–196.
- Dormaar, J.F. and S. Smoliak. 1985. Recovery of vegetative cover and soil organic matter during revegetation of abandoned farmland in a semiarid climate. *Journal of Range Management* 38:487–491.
- Guyette, Richard P., M.C. Stambaugh, D.C. Dey, and R.M. Muzika. (2012). Predicting fire frequency with chemistry and climate. *Ecosystems*, 15: 322-335.
- Harmony, K.R. 2007. Grazing and burning Japanese brome (*Bromus japonicus*) on mixed grass rangelands. *Rangeland Ecology and Management* 60:479–486.
- Heitschmidt, R.K. and L.T. Vermeire. 2005. An ecological and economic risk avoidance drought management decision support system. In: J.A. Milne (ed.) *Pastoral systems in marginal environments*, 20th International Grasslands Congress, July, 2005. Page 178.
- Knopf, F.L. 1996. Prairie legacies—Birds. In: F.B. Samson and F.L. Knopf (eds.) *Prairie conservation: Preserving North America's most endangered ecosystem*. Island Press, Washington, DC. Pages 135–148.
- Knopf, F.L. and F.B. Samson. 1997. Conservation of grassland vertebrates. In: F.B. Samson and F.L. Knopf (eds.) *Ecology and conservation of Great Plains vertebrates: Ecological Studies* 125. Springer-Verlag, New York, NY. Pages 273–289.
- Lauenroth, W.K., O.E. Sala, D.P. Coffin, and T.B. Kirchner. 1994. The importance of soil water in recruitment of *Bouteloua gracilis* in the shortgrass steppe. *Ecological Applications* 4:741–749.
- Laycock, W.A. 1988. History of grassland plowing and grass planting on the Great Plains. In: J.E. Mitchell (ed.) *Impacts of the Conservation Reserve Program in the Great Plains—symposium proceedings*, September 16–18, 1987. U.S. Dept. of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station, General Technical Report RM-158.
- Malloch, D.W., K.A. Pirozynski, and P.H. Raven. 1980. Ecological and evolutionary significance of mycorrhizal symbioses in vascular plants (a review). *Proceedings of the National Academy of Sciences* 77:2113–2118.
- Ogle, S.M., W.A. Reiners, and K.G. Gerow. 2003. Impacts of exotic annual brome grasses (*Bromus* spp.) on ecosystem properties of the northern mixed grass prairie. *American Midland Naturalist* 149:46–58.
- Roath, L.R. 1988. Implications of land conversions and management for the future. In: J.E. Mitchell (ed.) *Impacts of the Conservation Reserve Program in the Great Plains—symposium proceedings*, September 16–18, 1987. U.S. Dept. of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station, General Technical Report RM-158.
- Smoliak, S. and J.F. Dormaar. 1985. Productivity of Russian wildrye and crested wheatgrass and their effect on prairie soils. *Journal of Range Management* 38:403–405.

- Smoliak, S., J.F. Dormaar, and A. Johnston. 1972. Long-term grazing effects on Stipa-Bouteloua prairie soils. *Journal of Range Management* 25:246–250.
- Soil Survey Division Staff. 2017. Soil survey manual. U.S. Dept. of Agriculture Handbook 18.
- Soil Survey Staff. Official Soil Series Descriptions. U.S. Dept. of Agriculture, Natural Resources Conservation Service. Available online. https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2_053587. Accessed 15 November, 2017.
- Soil Survey Staff. Soil Survey Geographic (SSURGO) database. U.S. Dept. of Agriculture, Natural Resources Conservation Service.
- Soil Survey Staff. 2014. Keys to Soil Taxonomy, 12th edition. U.S. Dept. of Agriculture, Natural Resources Conservation Service.
- Soil Survey Staff. 2018. Web Soil Survey. U.S. Dept. of Agriculture, Natural Resources Conservation Service. Available online. <https://websoilsurvey.nrcs.usda.gov/app/>. Accessed 15 February, 2018.
- Soller, D.R. 2001. Map showing the thickness and character of Quaternary sediments in the glaciated United States east of the Rocky Mountains. U.S. Geological Survey Miscellaneous Investigations Series I-1970-E, scale 1:3,500,000.
- Stewart, Omer C. 2002. *Forgotten Fires*. Univ. of Oklahoma Press, Norman, OK.
- United States Department of Agriculture, Natural Resources Conservation Service. Glossary of landform and geologic terms. National Soil Survey Handbook, Title 430-VI, Part 629.02c. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2_054242 (Accessed 16 January, 2018).
- United States Army Corps of Engineers. 1987. Corps of Engineers wetlands delineation manual. Wetlands Research Program Technical Report Y-87-1. (<http://www.lrh.usace.army.mil/Portals/38/docs/USACE%2087%20Wetland%20Delineation%20Manual.pdf>). Waterways Experiment Station, Vicksburg, MS.
- United States Environmental Protection Agency, National Health and Environmental Effects Research Laboratory. 2013. Level III ecoregions of the continental United States. Available online. <https://www.epa.gov/eco-research/ecoregions>. Accessed 30 January, 2019.
- United States Department of Agriculture, Natural Resources Conservation Service. 2010a. Field indicators of hydric soils in the United States, version 7.0.
- United States Department of Agriculture, Natural Resources Conservation Service. 2013a. Climate data. National Water and Climate Center. <http://www.wcc.nrcs.usda.gov/climate> (Accessed 13 October, 2017).
- United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. Agriculture Handbook 296.
- United States Department of Agriculture, Natural Resources Conservation Service. 2013b. National Soil Information System. Available online. https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/geo/?Cid=nrcs142p2_053552. Accessed 30 October, 2017.
- United States Department of the Interior, Geological Survey. 2008. LANDFIRE 1.1.0 Vegetation Dynamics Models. <http://landfire.cr.usgs.gov/viewer/>.
- United States Department of the Interior, Geological Survey. 2011. LANDFIRE 1.1.0 Existing Vegetation Types. <http://landfire.cr.usgs.gov/viewer/>.

- Willeke, G.E. 1994. The national drought atlas [CD ROM]. U.S. Army Corps of Engineers, Water Resources Support Center, Institute for Water Resources Report 94-NDS-4.
- Wilson, S.D. and J.M. Shay. 1990. Competition, fire, and nutrients in a mixed-grass prairie. *Ecology* 71:1959–1967.
- With, K.A. 2010. McCown's longspur (*Rhynchophanes mccownii*). In: A. Poole (ed.) *The birds of North America* [online]. Cornell Lab of Ornithology, Ithaca, NY. <https://birdsna.org/Species-Account/bna/home>.
- Augustine, D.J., J. Derner, D. Milchunas, D. Blumenthal, and L. Porensky. 2017. Grazing moderates increases in C3 grass abundance over seven decades across a soil texture gradient in shortgrass steppe. In: *Journal of Vegetation Science*, DOI:10.1111/jvs.12508.
- Clark, J., E. Grimm, J. Donovan, S. Fritz, D. Engrstrom, and J. Almendinger. 2002. Drought cycles and landscape responses to past aridity on prairies of the Northern Great Plains, USA. *Ecology*, 83(3), Pages 595-601.
- Connell, L. C., J. D. Scasta, and L. M. Porensky. 2018. Prairie dogs and wildfires shape vegetation structure in a sagebrush grassland more than does rest from ungulate grazing. *Ecosphere* 9(8):e02390. 10.1002/ecs2.2390.
- Collins, S. and S. Barber. (1985). Effects of disturbance on diversity in mixed-grass prairie. In: *Vegetatio*, 64, pages 87-94.
- Egan, Timothy. 2006. *The Worst Hard Time*. Houghton Mifflin Harcourt Publishing Company, New York, NY.
- Guyette, R.P., M.C. Stambaugh, D.C. Dey, and R.M. Muzika. 2012. Predicting fire frequency with chemistry and climate. In: *Ecosystems*, 15: pages 322-335.
- Hart, R. and J. Hart. 1997. Rangelands of the Great Plains before European settlement. In: *Rangelands*, 19(1), pages 4-11.
- Hart, R. 2001. Plant biodiversity on shortgrass steppe after 55 years of zero, light, moderate, or heavy cattle grazing. In: *Plant Ecology*, 155, pages 111-118.
- Pellant, M., P. Shaver, D.A. Pyke, and J.E. Herrick. 2005. *Interpreting indicators of rangeland health, Version 4*. United States Department of the Interior, Bureau of Land Management.
- Porensky, L.M. and D.M. Blumenthal. 2016. Historical wildfires do not promote cheatgrass invasion in a western Great Plains steppe. In: *Biological Invasions* 18:3333-3349: DOI 10.1007/s10530-16-1225-z
- Porensky, L., J.D. Derner, and D.W. Pellatz. 2018. Plant community responses to historical wildfire in a shrubland-grassland ecotone reveal hybrid disturbance response. In: *Ecosphere*. DOI: 9(8):e02363. 10.1002/ecs2.2363.
- Mack, Richard N., and J.N. Thompson. 1982. Evolution in steppe with few large, hooved mammals. In: *The American Naturalist*. 119, No. 6, pages 757-773.
- Reyes-Fox, M., H. Stelzer, M.J. Trlica, G.S. McMaster, A.A. Andales, D.R. LeCain, and J.A. Morgan. 2014. Elevated CO2 further lengthens growing season under warming conditions. In: *Nature*, April 23, 2014. Available online. <http://www.nature.com/nature/journal/v510/n7504/full/nature13207.html>. Accessed 1 March, 2017.
- Schoeneberger, P.J., D.A. Wysockie, E.C. Benham, and Soil Survey Staff. 2012. *Field book for describing and sampling soils, Version 3.0*. U.S. Dept. of Agriculture, Natural Resources Conservation Service.
- Stahl, David W., E.R. Cook, M.K. Cleaveland, M.D. Therrell, D.M. Meko, H.D. Grissino-Mayer, E. Watson, and B.H. Luckman. Tree-ring data document 16th century megadrought over North America. 2000. In: *Eos*, 81(12), pages 121-125.
- Stewart, Omer C. 2002. *Forgotten Fires*. University of Oklahoma Press, Norman, OK.
- Stubbendieck, James, S.L. Hatch, and L.M. Landholt. 2003. *North American wildland plants*. Univ. of Nebraska

Press, Lincoln and London.

Zelikova, T.J., D.M. Blumenthal, D.G. Williams, L. Souza, D.R. LeCain, and J. Morgan. 2014. Long-term exposure to elevated CO₂ enhances plant community stability by suppressing dominant plant species in a mixed-grass prairie. In: Ecology, 2014 <https://www.pnas.org/content/111/43/15456>.

United States Department of Agriculture, Natural Resources Conservation Service. National Ecological Site Handbook, Title 190, Part 630, 1st Edition. Available online. <https://directives.sc.egov.usda.gov/>. Accessed 15 September, 2017.

United States Department of Agriculture, Natural Resources Conservation Service. 2009. Part 630, Hydrology, National Engineering Handbook

United States Department of Agriculture, Natural Resources Conservation Service. 1972-2012. National Engineering Handbook Hydrology Chapters. Available online. <http://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/water/?&cid=stelprdb1043063>. Accessed August, 2015.

United States Department of Agriculture, Natural Resources Conservation Service. 1997, revised 2003. National Range and Pasture Handbook. Available online. <http://www.glti.nrcs.usda.gov/technical/publications/nrph.html>. Accessed 26 February, 2018.

United States Department of Agriculture, Natural Resources Conservation Service. National Soil Survey Handbook title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. Web Soil Survey. Available online. <http://websoilsurvey.sc.egov.usda.gov/App/WebSoilSurvey.aspx>. Accessed 15 November, 2017.

Contributors

Everett Bainter
Glenn Mitchell

Approval

Kirt Walstad, 12/10/2024

Acknowledgments

Project Staff:

Kimberly Diller, Ecological Site Inventory Specialist, NRCS MLRA SSO, Pueblo CO

Mike Leno, Project Leader, NRCS MLRA SSO, Buffalo, WY

Partners/Contributors:

Joe Dyer, Soil Scientist, NRCS MLRA SSO, Buffalo, WY

Arnie Irwin, Soil Scientist, BLM, Buffalo, WY

Blaine Horn, Rangeland Extension Educator, UW Extension, Buffalo, WY

Isabelle Giuliani, Resource Soil Scientist, NRCS, Douglas, WY

Mary Jo Kimble, Project Leader, NRCS MLRA SSO, Miles City, MT

Ryan Murray, Rangeland Management Specialist, NRCS, Buffalo, WY

Lauren Porensky, Ph.D., Ecologist, ARS, Fort Collins, CO

Chadley Prosser, Rangeland Program Manager, USFS, Bismarck, ND

Bryan Christensen, Ecological Site Inventory Specialist, NRCS-MLRA SSO, Pinedale, WY

Marji Patz, Ecological Site Inventory Specialist, NRCS-MLRA SSO, Powell, WY

Rick Peterson, Ecological Site Inventory Specialist, NRCS-MLRA SSO, Rapid City, SD

Program Support:

John Hartung, WY State Rangeland Management Specialist-QC, NRCS, Casper, WY

David Kraft, NRCS MLRA Ecological Site Inventory Specialist-QA, Emporia, KS

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/01/2005
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:** Due to the wide slope range associated with this site, the number and extent of rills will vary from none on sites with slopes of < 9% to common on slopes > 25%.

- 2. Presence of water flow patterns:** Due to the wide slope range associated with this site, water flow patterns will vary from barely observable on sites with slopes of < 9% from broken and irregular in appearance to continuous on slopes > 25%.

- 3. Number and height of erosional pedestals or terracettes:** Not evident on slopes <9%. Erosional pedestals will be present with terracettes present at debris dams on slopes >9%.

- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground is 20 to 35%.

- 5. Number of gullies and erosion associated with gullies:** Active gullies restricted to concentrated water flow patterns.

- 6. Extent of wind scoured, blowouts and/or depositional areas:** None.

- 7. Amount of litter movement (describe size and distance expected to travel):** Little to no plant litter movement occurs on slopes < 9%. Litter movement does occur on slopes > 25%.

- 8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of**

values): Plant cover and litter is at 65% or greater of soil surface and maintains soil surface integrity. Stability class anticipated to be 5 or greater.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Use soil series description for depth and color of A-horizon.
-

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Plant canopy, very slow to slow infiltration rates, the amount of bare ground, and steepness of slopes results in a naturally high runoff rate on slopes > 25%, even in HCPC.
-

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer would be expected except for the naturally occurring rooting restriction (bedrock or decomposing shale) at 10 to 20 inches.
-

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional: Mid-stature bunch grasses > mid-stature rhizomatous grasses short stature grasses/grasslikes shrubs forbs

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Very low.
-

14. **Average percent litter cover (%) and depth (in):** Litter cover is in contact with soil surface.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 1300 lbs./acre
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Blue grama, buffalograss, greasewood, rabbitbrush, broom snakeweed, Plains prickly pear and species found on noxious weed list.
-

17. **Perennial plant reproductive capability:** No limitations.
