

Ecological site R058CY090ND Saline Lowland

Last updated: 10/31/2018 Accessed: 05/11/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Jeff Printz
Contact for lead author	mark.hayek@nd.usda.gov
Date	04/10/2012
Approved by	Jeff Printz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills: None.
- 2. Presence of water flow patterns: None.
- 3. Number and height of erosional pedestals or terracettes: None.
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): Bare ground is < 5%.
- 5. Number of gullies and erosion associated with gullies: None.
- 6. Extent of wind scoured, blowouts and/or depositional areas: None.
- 7. Amount of litter movement (describe size and distance expected to travel): None.

- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values): Stability class 3 or greater.
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Use soil series description for depth, color, and structure of A horizon/surface layer.
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Combination of shallow and deep rooted species (short-, mid-, and tall rhizomatous and tufted perennial cool- and warm-season grasses) with fine and coarse roots positively influences infiltration.
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None.
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Mid, cool-season rhizomatous grass

Sub-dominant: Short, warm-season bunchgrasses

Other: Mid, cool-season bunchgrasses forbs shrubs

Additional: Due to differing root structure and distribution, Kentucky bluegrass and smooth brome do not fit into Reference Plant Community F/S groups.

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): None.
- 14. Average percent litter cover (%) and depth (in): Plant litter is in contact with soil surface.
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction): Representative value (RV) is 2200 lbs./acre air dry.
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: State and local noxious, Kentucky bluegrass, smooth bromegrass, Russian olive

17. **Perennial plant reproductive capability:** All species exhibit high vigor relative to climatic conditions. Do not rate based solely on seed production. Perennial grasses should have vigorous rhizomes or tillers.