

Ecological site R086BY005TX Blackland

Last updated: 9/21/2023
Accessed: 05/10/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 086B–Texas Blackland Prairie, Southern Part

MLRA 86B, the Southern Part of the Texas Blackland Prairie is located in east-central Texas. It makes up about 2,925 square miles (7,585 square kilometers). The towns of Brenham, Caldwell, La Grange, Schulenberg, Hallettsville, and Navasota are in the eastern part while Lincoln, Benchley, and Normangee are in the western part. The area supports tall and mid-grass prairies, but improved pasture, croplands, and urban development account for the majority of the acreage.

Classification relationships

USDA-Natural Resources Conservation Service, 2006.
-Major Land Resource Area (MLRA) 86B

Ecological site concept

The Blackland ecological site is distinguished by its deep productive soils that are richly black in color, and characterized by their shrink-swell nature. The sites are widely distributed across the region.

Associated sites

R086BY001TX	Chalky Ridge The Chalky Ridge site is often upslope from the Blackland. It differs from the site by having shallow soils and low soil fertility.
R086BY002TX	Claypan Prairie The Claypan Prairie site is often adjacent to the Blackland. It differs from the site by having a sandy loam surface soil layer over clay subsoils.
R086BY006TX	Loamy Bottomland The Blackland Site occurs at higher elevation in the drainage and provides runoff to the Loamy Bottomland Site.
R086BY007TX	Clayey Bottomland The Blackland Site occurs at higher elevation in the drainage and provides runoff to the Clayey Bottomland Site.

Similar sites

R086BY004TX	Eroded Blackland The Eroded Blackland site is similar to the Blackland site by having similar soil types and topography. It differs by having extensive erosion indicated by a partial or lost A horizon, active rills and/or gullies, and lower production.
-------------	--

Table 1. Dominant plant species

Tree	(1) <i>Celtis laevigata</i> var. <i>reticulata</i> (2) <i>Ulmus crassifolia</i>
Shrub	(1) <i>Symphoricarpos orbiculatus</i>
Herbaceous	(1) <i>Schizachyrium scoparium</i> (2) <i>Andropogon gerardii</i>

Physiographic features

This site includes nearly level, moderately sloping, gently rolling, and hilly topography. Slopes range from 0 to 8 percent, but are typically 1 to 5 percent.

Table 2. Representative physiographic features

Landforms	(1) Plains > Terrace (2) Plains > Ridge
Runoff class	High to very high
Flooding frequency	None
Ponding frequency	None
Elevation	200–500 ft
Slope	1–5%
Aspect	Aspect is not a significant factor

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	Not specified
Slope	0–8%

Climatic features

The climate for MLRA 86B is humid subtropical and is characterized by hot summers, especially in July and August, and relatively mild winters. Tropical maritime air controls the climate during spring, summer and fall. In winter and early spring, frequent surges of Polar Canadian air cause sudden drops in temperatures and add considerable variety to the daily weather. When these cold air masses stagnate and are overrun by moist air from the south, several days of cold, cloudy, and rainy weather follow. Generally, these occasional cold spells are of short duration with rapid clearing following cold frontal passages. The summer months have little variation in day-to-day weather except for occasional thunderstorms that dissipate the afternoon heat. The moderate temperatures in spring and fall are characterized by long periods of sunny skies, mild days, and cool nights. Rainfall during the spring and summer months generally falls during thunderstorms, and fairly large amounts of rain may fall in a short time. High intensity rains of short duration are likely to produce rapid runoff almost anytime during the year. The amount of rain that falls varies considerably from month-to-month and from year-to-year.

Table 4. Representative climatic features

Frost-free period (average)	254 days
Freeze-free period (average)	280 days
Precipitation total (average)	42 in

Climate stations used

- (1) LEXINGTON [USC00415193], Lexington, TX
- (2) BRENHAM [USC00411048], Brenham, TX
- (3) FLATONIA 4SE [USC00413183], Moulton, TX
- (4) WASHINGTON SP [USC00419491], Navasota, TX
- (5) HALLETTSVILLE 2 N [USC00413873], Hallettsville, TX

Influencing water features

The plant community of this site is not influenced by a stream.

Wetland description

Wetlands are not associated with this site.

Soil features

These soils are deep, dark colored to olive, slightly acid to calcareous clays. The soils have high shrink-swell properties often forming microdepressions and microknolls. When they are dry and cracked, they take in water rapidly. When they are wet and the cracks have closed, however, the water intake is slow. The soils hold large amounts of water and have high natural fertility. They seal over when cover is inadequate and erosion is then a hazard. The soils are sticky and plastic when wet. When moist, they are crumbly. Good soil structure is generally very evident under grass cover. In this condition, they favor air and water movement and root growth.

Soils correlated to this site include: Bleiblerville, Branyon, Denhawken, Denvaca, Dimebox, Elmendorf, Elmenwood, Frelsburg, Greenvine, Heiden, Houston Black, Leson, and Luling.

Table 5. Representative soil features

Parent material	(1) Residuum—shale
Surface texture	(1) Clay
Family particle size	(1) Clayey
Drainage class	Moderately well drained to well drained
Soil depth	80 in


Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	6–10 in
Calcium carbonate equivalent (0-40in)	0–40%
Electrical conductivity (0-40in)	0–4 mmhos/cm
Sodium adsorption ratio (0-40in)	0–2
Soil reaction (1:1 water) (0-40in)	5.6–8.4
Subsurface fragment volume <=3" (Depth not specified)	0–3%
Subsurface fragment volume >3" (Depth not specified)	0–1%

Ecological dynamics

This tallgrass prairie site evolved and was maintained by the grazing and herding effects of native large ungulates, by rodents and rabbits, by insects and disease, and the occurrence of periodic fire. Extreme climatic fluctuations over time may also have been important in the maintenance of the historic plant community. Continuous overgrazing by confined domestic livestock and the suppression of fire degrades the plant community. Continuous grazing will remove big bluestem (*Andropogon gerardii*), Indiangrass (*Sorghastrum nutans*), little bluestem (*Schizachyrium scoparium*), switchgrass (*Panicum virgatum*), and preferred forbs such as Engelmann's daisy (*Engelmannia peristeria*), Maximilian sunflower (*Helianthus maximilliani*), Illinois bundleflower (*Desmanthus illinoensis*), and compass plant (*Silphium* spp.). These plants will be replaced by less productive perennial and annual grasses and forbs including silver bluestem (*Bothriochloa saccharoides*), Texas wintergrass (*Nassella leucotricha*), windmillgrass (*Chloris* spp.), white tridens (*Tridens albescens*), annual broomweed (*Amphiachyris dracunculoides*), and snow on the prairie (*Euphorbia bicolor*). With continued continuous grazing and the absence of periodic fire, a community dominated by woody species including mesquite (*Prosopis glandulosa*), cedar elm (*Ulmus crassifolia*), netleaf hackberry (*Celtis laevigata*), and eastern red cedar (*Juniperus virginiana*) will replace the grassland.

State and transition model

Ecosystem states




T1A - No fire, no brush management, improper grazing management, drought

T1B - Brush management, crop cultivation, pasture planting, nutrient management, pest management


- R2A** - Fire, brush management, proper grazing, range planting
- T2A** - Brush management, crop cultivation, pasture planting, nutrient management, pest management
- R3A** - Fire, brush management, proper grazing, range planting
- T3A** - No fire, no brush management, heavy continuous grazing, no pest management

State 1 submodel, plant communities




- 1.1A** - No fire, no brush management, improper grazing management, drought
- 1.2A** - Fire, brush management, proper grazing

State 2 submodel, plant communities



- 2.1A** - No fire, no brush management, improper grazing management, drought

State 3 submodel, plant communities



**State 1
Prairie/Savannah**

Two communities exist in the Prairie/Savannah State: the 1.1 Tallgrass Prairie Community and the 1.2 Mesquite/Midgrass Savannah Community. Community 1.1 is characterized by tallgrasses dominating the understory, with woody cover less than 10 percent. Community 1.2 is characterized by a decrease in tallgrasses and an increase in midgrasses and forbs. Woody cover increases to 20 to 40 percent.

**Community 1.1
Tallgrass Prairie**



The characteristic plant community of this site is the reference plant community. This site is a true prairie with few large live oak (*Quercus virginiana*), elm, and hackberry trees along the draws and in occasional motts. Little bluestem dominates the plant community, making up 40 to 50 percent of the total annual yield. Indiangrass and big bluestem make up another 30 percent. Numerous other grasses occur in the potential plant community but less frequently and in smaller amounts. Texas wintergrass and Virginia wildrye (*Elymus virginiana*) are perennial cool season plants that occur on the site. A variety of forbs, including Illinois bundleflower, Maximilian sunflower, Engelmann's daisy, and black sampson (*Echinacea angustifolia*) occur in this community. Grazing prescriptions that permit acceptable grazing periods and allow adequate rest periods along with prescribed fire every three to five years are important in the maintenance of the reference plant community and the prairie landscape structure. Continuous overgrazing, or over-rest and the absence of fire, tend to favor a vegetative shift towards woody species such as mesquite, elm, hackberry, cedar, and locust (*Gleditsia triacanthos*). Without corrective measures, this shift will continue to a mesquite-shrub dominated community.

Table 6. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	3825	5100	6375
Forb	450	600	750
Tree	180	240	300
Shrub/Vine	45	60	75
Total	4500	6000	7500

Table 7. Ground cover

Tree foliar cover	0-10%
Shrub/vine/liana foliar cover	0-10%
Grass/grasslike foliar cover	30-60%
Forb foliar cover	5-10%
Non-vascular plants	0%
Biological crusts	0%
Litter	10-20%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	10-20%

Table 8. Soil surface cover

Tree basal cover	0-10%
Shrub/vine/liana basal cover	0-5%
Grass/grasslike basal cover	10-30%
Forb basal cover	1-3%
Non-vascular plants	0%
Biological crusts	0%
Litter	30-40%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	10-20%

Table 9. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	—	0-5%
>0.5 <= 1	—	—	5-10%	0-5%
>1 <= 2	—	—	10-20%	0-5%
>2 <= 4.5	—	0-10%	20-40%	5-10%
>4.5 <= 13	—	—	0-5%	0-5%
>13 <= 40	0-10%	—	—	—
>40 <= 80	—	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

Community 1.2

Mesquite/Midgrass Savannah



The Mesquite/Midgrass Savannah Community develops in the absence of fire or mechanical or chemical brush management treatments. It is usually the result of abandonment following cropping or yearly continuous grazing. Mesquite is the dominant woody plant on the site but eastern red cedar, prickly ash (*Zanthoxylum clava-herculis*), cedar elm, bumelia (*Sideroxylon lanuginosum*), coralberry (*Symphoricarpos orbiculatus*), and netleaf hackberry may

also occur. Remnants of little bluestem and Indiangrass may still occur, but the herbaceous component of the community becomes dominated by lesser producing grasses and forbs. Silver bluestem, windmillgrass, white tridens, fall witchgrass (*Digitaria cognata*), threeawn (*Aristida* sp.), Texas wintergrass, prickly pear (*Opuntia* sp.), tasajillo (*Opuntia leptocaulis*), western ragweed (*Ambrosia psilostachya*), croton (*Croton* sp.), annual broomweed, and snow on the prairie are common occupants. If the woody canopy has not exceeded 40 percent, prescribed burning on a three to five year interval in conjunction with prescribed grazing is a viable option for returning this community to a tallgrass prairie that may resemble the reference plant community. If the woody canopy has exceeded 40 percent, chemical or mechanical brush control must be applied to move the transitional community back towards the reference plant community.

Table 10. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	1200	1800	2400
Shrub/Vine	450	675	900
Forb	300	450	600
Tree	150	225	300
Total	2100	3150	4200

Table 11. Ground cover

Tree foliar cover	10-40%
Shrub/vine/liana foliar cover	10-20%
Grass/grasslike foliar cover	30-40%
Forb foliar cover	20-30%
Non-vascular plants	0%
Biological crusts	0%
Litter	5-20%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	20-40%

Table 12. Soil surface cover

Tree basal cover	0%
Shrub/vine/liana basal cover	5-15%
Grass/grasslike basal cover	10-20%
Forb basal cover	5-10%
Non-vascular plants	0%
Biological crusts	0%
Litter	5-20%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	20-40%

Table 13. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	5-10%	—
>0.5 <= 1	—	—	5-10%	—
>1 <= 2	—	—	20-40%	5-15%
>2 <= 4.5	—	—	5-15%	5-15%
>4.5 <= 13	—	10-20%	—	—
>13 <= 40	20-40%	—	—	—
>40 <= 80	—	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

Pathway 1.1A Community 1.1 to 1.2



Tallgrass Prairie




Mesquite/Midgrass Savannah

The Tallgrass Prairie Community will shift to the Mesquite/Midgrass Savannah Community when there is continued growing-season stress on reference grass species. These stresses include improper grazing management that creates insufficient critical growing-season deferment, excess intensity of defoliation, repeated, long-term growing-season defoliation, long-term drought, and/or other repeated critical growing-season stress. Increaser species (midgrasses and woody species) are generally endemic species released by disturbance. Woody species canopy exceeding 20 percent and a species shift from tallgrass to midgrass composition indicate a transition. The driver for the shift is heavy grazing, no fire, no brush management, and/or abandonment.

Pathway 1.2A Community 1.2 to 1.1



Mesquite/Midgrass Savannah



Tallgrass Prairie

The Mesquite/Midgrass Savannah Plant Community will return to the Tallgrass Prairie Plant Community under grazing management that provides sufficient critical growing season deferment in combination with proper grazing intensity as long as the seedbank or seed source is still present. Favorable moisture conditions will facilitate or accelerate this transition. The understory component may return to dominance by tallgrasses in the absence of fire or brush control. However, reduction of the woody component to reference conditions of 10 percent or less canopy cover will require inputs of fire or brush control.

State 2 Shrubland

Two communities exist in the Shrubland State: the 2.1 Mesquite/Midgrass Shrubland Community and the 2.2 Mesquite/Texas Wintergrass Woodland Community. Community 2.1 is characterized by an increase in shade-tolerant grass and an overstory canopy of greater than 40 percent. Community 2.2 is characterized by a cool-season grasses, especially Texas wintergrass, and a greater than 40 percent canopy cover.

Community 2.1
Mesquite/Midgrass Shrubland

This is a transitional community between the prairie, pastureland, or cropland states and the mesquite-woodland state. It develops in the absence of fire or mechanical or chemical brush management treatments. It is usually the result of abandonment following cropping or yearly continuous grazing. Mesquite is the dominant woody plant on the site but eastern red cedar, prickly ash, cedar elm, bumelia, coralberry, and netleaf hackberry may also occur. Remnants of little bluestem and Indiangrass may still occur, but the herbaceous component of the community becomes dominated by lesser producing grasses and forbs. Silver bluestem, windmillgrass, white tridens, fall witchgrass, threeawn, Texas wintergrass, prickly pear, tasajillo, western ragweed, croton, annual broomweed, and snow on the prairie are common occupants. Since the woody canopy has exceeded 40 percent, chemical or mechanical brush control must be applied to move the transitional community back towards the reference plant community.

Table 14. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	900	1300	1800
Shrub/Vine	400	600	800
Tree	375	560	750
Forb	200	325	450
Total	1875	2785	3800

Community 2.2
Mesquite/Texas Wintergrass Woodland



This plant community is dominated by mesquite but may include some live oak, eastern red cedar, hackberry, cedar elm, bumelia, and prickly ash. Canopy cover exceeds 40 percent. Understory shrubs and vines include coralberry, greenbriar (*Smilax* spp.), grape (*Vitis* spp.), prickly pear, and tasajillo. Texas wintergrass, silver bluestem, cedar sedge (*Carex planostachys*), threeawn, annual broomweed, and snow on the prairie are the most common herbaceous components. Herbaceous species composition and production are directly related to woody canopy cover. If the site is not abandoned cropland, chemical brush control combined with prescribed grazing and prescribed burning is a viable treatment option for moving this community back towards the reference plant community. Mechanical brush control and seeding is usually the most viable treatment option when the objective is to return this state to a community that resembles the reference plant community.

Table 15. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	650	925	1300
Tree	600	900	1200
Shrub/Vine	350	525	700
Forb	100	200	300
Total	1700	2550	3500

Table 16. Ground cover

Tree foliar cover	40-60%
Shrub/vine/liana foliar cover	20-40%
Grass/grasslike foliar cover	10-20%
Forb foliar cover	15-30%
Non-vascular plants	0%
Biological crusts	0%
Litter	5-10%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	20-40%

Table 17. Soil surface cover

Tree basal cover	20-40%
Shrub/vine/liana basal cover	15-25%
Grass/grasslike basal cover	5-10%
Forb basal cover	5-10%
Non-vascular plants	0%
Biological crusts	0%
Litter	5-10%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	20-40%

Table 18. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	5-10%	5-10%
>0.5 <= 1	—	—	—	—
>1 <= 2	—	—	5-10%	10-20%
>2 <= 4.5	—	—	5-10%	—
>4.5 <= 13	—	15-25%	—	—
>13 <= 40	40-60%	—	—	—
>40 <= 80	—	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

Pathway 2.1A

Community 2.1 to 2.2

The Mesquite/Midgrass Shrubland Community will transition to the Mesquite/Texas Wintergrass Woodland Community with continued heavy grazing, no fire, no brush management and/or abandonment. Indicators of the transition include an increased species composition of cool-season grasses and trees becoming more mature.

State 3

Converted Land

The Converted Land State contains one community, the 3.1 Converted Land Community. The state is characterized by the land manager farming crops or planted grasses.

Community 3.1

Converted Land



Conversion of the tallgrass prairie to cropland (primarily cotton, corn, and grain sorghum) began in the middle 1800's and continued into the early 1900's. Much of the prairie is in cropland today and includes additional crops of small grains (for grain and livestock grazing), soybeans, and rice. While restoration of this site to a semblance of the tallgrass prairie is possible with seeding, prescribed grazing, and prescribed burning - complete restoration of the reference plant community in a reasonable time is unlikely. If cropping is abandoned, this site is usually planted to introduced grasses and forbs and managed as pastureland. Following crop production or brush control, this site is often planted to native or introduced grasses and legumes for livestock grazing or hay production. Typical species planted include improved Bermudagrass (*Cynodon* spp.) varieties, dallisgrass (*Paspalum dilatam*), switchgrass, kleingrass (*Panicum coloratum*), old-world bluestem (*Bothriochloa* spp.) varieties, and various legume species. Many of the introduced species (Bermudagrass, dallisgrass, old-world bluestem) are invasive - moving by wind, water, and animals. Once established, they are extremely difficult to remove and will hinder the re-establishment of native species. The establishment and maintenance of these species requires cultivation, fertilization, weed control,

and prescribed grazing management. Without the yearly application of these cultural practices, this vegetative state will move towards the Shrubland State.

Table 19. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	3000	6500	10000
Total	3000	6500	10000

Transition T1A

State 1 to 2

The Prairie/Savannah State will transition to the Shrubland State when continued heavy grazing pressure, no fire, no brush management, and/or field abandonment continues. The transition is evident when woody species canopy cover exceeds 40 percent and grasses shift composition to more shade-tolerant species.

Transition T1B

State 1 to 3

The transition to the Converted State occurs when the site is plowed for planting crops or pasture. The driver for the transition is the land manager's decision to farm the site.

Restoration pathway R2A

State 2 to 1

Restoration of the Shrubland State to the Prairie/Savannah State requires substantial energy input. Mechanical or herbicidal brush control treatments can be used to remove woody species. A long-term prescribed fire program may sufficiently reduce brush density particularly if the woody component is dominated by species that are not re-sprouters. However, fire may not be sufficient to remove mature trees. Brush control in combination with prescribed fire, proper grazing management, and favorable growing conditions may be the most economical means of creating and maintaining the desired plant community. If remnant populations of tallgrasses, midgrasses, and desirable forbs are not present at sufficient levels, range planting will be necessary to restore the reference plant community.

Transition T2A

State 2 to 3

The transition to the Converted State occurs when the site is plowed for planting crops or pasture. The driver for the transition is the land manager's decision to farm the site.

Restoration pathway R3A

State 3 to 1

The restoration to State 1 can occur when the land manager ceases agronomic practices. Range planting of native species found in the reference community will be required to bring back a similar community as the State 1 plant composition. The extent of previous soil disturbances will determine how much seedbed preparation will be needed, as well as the ability to be restored. Proper grazing and brush management will be required to ensure success.

Transition T3A

State 3 to 2

The Converted Land State will transition to the Shrubland State by heavy continuous grazing, no fire, no brush management and/or abandonment. Woody species that encroach will eventually grow into the overstory and begin to shade out grasses.

Additional community tables

Table 20. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass/Grasslike					
1	Tallgrass			1500–3000	
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	1500–3000	–
2	Tallgrasses			750–1500	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	750–1500	–
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	750–1500	–
3	Mid/Shortgrasses			300–600	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	300–600	–
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	300–600	–
	silver beardgrass	BOLAT	<i>Bothriochloa laguroides</i> ssp. <i>torreyana</i>	300–600	–
	sedge	CAREX	<i>Carex</i>	300–600	–
	cylinder jointtail grass	COCY	<i>Coelorachis cylindrica</i>	300–600	–
	fall witchgrass	DICO6	<i>Digitaria cognata</i>	300–600	–
	Virginia wildrye	ELVI3	<i>Elymus virginicus</i>	300–600	–
	Texas cupgrass	ERSE5	<i>Eriochloa sericea</i>	300–600	–
	Texas wintergrass	NALE3	<i>Nassella leucotricha</i>	300–600	–
	Florida paspalum	PAFL4	<i>Paspalum floridanum</i>	300–600	–
	Hall's panicgrass	PAHA	<i>Panicum hallii</i>	300–600	–
	vine mesquite	PAOB	<i>Panicum obtusum</i>	300–600	–
	switchgrass	PAVI2	<i>Panicum virgatum</i>	300–600	–
	marsh bristlegrass	SEPA10	<i>Setaria parviflora</i>	300–600	–
	composite dropseed	SPCOC2	<i>Sporobolus compositus</i> var. <i>compositus</i>	300–600	–
	white tridens	TRAL2	<i>Tridens albescens</i>	300–600	–
	eastern gamagrass	TRDA3	<i>Tripsacum dactyloides</i>	300–600	–
	longspike tridens	TRST2	<i>Tridens strictus</i>	300–600	–
Forb					
4	Forbs			300–600	
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	300–600	–
	groovestem Indian plantain	ARPL4	<i>Arnoglossum plantagineum</i>	300–600	–
	bluebell bellflower	CARO2	<i>Campanula rotundifolia</i>	300–600	–
	compact prairie clover	DACOC	<i>Dalea compacta</i> var. <i>compacta</i>	300–600	–
	Illinois bundleflower	DEIL	<i>Desmanthus illinoensis</i>	300–600	–
	blacksamson echinacea	ECAN2	<i>Echinacea angustifolia</i>	300–600	–
	Engelmann's daisy	ENPE4	<i>Engelmannia peristenia</i>	300–600	–
	button eryngo	ERYU	<i>Eryngium yuccifolium</i>	300–600	–
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	300–600	–
	coastal indigo	INMI	<i>Indigofera miniata</i>	300–600	–
	pinkscale blazing star	LIEL	<i>Liatris elegans</i>	300–600	–
	Nuttall's sensitive-briar	MINU6	<i>Mimosa nuttallii</i>	300–600	–
	whitest evening	OEAL	<i>Oenothera albicaulis</i>	300–600	–

	primrose				
	Nuttall's prairie parsley	PONU4	<i>Polytaenia nuttallii</i>	300–600	–
	prairie snoutbean	RHLA5	<i>Rhynchosia latifolia</i>	300–600	–
	fuzzybean	STROP	<i>Strophostyles</i>	300–600	–
Shrub/Vine					
5	Vines			30–60	
	coralberry	SYOR	<i>Symphoricarpos orbiculatus</i>	30–60	–
Tree					
6	Trees			150–300	
	netleaf hackberry	CELAR	<i>Celtis laevigata</i> var. <i>reticulata</i>	150–300	–
	gum bully	SILAL3	<i>Sideroxylon lanuginosum</i> ssp. <i>lanuginosum</i>	150–300	–
	cedar elm	ULCR	<i>Ulmus crassifolia</i>	150–300	–

Table 21. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass/Grasslike					
1	Small Grains			6000–8000	
	common oat	AVSA	<i>Avena sativa</i>	6000–8000	–
	Italian ryegrass	LOPEM2	<i>Lolium perenne</i> ssp. <i>multiflorum</i>	6000–8000	–
	common wheat	TRAЕ	<i>Triticum aestivum</i>	6000–8000	–
2	Pastureland Species			3000–10000	
	Bermudagrass	CYDA	<i>Cynodon dactylon</i>	3000–10000	–
	kleingrass	PACO2	<i>Panicum coloratum</i>	3000–10000	–
	bahiagrass	PANO2	<i>Paspalum notatum</i>	3000–10000	–

Animal community

As a historic tall grass prairie, this site was occupied by bison, antelope, deer, prairie chickens, quail, turkey, and dove. This site was also used by many species of grassland songbirds, migratory waterfowl, coyotes, and wolves. This site now provides forage for livestock and is still used by quail, dove, migratory waterfowl, migratory and resident grassland birds, and coyotes. Deer use the site when mesquite and associated woody species occur in sufficient densities and proper structure to provide adequate cover.

Hydrological functions

Peak rainfall periods occur in May and June from frontal passage thunderstorms and in September and October from tropical systems, as well as frontal passage thunderstorms. Rainfall amounts may be high (three to five inches per event) and events may be intense. The site is subject to erosion where adequate herbaceous cover is not maintained and on heavy use areas such as roads and livestock trails. Extended periods (60 days) of little to no rainfall during the growing season are common.

Recreational uses

The site is used for hunting, fishing, and photography.

Wood products

Mesquite may be harvested for firewood, charcoal, and specialty uses including furniture, gun stocks, and veneer.

Other products

Hay, seed from native grasses, forbs, and legumes

Inventory data references

These site descriptions were developed as part a Provisional Ecological Site project using historic soil survey manuscripts, available site descriptions, and low intensity field traverse sampling. Future work to validate the information is needed. This will include field activities to collect low, medium, and high-intensity sampling, soil correlations, and analysis of that data. A final field review, peer review, quality control, and quality assurance review of the will be needed to produce the final document.

Other references

Diggs, G.M., B.L. Lipscomb, and R.J. O'Kennon. 1999. Illustrated Flora of North Central Texas Botanical Research Institute of Texas and Austin College, Fort Worth, TX.

Hatch, S.L, K.N. Gandhi, and L.E. Brown. 1990. Checklist of the vascular plants of Texas. Texas Agricultural Experiment Station, College Station, TX.

Contributors

Mark Moseley
Tyson Hart

Approval

Bryan Christensen, 9/21/2023

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Mike Stellbauer
Contact for lead author	Mike Stellbauer, Zone RMS, NRCS, Bryan, Texas
Date	05/23/2005
Approved by	Bryan Christensen
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** A few rills of small extent may naturally occur on this site.

-
2. **Presence of water flow patterns:** Some water flow patterns are normal for this site due to landscape position and slope but should be vegetated and stable.
-

3. **Number and height of erosional pedestals or terracettes:** Pedestals or terracettes are uncommon for this site when occupied by the reference community.
-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Expect no more than 20 percent bare ground distributed in small patches.
-
5. **Number of gullies and erosion associated with gullies:** No gullies should be present. Drainageways should be stable and covered with vegetation.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** None.
-
7. **Amount of litter movement (describe size and distance expected to travel):** This site has slowly permeable soils. On sloping sites, small to medium-size litter will move short distances with intense storms.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil surface is resistant to erosion. Stability class range is expected to be 5 to 6.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface structure is greater than 60 inches. Colors range from black to very dark brown and moderate fine to medium subangular block structure. SOM is one to three percent.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** This prairie site is dominated by tallgrasses and forbs, with adequate litter and little bare ground, which provides for maximum infiltration and little runoff under normal rainfall events.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No evidence of compaction.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Warm-season tallgrasses >>
- Sub-dominant: Warm-season midgrasses > Cool-season midgrasses > Forbs >
- Other: Trees > Shrubs/Vines
- Additional:
-

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** There should be little mortality or decadence for any functional group for the reference community.
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 4,500 pounds per acre during below average moisture years to 7,500 pounds per acre during above average moisture years.
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Potential invasive species include yellow bluestems, common Bermudagrass, mesquite, elm, huisache, eastern red cedar, and osage orange.
-

17. **Perennial plant reproductive capability:** All perennial plants should be capable of reproducing, except during drought conditions, heavy natural herbivory, and intense wildfires.
-