

Ecological site R102CY046NE Subirrigated

Last updated: 12/10/2024
Accessed: 05/11/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Classification relationships

"Subirrigated" range sites for NE NRCS Vegetation Zones 3 & 4

NE Natural Heritage Program/NE Game & Parks Commission: "Lowland Tallgrass Prairie"

General information for MLRA 102C:

Fenneman (1916) Physiographic Regions

Division - Interior Plains

East:

Province - Central Lowland

Section - Till Plains

West:

Province - Great Plains

Section - High Plains

USFS (2007) Ecoregions

Domain - Humid Temperate

Division - Prairie

Province - Prairie Parkland (Temperate)
Section - North-Central Glaciated Plains (251B)

EPA Ecoregions (Omernik 1997)

I - Great Plains (9)

II - Temperate Prairies (9.2)

III - Western Corn Belt Plains (9.2.3) IV - Loess Prairies (47a)

IV - Northeastern Nebraska Loess Hills (47k)

IV - Transitional Sandy Plain (47l)

Ecological site concept

This site has a seasonally high water table from 51 to 102 centimeters (1.5 to 3.5 feet), with additional moisture received from higher adjacent areas as run-on. This increases plant production while also “buffering” variability caused by fluctuating weather conditions. However, relatively minor changes in local elevation can dramatically affect the plant community and this site often occurs in a complex with other sites straddling this water table depth range.

Associated sites

R102CY044NE	WET LAND This site typically occurs in the lowest areas, or where hydrology otherwise supports a community heavily dominated by hydrophytic vegetation.
R102CY045NE	WET SUBIRRIGATED This site occurs on lower relief with a seasonally high water table within 24
R102CY048NE	Loamy Overflow This site occurs on surrounding higher areas with significantly lower production.

Similar sites

R102CY044NE	WET LAND This site is saturated at or near the surface and often ponded. Gleying is common.
R102CY048NE	Loamy Overflow This site does not show evidence of a seasonally high water table within 42
R102CY045NE	WET SUBIRRIGATED This site is seasonally saturated 0-24

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Andropogon gerardii</i> (2) <i>Schizachyrium scoparium</i>

Physiographic features

This site mainly occurs on nearly level to gently sloping floodplains and interdunes on sandhill valleys (0-3% slopes). A few areas are in swales, stream terraces, alluvial fans, and on foot slopes. It predominantly receives runoff from adjacent sites, has a seasonally high water table from 60 to 102 centimeters (24 to 40 inches) from November-May, does not pond, and may flood occasionally for a brief duration.

Table 2. Representative physiographic features

Landforms	(1) Flood plain (2) Interdune (3) Swale
-----------	---

Runoff class	Negligible to medium
Flooding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Flooding frequency	None to occasional
Elevation	600–1,500 ft
Slope	0–3%
Water table depth	17–40 in
Aspect	Aspect is not a significant factor

Climatic features

Most of the rainfall occurs as high-intensity, convective thunderstorms during the growing season. Peak precipitation occurs from the middle of spring to early in autumn. Winter precipitation occurs as snow (USDA/NRCS 2006).

The average annual temperature gradient trends higher from north (45°F/7°C) to south (51°F/11°C).

The average annual precipitation gradient trends higher from northwest (25 inches / 64 centimeters) to southeast (31 inches / 79 centimeters).

The annual snowfall ranges from about 60 centimeters (24 inches) in the southern part of the area to 85 centimeters (34 inches) in the northern part.

The following data summary includes weather stations representing the full geographic extent of the MLRA, and is based on 70% probabilities (NOAA/UNL) meaning that actual observed climate conditions may fall outside these ranges 30% of the time. Furthermore, climatic events can manifest many different ways. For example, abnormally dry periods could occur as 3 consecutive drought years out of 10, 3 individual years separated by “normal” years, or some combination. Tree-ring records indicate that portions of the Great Plains have also historically experienced droughts lasting several decades, so plant community response will largely depend on the manner in which climatic variability is realized in interaction with past and current land management.

Table 3. Representative climatic features

Frost-free period (characteristic range)	127-134 days
Freeze-free period (characteristic range)	144-156 days
Precipitation total (characteristic range)	27-29 in
Frost-free period (actual range)	124-139 days
Freeze-free period (actual range)	143-166 days
Precipitation total (actual range)	27-30 in
Frost-free period (average)	131 days
Freeze-free period (average)	151 days
Precipitation total (average)	28 in

Climate stations used

- (1) GENOA 2 W [USC00253185], Genoa, NE
- (2) CREIGHTON [USC00251990], Creighton, NE
- (3) FREMONT [USC00253050], Fremont, NE
- (4) FLANDREAU [USC00392984], Flandreau, SD

Influencing water features

The soil profile is endosatuated by an unperched water table produced by lateral subsurface flow from surrounding higher areas and/or adjacent water bodies.

Soil features

These are predominantly very deep, somewhat poorly to moderately well drained soils. The surface texture is predominantly silt loam, loam, or silty clay loam from 0 to 18 centimeters (0 to 7 inches) and the Subsurface Texture Groups are Loamy or Sandy from 18 to 203 centimeters (7 to 80 inches).

Rills, gullies, and water flow patterns are not inherent to this site. Pedestalling is none to slight. Soil aggregate stability should be high.

Major soils assigned to this site include Ackmore, Boel, Coleridge, Els, Elsmere, Gibbon, Lamo, Ord, Primghar, Spillco, Splitrock, Wann

Table 4. Representative soil features

Surface texture	(1) Silty clay loam (2) Silt loam (3) Loam
Family particle size	(1) Loamy
Drainage class	Somewhat poorly drained to moderately well drained
Permeability class	Slow to very rapid
Soil depth	80 in
Surface fragment cover <=3"	0–7%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	1.9–9.4 in
Calcium carbonate equivalent (0-40in)	0–3%
Electrical conductivity (0-40in)	0–2 mmhos/cm
Sodium adsorption ratio (0-40in)	0–5
Soil reaction (1:1 water) (0-40in)	5.6–9
Subsurface fragment volume <=3" (Depth not specified)	0–8%
Subsurface fragment volume >3" (Depth not specified)	0–3%

Ecological dynamics

The foremost driver influencing this site is a water table that serves to bolster production, especially in times of reduced precipitation. Relatively minor changes in local elevation can dramatically affect the plant community, and this site often occurs in a complex with other sites straddling this water table depth range, particularly Loamy Lowland, Loamy Overflow, and Wet Subirrigated. Plant community composition may also experience similar changes through disturbances that affect the water table itself, such as extended dry or wet cycles. Local and regional anthropogenic factors can further influence the water table through drainage, flow regulation, stream channelization, etc.

This site developed with occasional fires being part of the ecological processes. It is presumed that the historic fires generally occurred every 3-4 years and ameliorated the relatively rapid accumulation of excessive litter. It is also

believed that pre-European inhabitants may have used fire as a management tool for attracting herds of large migratory herbivores (bison, elk, and/or deer.) The impact of fire over the past 100 years has been relatively insignificant due to the human control of wildfires and the lack of acceptance of prescribed fire as a management tool.


The degree of herbivory (feeding on herbaceous plants) has a significant impact on the dynamics of the site. Historically, periodic grazing by herds of large migratory herbivores was a primary influence. Secondary influences of herbivory by species such as grasshoppers and root feeding organisms impacted the vegetation historically, and continue to this day. The management of herbivory by humans through grazing of domestic livestock and/or manipulation of wildlife populations has been a major influence on the ecological dynamics of the site. This management coupled with climate largely dictates the plant communities for the site.

The plant community for this site is dynamic due to the complex interaction of many ecological processes. The interpretive plant community for this site is the reference state. The reference state has been determined by the study of rangeland relic areas, areas protected from excessive disturbance and areas under compatible grazing strategies. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.


The following is a diagram that illustrates the common plant communities that can occur on the site and the pathways among communities. The ecological processes will be discussed in more detail in the plant community descriptions following the diagram.

State and transition model

Ecosystem states



States 1, 5 and 2 (additional transitions)



T1A - Colonization of cool season grasses

T1B - Major disturbance of site

T1C - Woody encroachment

R2A - Eradication of cool season grasses

T2A - Invasive grasses dominant

T2B - Site disturbance allows annuals to dominate

T2C - Woody species dominate due to absence of fire or grazing


T3A - Management inputs to restore native species dominance

T3A - Woody species gain dominance due to lack of fire and/or grazing.

R4A - Management inputs to restore native species dominance.

R5A - Restoration of site to native, tallgrass species.


State 1 submodel, plant communities



1.1A - Heavy grazing without recovery periods

1.2A - Reduced grazing pressure


State 2 submodel, plant communities




2.1A - Native species dominant but cool season grasses established on site

2.2A - Cool season grasses increase; native species still dominant


State 3 submodel, plant communities



State 4 submodel, plant communities



State 5 submodel, plant communities



State 1 Native tallgrass

This state comprises the communities within the range of natural variability under historic conditions and disturbance regimes. Patterns created by wildlife use and fire would have created a mosaic of communities across the landscape; however, warm-season tallgrasses are dominant, with a subdominant to minor contribution from

native cool-season grasses, forbs, and shrubs. Fire and bison herbivory were the dominant disturbance regimes that historically maintained the tallgrass dominance with a diverse forb component. Furthermore, bison grazing was closely linked to fire patterns as the animals preferred grazing burned areas offering lush regrowth devoid of decadence and of higher nutritive quality. Thus, historic plant communities were subjected to occasional burning and grazing, with substantial rest/recovery periods as the fuel load rebuilt to eventually start this process again. Fire return intervals of 3-4 years served to suppress woody species, particularly the various deciduous tree and shrub species prevalent in adjacent riparian corridors. The degree to which observed conditions represent this state largely depends on how closely the management has mimicked these past disturbance effects.

Dominant plant species

- big bluestem (*Andropogon gerardii*), grass
- little bluestem (*Schizachyrium scoparium*), grass

Community 1.1

Big bluestem-Little bluestem (*Andropogon gerardii*-*Schizachyrium scoparium*)



Figure 8. Subirrigated 1.1

This is the interpretive plant community and can be found on areas that are properly managed with prescribed grazing that allows for adequate recovery periods following each grazing event. The plant community consists of 80-95% grasses and grass-like, 5-10% forbs and 0-5% shrubs. Dominant grasses include big bluestem, indiangrass, and switchgrass. Other grasses and grass-like are little bluestem, sideoats grama, western wheatgrass, and sedges. Forb species are diverse and often include western ragweed and Missouri goldenrod. This plant community is diverse, stable, and productive. Plant community dynamics, nutrient cycles, water cycles, and energy flow are functioning properly. Plant litter is properly distributed with negligible movement off-site and natural plant mortality is very low. This community is resistant to many disturbances except continuous, season-long heavy grazing, tillage, or non-use. Broadcast herbicide application will dramatically reduce forb diversity and abundance. Total annual production, during an average year, ranges from 4300 to 5600 pounds per acre air-dry weight and will average 5100 pounds.

Dominant plant species

- big bluestem (*Andropogon gerardii*), grass
- little bluestem (*Schizachyrium scoparium*), grass

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	4115	4590	4710
Forb	220	383	610
Shrub/Vine	0	128	290
Total	4335	5101	5610

Figure 10. Plant community growth curve (percent production by month). NE1021, 102C Warm-season. Warm-season grass, MLRA 102C.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	2	7	18	23	26	16	6	2	0	0

Community 1.2
 Prairie cordgrass-Big bluestem (*Spartina pectinata*-*Andropogon gerardii*)



Figure 11. Subirrigated 1.2

Prairie cordgrass has replaced big bluestem as the dominant species. Other species, such as reed canarygrass, little bluestem, and western wheatgrass have also increased. While still within the range of natural variability, energy capture, nutrient cycling, and hydrology are not functioning at their full potential relative to the reference condition.


Dominant plant species

- prairie cordgrass (*Spartina pectinata*), grass
- big bluestem (*Andropogon gerardii*), grass

Figure 12. Plant community growth curve (percent production by month). NE1021, 102C Warm-season. Warm-season grass, MLRA 102C.


Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	2	7	18	23	26	16	6	2	0	0

Pathway 1.1A
 Community 1.1 to 1.2



Grazing management which does not provide adequate recovery periods will cause a shift from big bluestem and Indiangrass towards less palatable species, particularly prairie cordgrass.

Pathway 1.2A
 Community 1.2 to 1.1



Prairie cordgrass-Big bluestem (*Spartina pectinata*-*Andropogon gerardii*)

Big bluestem-Little bluestem (*Andropogon gerardii*-*Schizachyrium scoparium*)

Management that provides adequate recovery periods and does not annually prevent tallgrass seedset or otherwise impair vigor will facilitate a return to community phase 1.1. In the case of drought, the return to more typical precipitation patterns will promote shift towards tallgrass species.

State 2

Native/invaded mix

This state can manifest three ways: 1) the appearance of introduced cool-season grasses, 2) the expansion of deciduous shrubs and/or trees, or 3) some combination of these. Kentucky bluegrass and smooth brome are the primary cool-season grass invaders in this region, commonly found in roadsides, disturbed areas, and pastures intentionally seeded for cool-season forage. Management practices and/or environmental conditions that are not favorable to native grass vigor may allow introduced grasses to invade the site thereby decreasing native diversity and abundance, particularly of forbs. While reed canarygrass is a native, it may act in much the same way as the introduced species. In the absence of the historic fire regime, woody deciduous species may also expand to become an influential component of the community. The invasive component tends to have very high resilience, is extremely difficult to eradicate, and what might be considered a new "contemporary" range of natural variability is seen as competition between the native grasses and introduced/woody species for space and resources.

Dominant plant species

- big bluestem (*Andropogon gerardii*), grass
- little bluestem (*Schizachyrium scoparium*), grass
- smooth brome (*Bromus inermis*), grass
- reed canarygrass (*Phalaris arundinacea*), grass

Community 2.1

Subdominant Smooth brome-Reed canarygrass (*Bromus inermis*-*Phalaris arundinacea*)

While native warm-season grasses still dominate the site, introduced cool-season species have established a foothold in the system and can be found interspersed throughout the stand. The stand may still have a native tallgrass appearance overall, but brome and/or reed canarygrass can be easily found. Deciduous shrub/tree species may also have begun to expand into areas where they did not persist historically, but the overall appearance can vary depending on the propagation method of a particular species.

Dominant plant species

- big bluestem (*Andropogon gerardii*), grass
- little bluestem (*Schizachyrium scoparium*), grass
- reed canarygrass (*Phalaris arundinacea*), grass
- smooth brome (*Bromus inermis*), grass

Figure 13. Plant community growth curve (percent production by month). NE1022, Warm-season dominant, cool-season subdominant.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	2	9	19	23	24	13	7	3	0	0

Community 2.2

Codominant Smooth brome-Reed canarygrass (*Bromus inermis*-*Phalaris arundinacea*)



Figure 14. Subirrigated 2.2

This community is comprised of a relatively even mix of native grasses and invasive species overall. This may manifest as a well-distributed interspersion of natives and invaders, as distinct patches wherein competitors dominate locally, or some combination. Forb diversity and abundance is further diminished.

Dominant plant species

- big bluestem (*Andropogon gerardii*), grass
- reed canarygrass (*Phalaris arundinacea*), grass
- little bluestem (*Schizachyrium scoparium*), grass
- smooth brome (*Bromus inermis*), grass

Figure 15. Plant community growth curve (percent production by month). NE1023, Warm-season, cool-season codominant.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	10	23	26	16	10	7	4	1	0

Pathway 2.1A
Community 2.1 to 2.2

Management and/or environmental conditions have afforded a persisting competitive advantage to introduced cool-season grasses, and they begin to dominate the ecological dynamics of the site. The robust invasive component is able to quickly and effectively exploit opportunities to outcompete and displace natives. Repeated summer use of an area will place the bulk of stressor impacts on native plants, reducing native vigor and allowing invaders to thrive. Likewise, a climate pattern limiting natural moisture to the spring and fall months coincides with peak cool-season growth and may support a similar process.

Pathway 2.2A
Community 2.2 to 2.1

The native component remains in an abundance that can facilitate a return towards more historic conditions if management is modified to shift stressor impacts to the invasive species, and promote warm-season grass vigor. Environmental conditions and/or disturbance regimes that strongly favor warm-season grasses can also trend the site towards the reference.

State 3
Invasive dominant

Introduced cool-season invasion has progressed to the point that native species comprise a negligible portion of the community and the aggressively rhizomatous invasives preclude native germination and seedling survival. The native component may be completely absent, and the site resembles a seeded pasture. Alternatively, the dominant invasives may be deciduous woody species. Woody competitiveness for sunlight, water, space, and other resources continues to increase as desirable herbaceous species are shaded out, crowded out, or otherwise suppressed.

Dominant plant species

- smooth brome (*Bromus inermis*), grass
- reed canarygrass (*Phalaris arundinacea*), grass

Community 3.1
Smooth brome-Reed Canarygrass (*Bromus inermis*-*Phalaris arundinacea*)



Figure 16. Subirrigated 3.1

This community is typically composed of smooth brome with bluegrass interspersed among the brome tillers. Warm-season natives, if present, are sparse yet often conspicuous due to pronounced differences in growth habits and metabolic pathways. Community structure and function have been dramatically simplified relative to the reference condition, and very few biotic functional groups are represented in amounts that would influence ecological function. The invasive grass root skein provides good site stability; however, replacement of the deeper roots and complex bunchgrass canopy with the shallower roots and erect tiller canopy of the invaders results in reduced interception and infiltration rates.

Dominant plant species

- smooth brome (*Bromus inermis*), grass
- reed canarygrass (*Phalaris arundinacea*), grass

Figure 17. Plant community growth curve (percent production by month).
NE1024, Cool-season. Smooth brome/Kentucky bluegrass.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	4	13	25	28	8	6	9	5	2	0

State 4
Annual/pioneer

Nutrient cycling, hydrologic function, and/or soil stability have been severely altered, and possibly compromised. This is a highly variable state in which the specific plants observed will depend largely on the original community and the nature of the disturbance. This condition encompasses (but is not necessarily limited to) events such as severe fire impacts, heavy continuous grazing, heavy nutrient inputs, and abandoned cropland.

Dominant plant species

- ragweed (*Ambrosia*), other herbaceous
- hoary verbena (*Verbena stricta*), other herbaceous
- pigweed (*Amaranthus*), other herbaceous
- Canada thistle (*Cirsium arvense*), other herbaceous

Community 4.1

Variable native and introduced



Figure 18. Subirrigated 4.1

This community is heavily dominated by annual plants that thrive in disturbed areas and often includes annual ragweed, hoary verbena, or amaranths. It is also particularly vulnerable to noxious weed invasion with the most common species being musk and Canada thistles. Leafy spurge becomes more common northward in the MLRA.

Dominant plant species

- ragweed (*Ambrosia*), other herbaceous
- hoary verbena (*Verbena stricta*), other herbaceous
- pigweed (*Amaranthus*), other herbaceous
- Canada thistle (*Cirsium arvense*), other herbaceous

State 5

Woody dominant

Under historic disturbance regimes, frequent and uncontrolled fire and wildlife browsing served to keep woody species in check. However, in the absence of fire (either wild or prescribed), it's not uncommon for the woody trees and shrubs normally limited to riparian areas to expand into the floodplains, regardless of herbaceous community composition. Wildlife may introduce a seed source to areas not associated with a waterway, such as interdunal depressions.

Dominant plant species

- maple (*Acer*), tree
- eastern cottonwood (*Populus deltoides*), tree
- boxelder (*Acer negundo*), tree
- green ash (*Fraxinus pennsylvanica*), tree

Community 5.1

Native deciduous overstory



Figure 19. Subirrigated 5.1a



Figure 20. Subirrigated 5.1b early encroachment

Deciduous woody species have encroached and established, typically with species such as maples, cottonwood, boxelder, green ash, and swamp oak. Eastern redcedar may also establish, but is not usually as dominant as seen on drier sites.

Dominant plant species

- maple (*Acer*), tree
- eastern cottonwood (*Populus deltoides*), tree
- boxelder (*Acer negundo*), tree
- green ash (*Fraxinus pennsylvanica*), tree

Transition T1A

State 1 to 2

In the presence of introduced cool-season grasses, environmental conditions and/or management that reduces native vigor and stand resilience, and frees up resources (space, sunlight, nutrients, water) will allow for colonization of Kentucky bluegrass, smooth brome and/or reed canarygrass. Likewise, similar processes may also allow for woody species to expand, particularly willows and cottonwoods.

Transition T1B

State 1 to 4

There are many possible triggers for this transition that may occur as acute events (e.g. plowing) or cumulative impacts of chronic events (e.g. long-term undermanaged grazing.) The absence of deep-rooted perennial cover exposes the site to topsoil loss, open nutrient cycle, and free space which collectively allow for opportunistic annual species to dominate.

Transition T1C

State 1 to 5

All herbaceous communities are vulnerable to woody encroachment in the absence of fire and/or browsing and hoof action impacts. This is particularly prominent in areas adjacent to riparian corridors which supply a constant seed source. As tree establishment progresses, the conditions grow increasingly favorable for woody deciduous germination and growth.

Restoration pathway R2A

State 2 to 1

Eradication of introduced cool-season grasses from this site will require long-term, targeted management efforts to create an adverse environment during the spring and late fall when bluegrass and brome are most actively growing, with favorable conditions during the summer to promote native warm-season species. Targeted practices such as prescribed burning, flash grazing, and herbicide are often employed at strategic times of the year to set back undesirable species. The combination of practices should strive to mimic the historic disturbance regimes to which the desirable native species are best adapted.

Transition T2A

State 2 to 3

If the conditions which initiated and fomented the colonization and expansion of cool-season invasion are not removed or mitigated, stand composition will continue to shift in this direction and begin to resemble a monoculture of brome and/or canarygrass. Due to the dense rhizomatous root mat of brome and bluegrass, native species suffer decreasing opportunities to contribute propagules, and individual plants lost are not replaced by desirable natives.

Transition T2B

State 2 to 4

There are many possible triggers for this transition that may occur as acute events (e.g. plowing) or cumulative impacts of chronic events (e.g. long-term undermanaged grazing.) The absence of deep-rooted perennial cover exposes the site to topsoil loss, open nutrient cycle, and free space which collectively allow for opportunistic annual species to dominate.

Transition T2C

State 2 to 5

All herbaceous communities are vulnerable to deciduous encroachment in the absence of fire and/or browsing impacts. This is particularly prominent in areas adjacent to riparian corridors which supply a constant seed source. As tree establishment progresses, the conditions grow increasingly favorable for woody deciduous germination and growth.

Transition T3A

State 3 to 2

Aggressive intervening actions will be required to simultaneously recolonize native grasses and suppress vigor in undesirable species. Restoration follows the same principles as the R2A pathway, but may also require native range seeding if the latent seedbank is inadequate.

Transition T3B

State 3 to 4

Nutrient cycling, hydrologic function, and/or soil stability have been severely altered, and possibly compromised. This is a highly variable state in which the specific plants observed will depend largely on the original community and the nature of the disturbance.

Context dependence. Site disturbance resulting in annual plant dominance

Transition T3A

State 3 to 5

All herbaceous communities are vulnerable to deciduous encroachment in the absence of fire and/or browsing impacts. This is particularly prominent in areas adjacent to riparian corridors which supply a constant seed source. As tree establishment progresses, the conditions grow increasingly favorable for woody deciduous germination and growth.

Restoration pathway R4A

State 4 to 1

Restoration strategies will depend on the nature of the disturbance and the viability of the seedbank. On pastures, changes to grazing management and favorable moisture conditions may produce a perennial community. However, in abandoned cropland range seeding will likely be necessary to recolonize desirable perennial species.

Restoration pathway R5A

State 5 to 1

The combination of tree size, reduced herbaceous understory, and more mesic conditions makes it increasingly difficult for natural disturbances to restore/maintain the historic tallgrass community, and mature woodlands can no longer be restored with fire. Intensive brush management will be required to mechanically remove the established overstory. Woody control and maintenance will be an ongoing process and may also require chemical methods if sprouting species are present.

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass/Grasslike					
1	Tall warm-season			2295–3825	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	1275–2040	–
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	510–1020	–
	switchgrass	PAVI2	<i>Panicum virgatum</i>	255–765	–
	prairie cordgrass	SPPE	<i>Spartina pectinata</i>	255–510	–
2				765–1785	
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	765–1020	–
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	0–510	–
3	Cool-season			255–1020	
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	0–255	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	0–255	–
	reed canarygrass	PHAR3	<i>Phalaris arundinacea</i>	0–255	–
	porcupinegrass	HESP11	<i>Hesperostipa spartea</i>	0–255	–
	foxtail barley	HOJU	<i>Hordeum jubatum</i>	0–255	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0–105	–
	prairie wedgescale	SPOB	<i>Sphenopholis obtusata</i>	0–105	–
	needle and thread	HECO26	<i>Hesperostipa comata</i>	0–105	–
	Scribner's rosette grass	DIOLS	<i>Dichanthelium oligosanthes</i> var. <i>scribnerianum</i>	0–105	–
4	Grass-like			255–510	
	Germoid (grass or grass-like)	20PAM	Germoid (grass or grass-like)	51–255	

	Graminoid (grass or grass-like)	2GRAM	Graminoid (grass or grass-like)	0-255	—
	sedge	CAREX	<i>Carex</i>	0–255	—
	broom sedge	CASC11	<i>Carex scoparia</i>	0–255	—
	awlfruit sedge	CAST5	<i>Carex stipata</i>	0–255	—
	rush	JUNCU	<i>Juncus</i>	0–255	—
	cloaked bulrush	SCPA8	<i>Scirpus pallidus</i>	0–255	—
	common threesquare	SCPU10	<i>Schoenoplectus pungens</i>	0–255	—
Forb					
5				255–510	
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	0–255	—
	Forb (herbaceous, not grass nor grass-like)	2FORB	<i>Forb (herbaceous, not grass nor grass-like)</i>	0–255	—
	Illinois bundleflower	DEIL	<i>Desmanthus illinoensis</i>	0–255	—
	scouringrush horsetail	EQHY	<i>Equisetum hyemale</i>	0–255	—
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	0–255	—
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	0–255	—
	stiff sunflower	HEPA19	<i>Helianthus pauciflorus</i>	0–255	—
	Pennsylvania smartweed	POPE2	<i>Polygonum pensylvanicum</i>	0–255	—
	blackeyed Susan	RUHI2	<i>Rudbeckia hirta</i>	0–255	—
	giant goldenrod	SOGI	<i>Solidago gigantea</i>	0–255	—
	swamp verbena	VEHA2	<i>Verbena hastata</i>	0–255	—
	white heath aster	SYER	<i>Symphotrichum ericoides</i>	0–105	—
	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	0–105	—
	stiff goldenrod	OLRI	<i>Oligoneuron rigidum</i>	0–105	—
	Virginia strawberry	FRVI	<i>Fragaria virginiana</i>	0–105	—
	onion	ALLIU	<i>Allium</i>	0–105	—
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	0–105	—
	false boneset	BREU	<i>Brickellia eupatorioides</i>	0–105	—
	white prairie clover	DACA7	<i>Dalea candida</i>	0–105	—
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	0–105	—
Shrub/Vine					
6				0–255	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0–255	—
	leadplant	AMCA6	<i>Amorpha canescens</i>	0–255	—
	western snowberry	SYOC	<i>Symphoricarpos occidentalis</i>	0–255	—
	eastern poison ivy	TORA2	<i>Toxicodendron radicans</i>	0–175	—
	prairie rose	ROAR3	<i>Rosa arkansana</i>	0–175	—

Animal community

This site is well adapted to managed grazing by domestic livestock. The predominance of herbaceous plants across all plant community phases best lends these sites to grazing by cattle but browsing livestock such as goats or sheep that will more heavily utilize invasive forbs and brush. Carrying capacity and production estimates are conservative estimates that should be used only as guidelines in initial stages of grazing lands planning.

Often, the plant community does not entirely match any particular plant community (as described in the ecological site description). Because of this, a resource inventory is necessary to document plant composition and production.

Proper interpretation of this inventory data will permit the establishment of a safe, initial stocking rate for the type and class of animals and level of grazing management. Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide year-long forage for cattle, sheep, or horses. During the dormant period, the protein levels of the forage may be lower than the minimum needed to meet livestock (primarily cattle and sheep) requirements.

Suggested stocking rates (carrying capacity*) for cattle under continuous season-long grazing under normal growing conditions are listed below:

- 1.1 Big bluestem-Little bluestem; 5100 lbs/acre production and 1.40 AUM/acre
- 1.2 Prairie cordgrass-Big bluestem; 4050 lbs/acre production and 1.11 AUM/acre
- 2.1 Subdominant smooth brome-reed canarygrass; 4350 lbs/acre production and 1.19 AUM/acre
- 2.2 Codominant smooth brome-reed canarygrass; 4650 lbs/ac and 1.27 AUM/acre with 50% or more introduced cool-season component
- 3.1 Smooth brome-Reed canarygrass; 3900 lbs/ac and 1.07 AUM/ac, unfertilized, non-irrigated naturalized community. Refer to Forage Suitability Groups for cool-season pasture under a higher management level.

*Carrying capacity based on continuous season-long grazing by cattle under average growing conditions, 25% harvest efficiency. Air dry forage requirements based on 3% of animal body weight, or 912 lbs/AU/month.

If grazing distribution problems occur, stocking rates must be reduced to maintain plant health and vigor. Carrying capacity and production estimates are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Utilizing a rotational grazing system that allows for adequate rest and recovery will increase plant vigor and carrying capacity. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended to document plant composition and production. More precise carrying capacity estimates can be calculated based on actual site information along with animal preference data, particularly when livestock other than cattle are involved. With consultation of the land manager, more intensive grazing management may result in improved harvest efficiencies and increased carrying capacity.

Inventory data references

Information presented here has been derived from RANGE-417 archives, Rangeland NRI, and other inventory data. Field observations from range-trained personnel were also used. In addition to the multitude of NRCS field office employees and private landowners that helped with site visits and local knowledge, those involved in developing this site include:

Nebraska NRCS:

Nadine Bishop, State Rangeland Management Specialist
Patrick Cowsert, Resource Soil Scientist
Cassidy Gerdes, Biologist
Dirk Schultz, Soil Conservationist
Dan Shurtliff, Asst State Soil Scientist

South Dakota NRCS:

Stan Boltz, State Rangeland Management Specialist
Shane Deranleau, Area Rangeland Management Specialist
Kevin Luebke, State Biologist

Iowa NRCS:

Jess Jackson, Area Grazing Specialist

Minnesota NRCS:

Lance Smith, Area Grazing Specialist

MLRA Office 10:

Stu McFarland, Ecological Site Inventory Specialist, QC
Stacey Clark, Ecological Site Inventory Specialist, QA
Michael Whited, Soil Data Quality Specialist
Jo Parsley, Soil Scientist/10-3 MSSO Leader

National Soil Survey Center:

Mike Kucera, National Agronomist, Soil Quality & Ecosystems
Steve Peaslee, GIS Specialist, Soil Survey Interpretations

Nebraska Game & Parks Commission:

Gerry Steinauer, Botanist
Scott Wessel, Biologist
Russ Hamer, Biologist
Rebekah Jessen, Biologist

Nebraska Forest Service:

Steve Rasmussen, District Forester

Other references

Bestelmeyer, Brandon, et al. 2010. Practical Guidance for Developing State-and-transition Models. Rangelands 32:6 pp 2-64. Wheat Ridge, CO: Society for Range Management.

Fenneman, Nevin M. 1916. Physiographic Subdivision of the United States. Annals of the Association of American Geographers.

Helzer, Chris. 2010. The Ecology and Management of Prairies in the Central U.S. Iowa City, IA: University of Iowa Press/The Nature Conservancy.

Kaul, Robert B., David Sutherland, and Steven Rolfsmeier. 2006. The Flora of Nebraska. Lincoln, NE: University of Nebraska – Lincoln (Conservation and Survey Division, School of Natural Resources.)

NOAA/UNL – High Plains Regional Climate Center. Historical Data Summaries:
<http://www.hprcc.unl.edu/data/historical/>

Omerik, J.M. 1997. Ecoregions of the Conterminous United States. Annals of the Association of American Geographers, v.77, no. 1, p.118-125.

Steinauer, Gerry and Steve Rolfsmeier. 2010. Terrestrial Ecological Systems and Natural Communities of Nebraska. Lincoln, NE: Nebraska Natural Heritage Program and Nebraska Game and Parks Commission.

USDA/USFS. 2007. Ecological Subregions: Sections and Subsections for the Conterminous United States. Washington, DC: USDA - Forest Service.

USDA/SCS. 1977. Rangeland Resources of Nebraska. Lincoln, NE: Society for Range Management. USDA/NRCS. 2011. ESD User Guide. Fort Worth, TX: Central National Technology Support Center.

USDA/NRCS 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

USDA/NRCS. 2012. Field Office Technical Guide (Nebraska, Natural Resources Information, Statewide Soil and Site Information, Rangeland Interpretations, Nebraska Range Site Descriptions – Vegetative Zones 3 and 4), U.S. Department of Agriculture, Natural Resources Conservation Service, Nebraska Ecological Sciences.

Contributors

Approval

Suzanne Mayne-Kinney, 12/10/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Original Authors: Stu McFarland and Nadine Bishop. 08/01/2013 Revision Authors: Nadine Bishop, Emily Helms, Jeff Nichols
Contact for lead author	jeffrey.nichols@usda.gov
Date	12/04/2024
Approved by	Suzanne Mayne-Kinney
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None. Rills are not expected on this site.

2. **Presence of water flow patterns:** None. Water flow patterns are not expected on this site.

3. **Number and height of erosional pedestals or terracettes:** None. Pedestals and terracettes are not expected to occur on this site.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground is 5 percent or less.

Bare ground is exposed mineral soil that is not covered by vegetation (basal and/or foliar canopy), litter, standing dead vegetation, gravel/rock, and visible biological crust (e.g., lichen, mosses, algae).

5. **Number of gullies and erosion associated with gullies:** None. Gullies are not expected on this site.

6. **Extent of wind scoured, blowouts and/or depositional areas:** None. Wind-scoured and/or depositional areas should not be present.

7. **Amount of litter movement (describe size and distance expected to travel):** None. Litter falls into place. Litter movement is not expected on this site.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability ratings will be 5 to 6, typically 6. Surface organic matter adheres to the soil surface. Soil surface fragments will typically retain structure indefinitely when dipped in distilled water.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** A-horizon varies from 6 to 38 inches (15-97 cm) thick. Soil colors range from very dark gray, dark gray, gray to dark grayish brown (Hue: 10YR; value: 3 to 5; chroma: 1 or 2) when dry and black, very dark brown, very dark grayish brown, dark grayish brown, or very dark gray (Hue: 10YR; value 2 to 4; chroma: 1 or 2).

Soil structure of the A-horizon varies significantly with soil series and ranges from weak very fine granular to moderate fine granular, moderate very fine granular to medium subangular blocky.

See Official Soils Descriptions for additional details; major soil series correlated to the site are Ackmore, Boel, Coleridge, Els, Elsmere, Gibbon, Lamo, Ord, Primghar, Spillco, Splitrock, and Wann.

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Plant community composition is approximately 80 to 95 percent grasses or grass-like plants, 5 to 10 percent forbs, and 0 to 5 percent shrubs which optimizes infiltration on the site. The perennial grass and grass-like component is made up of warm-season (C4), tallgrasses; warm-season (C4), midgrasses; cool-season (C3) grasses; and grass-likes. The functional/structural groups provide a combination of rooting depths and structure which positively influences infiltration. \

Invasion of introduced cool-season grasses such as grasses such as reed canarygrass, creeping foxtail, smooth brome, or Kentucky bluegrass may negatively impact infiltration. Tree encroachment will also adversely impact infiltration.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. No compaction layers are expected on this site.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Phase 1.1

1. Native, perennial, warm-season, tallgrasses, 2295-3825 #/ac, 45-75% (4 species minimum): big bluestem, switchgrass, Indiangrass, prairie cordgrass.

2. Native, perennial, warm-season, midgrass, 765-1785 #/ac, 15-35%, (2 species minimum): Little bluestem, sideoats grama.

Phase 1.2

1. Native, perennial, warm-season, tallgrasses (3 species minimum): big bluestem, switchgrass, Indiangrass, prairie cordgrass.

Sub-dominant: Phase 1.1

1. Native, perennial, cool-season grass, 255-1020 #/ac, 5-20% (2 species minimum): western wheatgrass, reed canarygrass, porcupinegrass, foxtail barley, prairie Junegrass, prairie wedgescale, needle and thread, Scribner's rosettegrass.

Phase 1.2

1. Native, perennial, warm-season, midgrass (1 species minimum): little bluestem, sideoats grama.
2. Native, perennial, cool-season grass (2 species minimum): western wheatgrass, reed canarygrass, porcupinegrass, foxtail barley, prairie Junegrass, prairie wedgescale, needle and thread, Scribner's rosettegrass.

Other: Minor - Phase 1.1

1. Grass-likes, 255-510 #/ac, 5-10%: sedges, broom sedge, awlfruit sedge, rush, cloaked bulrush, common threesquare.
2. Native forbs, 255-510 #/ac, 5-10%: forbs present vary from location to location.
3. Shrubs, 0-255 #/ac, 0-5%: shrubs present will vary from location to location.

Minor - Phase 1.2

1. Grass-likes: sedges, broom sedge, awlfruit sedge, rush, cloaked bulrush, common threesquare.
2. Native forbs: forbs present vary from location to location.
3. Shrubs: shrubs present will vary from location to location.

Additional: The Big Bluestem-Little Bluestem Community or Reference Community (1.1) includes six F/S groups which include in order of relative abundance, native, perennial, warm-season (C 4) tallgrass; native, perennial, warm-season (C4) midgrass; native, perennial, cool-season (C3) grass; grass-likes; native forbs; shrubs.

The Prairie Cordgrass – Big Bluestem Community (1.2) includes six F/S groups which include in order of relative abundance, native, perennial, warm-season (C4) tallgrass; native, perennial, warm-season (C4) midgrass; native, perennial, cool-season (C3) grass; grass-likes; native, perennial, warm-season (C4) shortgrass; native forbs; and shrubs.

-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Bunchgrasses have strong, healthy centers with few (less than 3 percent) dead centers. Shrubs may show some dead branches (less than 5 percent) as plants age.
-
14. **Average percent litter cover (%) and depth (in):** Plant litter cover is evenly distributed throughout the site and is expected to be 80 to 90 percent and at a depth of approximately 0.5 to 1.0 inch (1.3 to 2.5 cm).
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** The representative value (RV) for annual production is 5,100 pounds per acre in a year with normal precipitation and temperatures. Low and High production years should yield 4,300 and 5,600 pounds per acre respectively.
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** No non-native invasive species are present. Kentucky bluegrass, smooth brome, reed canarygrass, leafy spurge, Canada thistle, eastern red cedar, roughleaf dogwood, buckbrush, and Siberian elm are

known invasives that have the potential to become dominant or co-dominant on the site. Note: species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants.

17. **Perennial plant reproductive capability:** All perennial species exhibit high vigor relative to climatic conditions. Perennial grasses should have vigorous rhizomes or tillers; vegetative and reproductive structures are not stunted. All perennial species should be capable of reproducing annually.
-