

Ecological site R102DY020SD

Loamy Overflow

Last updated: 8/14/2024
Accessed: 05/11/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 102D–Prairie Coteau

This area makes up about 7,867 square miles (20,375 square kilometers), consisting mostly of nearly level to undulating till plains with potholes and moraines. Elevation ranges from 1,150 to 2,130 feet (350 to 650 meters). The average annual precipitation is 22 to 29 inches (559 to 734 millimeters). The average annual temperature is 42 to 45 degrees F (6 to 7 degrees C). The dominant soil order in this MLRA is Mollisols. The soils in this area dominantly have a frigid temperature regime, and an aquic or udic moisture regime. They are generally very deep and loamy. Soils range from well drained to very poorly drained. Parent materials are dominantly fine-loamy till to clayey material, with smaller amounts of outwash, glaciofluvial deposits, eolian deposits, alluvium, and, to a lesser extent, loess and organic materials.

Classification relationships

Fenneman (1916) Physiographic Regions

Division - Interior Plains

East:

Province - Central Lowland

Section - Western Lake / Dissected Till Plains (12b/12e)

USFS (2007) Ecoregions

Domain - Humid Temperate

Division - Prairie

Province - Prairie Parkland (Temperate)

Section - North-Central Glaciated Plains (251B)

EPA Ecoregions (Omernik 1997)

I - Great Plains (9)

II - Temperate Prairies (9.2)

III - Aspen Parkland/Northern Glaciated Plains (9.2.1)

Ecological site concept

The Loamy Overflow ecological site occurs in upland swales which receive additional run off moisture from adjoining slopes. Soils are moderately well drained which have water flow into and over/through the site and have less than 40 percent clay in the surface and subsoil.

Vegetation in the Reference State includes big bluestem, Indiangrass, and Switchgrass. Forbs include goldenrods, cudweed sagewort, heath aster, western yarrow. Non-native grasses such as Kentucky bluegrass smooth brome grass may invade the site due to changes in disturbance regime.

Associated sites

R102DY010SD	Loamy These sites occur on upland areas. The soils are well drained and have less than 40 percent clay in the surface and subsoil.
R102DY012SD	Thin Upland These sites occur on uplands. Soils are well drained and will effervesce with acid at or near the surface.

Similar sites

R102DY010SD	Loamy The Loamy site occurs in a backslope landscape position. Soils are well drained and do not have water flow into and over/through the site. The Loamy site will have less big bluestem, more needlegrass and lower production than the Loamy Overflow site.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Andropogon gerardii</i> (2) <i>Sorghastrum nutans</i>

Physiographic features

This site occurs on upland swales which receive additional run-on moisture from adjoining slopes.

Table 2. Representative physiographic features

Landforms	(1) Swale
Runoff class	Low to medium
Flooding frequency	None
Ponding frequency	None
Elevation	1,000–2,000 ft
Slope	0–2%
Water table depth	30–60 in
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

The average annual precipitation is 22 to 28 inches. Half or more of the precipitation falls during the growing season. Rainfall typically occurs during high-intensity, convective thunderstorms in summer. In the western part of the MLRA, rainfall is less abundant and not always adequate for full maturation of crops. Precipitation in winter is typically snow. The average annual temperature is 42 to 45 degrees F. The freeze-free period averages 143 days and ranges from 131 to 151 days.

Table 3. Representative climatic features

Frost-free period (characteristic range)	114-129 days
Freeze-free period (characteristic range)	138-149 days
Precipitation total (characteristic range)	24-27 in
Frost-free period (actual range)	110-131 days
Freeze-free period (actual range)	131-151 days
Precipitation total (actual range)	22-28 in

Frost-free period (average)	122 days
Freeze-free period (average)	142 days
Precipitation total (average)	25 in




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) ROY LAKE [USC00397326], Lake City, SD
- (2) WEBSTER [USC00399004], Webster, SD
- (3) WAUBAY NWR [USC00398980], Waubay, SD
- (4) WATERTOWN RGNL AP [USW00014946], Watertown, SD
- (5) WATERTOWN 1W [USC00398930], Watertown, SD
- (6) CASTLEWOOD [USC00391519], Castlewood, SD
- (7) CLEAR LAKE [USC00391777], Clear Lake, SD
- (8) ARLINGTON 1 W [USC00390281], Arlington, SD
- (9) BROOKINGS 2 NE [USC00391076], Brookings, SD
- (10) TYLER [USC00218429], Tyler, MN

Influencing water features

No riparian areas or wetland features are directly associated with this site.

Soil features

Soils are typically formed in loess or till. Surface textures are silty clay loam or loam. Soils are moderately well-drained.

Table 4. Representative soil features

Parent material	(1) Loess (2) Till
Surface texture	(1) Silty clay loam (2) Loam
Drainage class	Moderately well drained
Permeability class	Moderately rapid
Soil depth	80 in
Surface fragment cover ≤3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	7.3–8.2 in
Soil reaction (1:1 water) (0-10in)	5.6–7.8
Subsurface fragment volume ≤3" (0-60in)	0–2%
Subsurface fragment volume >3" (0-60in)	0–1%

Ecological dynamics


The site which is located in the Prairie Pothole Region developed under Northern Great Plains climatic conditions and included natural influence of large herding herbivores and occasional fire. Changes will occur in the plant communities due to weather fluctuations and/or management actions. Under adverse impacts, a relatively rapid decline in vegetative vigor and composition can occur. Under favorable conditions the site has the potential to resemble the Reference State. Interpretations for this site are based primarily on the 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase. This community phase and the Reference State have been determined by study of rangeland relic areas, areas protected from excessive disturbance, and areas under long-term rotational grazing regimes. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts also have been considered.

This ecological site (ES) has been grazed by domestic livestock since they have been introduced into the area. The introduction of domestic livestock and the use of fencing and reliable water sources have changed the ecological dynamics of this site. Continuous grazing without adequate recovery periods following each grazing occurrence over several years causes this site to depart from the interpretive plant community. Species such as little bluestem (*Schizachyrium scoparium*), sideoats grama (*Bouteloua curtipendula*), and sedge (*Carex*) will initially increase. Big bluestem, Indiangrass, and switchgrass will decrease in frequency and production. Heavy continuous grazing causes Kentucky bluegrass (*Poa pratensis*) to increase and eventually develop into a sod condition. Extended periods of nonuse and no fire will result in a plant community having high litter levels which favors an increase in Kentucky bluegrass and smooth brome (*Bromus inermis*). In time, shrubs such as western snowberry (*Symphoricarpos occidentalis*) and chokecherry (*Prunus virginiana*) will also increase.


Following the state and transition diagram are narratives for each of the described states and community phases. These may not represent every possibility, but they are the most prevalent and repeatable states/community phases. The plant composition tables shown below have been developed from the best available knowledge at the time of this revision. As more data are collected, some of these community phases and/or states may be revised or removed, and new ones may be added. The main purpose for including the descriptions here is to capture the current knowledge and experience at the time of this revision.

State and transition model

Ecosystem states



States 1, 5 and 2 (additional transitions)



T1A - Heavy continuous grazing

T1B - Heavy continuous grazing, non-use, no fire, invasion

T1C - Tillage

T2A - Non-native encroachment/seeding

T2B - Tillage

T3A - Non-use, no fire, heavy continuous grazing

T3C - Tillage


R4A - Long-term prescribed grazing, prescribed burning

T4A - Tillage

R5A - Seeding

R5B - Seeding, abandonment of cropping

State 1 submodel, plant communities




1.1A - Heavy continuous grazing

1.2A - Prescribed grazing with recovery periods, prescribed burning

State 2 submodel, plant communities




State 3 submodel, plant communities



3.1A - Heavy continuous grazing

3.2A - Prescribed grazing with recovery periods

State 4 submodel, plant communities




4.1A - Heavy continuous grazing

4.2A - Prescribed grazing with recovery periods

4.3A - Time without disturbances

State 5 submodel, plant communities



State 1 Reference State

The Reference State represents the natural range of variability that dominated the dynamics of this ecological site (ES). This state was dominated by warm-season grasses. In pre-European times, the primary disturbance mechanisms for this site in the reference condition included periods of below and/or above average precipitation, periodic fire, and herbivory by insects and large ungulates. Timing of fires and herbivory coupled with weather events dictated the dynamics that occurred within the natural range of variability. In some locations, this site likely received relatively heavy grazing pressure. Tall warm-season grasses would have declined and shorter warm-season grasses would have increased. Today, a similar state, the Native/Invaded State (State 3) can be found on areas that are properly managed with grazing and/or prescribed burning and sometimes on areas receiving occasional short periods of rest.

Dominant plant species

- snowberry (*Symphoricarpos*), shrub
- leadplant (*Amorpha canescens*), shrub

- big bluestem (*Andropogon gerardii*), grass
- Indiangrass (*Sorghastrum nutans*), grass
- little bluestem (*Schizachyrium scoparium*), grass
- green needlegrass (*Nassella viridula*), grass
- slender wheatgrass (*Elymus trachycaulus*), grass
- American licorice (*Glycyrrhiza lepidota*), other herbaceous
- Maximilian sunflower (*Helianthus maximiliani*), other herbaceous
- blazing star (*Liatris*), other herbaceous

Community 1.1

Big Bluestem-Indiangrass-Switchgrass

Interpretations are based primarily on the 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase (this is also considered to be climax). The potential vegetation was about 80 percent grasses or grass-like plants, 10 percent forbs, and 10 percent shrubs. The community was dominated by warm-season grasses. The major grasses included big bluestem, Indiangrass, and switchgrass. Other grass or grass-like species included porcupine grass (*Hesperostipa spartea*), green needlegrass (*Nassella viridula*), slender wheatgrass (*Elymus trachycaulus*), and little bluestem. This plant community was resilient and well adapted to the Northern Great Plains climatic conditions. The diversity in plant species allowed for high drought tolerance. This was a sustainable plant community in regards to site/soil stability, watershed function, and biologic integrity.

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	3400	3910	4300
Forb	200	345	550
Shrub/Vine	200	345	550
Total	3800	4600	5400




Figure 8. Plant community growth curve (percent production by month). SD0205, Rolling Till Prairie, warm-season dominant.. Warm-season dominant..

Community 1.2

Little Bluestem-Sideoats Grama-Big Bluestem-Forbs

This plant community evolved under heavy continuous grazing or from over utilization during extended drought periods. The potential plant community was made up of approximately 75 percent grasses and grass-like species, 15 percent forbs, and 10 percent shrubs. Dominant grasses included little bluestem, sideoats grama, and big bluestem. Grasses of secondary importance included Indiangrass, switchgrass, green needlegrass, tall dropseed (*Sporobolus compositus*), and slender wheatgrass. Forbs commonly found in this plant community included Canada goldenrod (*Solidago Canadensis*), cudweed sagewort (*Artemisia ludoviciana*), heath aster (*Symphotrichum ericoides*), scurfpea (*Psoralegium*), stiff goldenrod (*Oligoneuron rigidum*), and western yarrow (*Achillea millefolium*). This plant community had similar plant composition to the 3.2 Little Bluestem-Kentucky Bluegrass-Big Bluestem Plant Community Phase. The main difference is that this plant community phase did not have the presence of nonnative invasive species such as Kentucky bluegrass and smooth brome grass. When

compared to the 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase, sideoats grama and little bluestem increased. Production of tall warm-season grasses was reduced. This plant community was moderately resistant to change. The herbaceous species present were well adapted to grazing; however, species composition could be altered through long-term overgrazing. If the herbaceous component was intact, it tended to be resilient if the disturbance was not long-term. Most of the components of the ecological processes would have been functioning at optimum levels. However, the vigor and reproductive capability of the tall warm-season grasses would have been reduced due to grazing pressure or a combination of stressors. A reduction of this dominant functional group allowed for an increase in shorter-statured (and shallower rooted) species.




Figure 9. Plant community growth curve (percent production by month).
SD0205, Rolling Till Prairie, warm-season dominant.. Warm-season dominant..

Pathway 1.1A

Community 1.1 to 1.2

Heavy continuous grazing which includes herbivory at moderate to heavy levels at the same time of year each year without adequate recovery periods, or during periods of below normal precipitation when grazing frequency and intensity increases on these sites due to limited forage availability on adjacent upland sites will shift this community to the 1.2 Little Bluestem-Sideoats Grama-Big Bluestem-Forbs Plant Community Phase.

Pathway 1.2A

Community 1.2 to 1.1

Prescribed grazing, and/or prescribed burning returned to normal disturbance regime levels and frequencies or periodic light to moderate grazing possibly including periodic rest will convert this plant community to the 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase.

State 2

Shortgrass Sod State

This state is the result of heavy continuous grazing, and in the absence of periodic fire due to fire suppression. This state is dominated by blue grama and buffalograss forming a dense sod layer that effectively blocks introduction of other plants into the system. Taller cool-season species will decline and a corresponding increase in short statured grass will occur. Once the threshold is crossed, a change in grazing management alone cannot cause a reduction in the sod grass dominance.

Dominant plant species

- snowberry (*Symphoricarpos*), shrub
- leadplant (*Amorpha canescens*), shrub
- rose (*Rosa*), shrub
- blue grama (*Bouteloua gracilis*), grass
- sideoats grama (*Bouteloua curtipendula*), grass
- little bluestem (*Schizachyrium scoparium*), grass
- threadleaf sedge (*Carex filifolia*), grass
- white sagebrush (*Artemisia ludoviciana*), other herbaceous
- goldenrod (*Solidago*), other herbaceous

- western yarrow (*Achillea millefolium* var. *occidentalis*), other herbaceous

Community 2.1

Sideoats Grama-Little Bluestem-Sedges-Forbs

This plant community evolved under heavy continuous season grazing or from over utilization during extended drought periods. The potential plant community was made up of approximately 80 percent grasses and grass-like species, 15 percent forbs, and 5 percent shrubs. Dominant grass and grass-like species included sideoats grama, little bluestem, and sedge. Grasses of secondary importance included big bluestem, switchgrass, green needlegrass, slender wheatgrass, and tall dropseed. Forbs commonly found in this plant community included cudweed sagewort, goldenrod (*Solidago*), and western yarrow. When compared to the 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase, tall warm-season grasses were reduced, and the more grazing tolerant species such as sideoats grama, little bluestem and sedge were dominant on this plant community. With the exception of green needlegrass, cool-season grasses decreased significantly. This vegetation state was very resistant to change, especially if the disturbance continued and the short-statured species such as sedge increased. The herbaceous species present were well adapted to grazing. This plant community was less productive than other phases.




Figure 10. Plant community growth curve (percent production by month).
SD0204, Rolling Till Prairie, warm-season dominant, cool-season subdominant.. Warm-season dominant, cool-season subdominant..

State 3

Native/Invaded State

This state represents the more common range of variability that exists with higher levels of grazing management but in the absence of periodic fire due to fire suppression. This state is dominated by warm-season grasses, with cool-season grasses being subdominant. It can be found on areas that are properly managed with grazing and/or prescribed burning, and sometimes on areas receiving occasional short periods of rest. Taller warm-season species can decline and a corresponding increase in short statured grass will occur.

Dominant plant species

- snowberry (*Symphoricarpos*), shrub
- rose (*Rosa*), shrub
- big bluestem (*Andropogon gerardii*), grass
- little bluestem (*Schizachyrium scoparium*), grass
- Kentucky bluegrass (*Poa pratensis*), grass
- smooth brome (*Bromus inermis*), grass
- green needlegrass (*Nassella viridula*), grass
- goldenrod (*Solidago*), other herbaceous
- white heath aster (*Symphyotrichum ericoides*), other herbaceous
- white sagebrush (*Artemisia ludoviciana*), other herbaceous

Community 3.1

Big Bluestem-Indiangrass-Switchgrass

This plant community phase is similar to the 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase but

it also contains minor amounts of non-native invasive grass species such as Kentucky bluegrass and smooth brome grass (up to about 10 percent by air-dry weight). The potential vegetation is about 80 percent grasses or grass-like plants, 10 percent forbs, and 10 percent shrubs. This community is dominated by warm-season grasses. The major grasses include big bluestem, Indiangrass, and switchgrass. Other grass or grass-like species include porcupine grass, green needlegrass, slender wheatgrass, and little bluestem. This plant community is resilient and well adapted to the Northern Great Plains climatic conditions. The diversity in plant species allows for high drought tolerance. This is a sustainable plant community in regards to site/soil stability, watershed function, and biologic integrity.




Figure 11. Plant community growth curve (percent production by month). SD0205, Rolling Till Prairie, warm-season dominant.. Warm-season dominant..

Community 3.2 Little Bluestem-Kentucky Bluegrass-Big Bluestem

This plant community is a result of heavy continuous grazing or from over utilization during extended drought periods. The potential plant community is made up of approximately 75 percent grasses and grass-like species, 15 percent forbs, and 10 percent shrubs. Dominant grasses include little bluestem, big bluestem, and Kentucky bluegrass. Grasses of secondary importance include Indiangrass, switchgrass, smooth brome grass, sideoats grama, green needlegrass, tall dropseed, slender wheatgrass, and sedge. Forbs commonly found in this plant community include cudweed sagewort, heath aster, Canada goldenrod, stiff goldenrod, scurfpea, Indian hemp (*Apocynum cannabinum*), and western yarrow. When compared to the 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase, little bluestem has increased and Kentucky bluegrass has invaded and become a codominant. Production of tall warm-season grasses is reduced. This plant community is moderately resistant to change. The herbaceous species present are well adapted to grazing; however, species composition can be altered through long-term overgrazing. If the herbaceous component is intact, it tends to be resilient if the disturbance is not long-term. Most of the components of the ecological processes are functioning at optimum levels. However, the vigor and reproductive capability of the tall warm-season grasses are reduced due to grazing pressure or a combination of stressors. A reduction of this dominant functional group allows for an increase in shorter-statured (and shallower rooted) species. The introduction of nonnative invasive species such as Kentucky bluegrass and smooth brome grass results in alterations to the soil profile. Organic matter levels tend to decrease and begin to be concentrated more in the surface layers and the structure will begin to be modified. These changes favor the shallow-rooted species and hasten their eventual dominance if steps are not taken to reduce these species.

Table 6. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	2285	2856	3290
Forb	150	340	610
Shrub/Vine	65	204	400
Total	2500	3400	4300




Figure 13. Plant community growth curve (percent production by month).
SD0204, Rolling Till Prairie, warm-season dominant, cool-season subdominant.. Warm-season dominant, cool-season subdominant..

Pathway 3.1A Community 3.1 to 3.2

Heavy continuous grazing which includes herbivory at moderate to heavy levels at the same time of year each year without adequate recovery periods, or during periods of below normal precipitation when grazing frequency and intensity increases on these sites due to limited forage availability on adjacent upland sites will shift this community to the 3.2 Little Bluestem-Kentucky Bluegrass-Big Bluestem Plant Community Phase.

Pathway 3.2A Community 3.2 to 3.1

Prescribed grazing (alternating season of use and providing adequate recovery periods) or periodic light to moderate grazing possibly including periodic rest will convert this plant community to the 3.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase.

Conservation practices

Prescribed Grazing

State 4 Invaded State

This state is a result of encroachment mainly by invasive introduced cool-season grasses. The ecological processes are not functioning, especially the biotic processes and the hydrologic functions. The introduced cool-season grasses cause reduced infiltration and increased runoff. Preliminary studies would tend to indicate this threshold may exist when Kentucky bluegrass exceeds 30 percent of the plant community and native grasses represent less than 40 percent of the plant community composition. The opportunity for high intensity spring burns is severely reduced by early greenup and increased moisture and humidity at the soil surface and grazing pressure cannot cause a reduction in sodgrass dominance. Production is limited to the sod forming species. Infiltration continues to decrease and runoff increases and energy capture into the system is restricted to early season low producing species. Nutrient cycling is limited by root depth of the dominant species.

Dominant plant species

- snowberry (*Symphoricarpos*), shrub
- rose (*Rosa*), shrub
- smooth brome (*Bromus inermis*), grass
- Kentucky bluegrass (*Poa pratensis*), grass
- big bluestem (*Andropogon gerardii*), grass
- switchgrass (*Panicum virgatum*), grass
- threadleaf sedge (*Carex filifolia*), grass
- goldenrod (*Solidago*), other herbaceous
- white heath aster (*Symphyotrichum ericoides*), other herbaceous
- white clover (*Trifolium repens*), other herbaceous

- white sagebrush (*Artemisia ludoviciana*), other herbaceous

Community 4.1

Smooth Brome-grass-Kentucky Bluegrass

This plant community phase is a result of extended periods of nonuse and no fire or occasionally light levels of grazing over several years. It is characterized by dominance of smooth brome-grass and to a lesser extent Kentucky bluegrass. The dominance is at times so complete that other species are difficult to find on the site. A thick duff layer also accumulates at or above the soil surface and eventually a thatch-mat layer may develop at the surface. Nutrient cycling is greatly reduced and native plants have great difficulty becoming established. When dominated by smooth brome-grass, infiltration is moderately reduced and runoff is moderate. Production can be equal to or higher than the interpretive plant community. However, when dominated by Kentucky bluegrass, infiltration is greatly reduced and runoff is high. Production in this case will likely be significantly less. In either case, the period that palatability is high is relatively short as these cool-season species mature rapidly. Energy capture is also reduced. The dominance of these introduced species has been shown to alter the biotic component of the soil, as well as, organic matter levels and eventually the soil structure. These alterations perpetuate the dominance of Kentucky bluegrass and smooth brome-grass and tend to make establishment of native species extremely difficult.

Table 7. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	2750	3460	4050
Forb	175	300	475
Shrub/Vine	75	240	475
Total	3000	4000	5000




Figure 15. Plant community growth curve (percent production by month). SD0201, Rolling Till Prairie, cool-season dominant.. Cool-season dominant..

Community 4.2

Kentucky Bluegrass-Quackgrass

This plant community phase is a result of heavy, continuous seasonal grazing or heavy, continuous season-long grazing. It is characterized by a dominance of Kentucky bluegrass and quackgrass. The dominance is at times so complete that other species are difficult to find on the site. A relatively thick duff layer can sometimes accumulate at or above the soil surface and a thatch-mat layer often develops at the surface. Nutrient cycling is greatly reduced and native plants have great difficulty becoming established. Infiltration is greatly reduced and runoff is high. Production will be significantly reduced when compared to the interpretive plant community. The period that palatability is high is relatively short as Kentucky bluegrass matures rapidly. Energy capture is also reduced. Biological activity in the soil is likely reduced significantly in this phase.

Table 8. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	1660	2249	2765
Forb	115	260	460
Shrub/Vine	25	91	175
Total	1800	2600	3400




Figure 17. Plant community growth curve (percent production by month).
SD0201, Rolling Till Prairie, cool-season dominant.. Cool-season dominant..

Community 4.3

Annual Pioneer-Perennial Pioneer

This plant community developed under continuous heavy grazing or other excessive disturbances. The potential plant community is made up of approximately 40 to 80 percent grasses and grass-like species, 20 to 60 percent forbs, and 0 to 5 percent shrubs. The species present in this phase are highly variable but often include non-native invasive and/or early seral species. Plant diversity is low (plant richness may be high, but areas are often dominated by a few species). The ecological processes are difficult to restore because of the loss of plant diversity and overall soil disturbance. Soil erosion is potentially very high because of the bare ground and shallow rooted herbaceous plant community. Water runoff will increase and infiltration will decrease due to animal related soil compaction and loss of root mass due to low plant diversity and vigor. This plant community will require significant economic inputs and time to move towards another plant community. This movement is highly variable in its succession. This is due to the loss of diversity (including the loss of the seed bank), within the existing plant community, and the plant communities on adjacent sites. This community can be renovated to improve the production capability; however, if management changes are not made the vegetation could revert back to early seral species.

Pathway 4.1A

Community 4.1 to 4.2

Heavy continuous grazing which includes herbivory at moderate to heavy levels at the same time of year each year without adequate recovery periods, or during periods of below normal precipitation when grazing frequency and intensity increases on these sites due to limited forage availability on adjacent upland sites will shift this community to the 4.2 Kentucky Bluegrass-Quackgrass Plant Community Phase.

Pathway 4.2A

Community 4.2 to 4.1

Prescribed grazing (alternating season of use and providing adequate recovery periods) or periodic light to moderate grazing possibly including periodic rest will convert this plant community to the 4.1 Smooth Bromegrass-Kentucky Bluegrass Plant Community Phase.

Conservation practices

Prescribed Grazing

Pathway 4.3A

Community 4.3 to 4.2

This community pathway occurs with the passage of time as successional processes take place and perennial plants gradually begin to establish on the site again. This pathway will lead to the 4.2 Kentucky Bluegrass-Quackgrass Plant Community Phase.

State 5

Crop Production State

This state is characterized by the production of annual crops using a variety of tillage and cropping systems along with management practices. Cropping on this site is enabled during years with drier than normal precipitation or with artificial drainage (surface or subsurface).

Dominant plant species

- corn (*Zea*), grass
- wheat (*Triticum*), grass
- soybean (*Glycine*), other herbaceous

Community 5.1

Annual Crops

This plant community developed with the use of a variety of tillage systems and cropping systems for the production of annual crops including corn, soybeans, wheat, sugar beet and a variety of other crops.

Transition T1A

State 1 to 2

Heavy continuous grazing (stocking levels well above carrying capacity for extended portions of the growing season, and often at the same time of year each year, typically beginning early in the season) will convert this plant community to the 2.1 Sideoats Grama-Little Bluestem-Sedges-Forbes Plant Community Phase within the Shortgrass Sod State.

Transition T1B

State 1 to 3

Non-use and/or no surface fire for extended periods of time (typically for 10 or more years) causing litter levels to become high enough to reduce native grass vigor, diversity, and density, and/or heavy continuous grazing or invasion of non-native plant species will likely lead this state over a threshold resulting in the Native/Invaded State (State 3).

Transition T1C

State 1 to 5

Tillage will cause a shift over a threshold leading to the 5.1 Annual Crops within the Crop Production State (State 5).

Transition T2A

State 2 to 4

Encroachment of non-native invasive/noxious species, abandonment of cropping, or seeding of introduced and/or native improved varieties of forage species may lead this plant community phase over a threshold to the Invaded State (State 4) and more specifically to the 4.3 Annual/Pioneer, Non-native Perennial Plant Community Phase. In the case of a seeding, refer to the corresponding Forage Suitability Group (FSG) description for adapted species and expected production (production estimates in the FSG description may be unrealistically high due to the degraded condition of the site at this phase).

Transition T2B
State 2 to 5

Tillage will cause a shift over a threshold leading to the 5.1 Annual Crops within the Crop Production State (State 5).

Transition T3A
State 3 to 4

Non-use and/or no surface fire for extended periods of time (typically for 10 or more years) causing litter levels to become high enough to reduce native grass vigor, diversity, and density, will likely lead this state over a threshold leading to the 4.1 Smooth Bromegrass-Kentucky Bluegrass Community Phase within the Invaded State (State 4). Heavy continuous grazing (stocking levels well above carrying capacity for extended portions of the growing season and often at the same time of year each year), will likely lead this state over a threshold leading to the 4.2 Kentucky Bluegrass-Quackgrass Community Phase within the Invaded State (State 4). Grazing repeatedly in the early growing season can expedite this shift by causing mechanical disturbance due to trampling.

Transition T3C
State 3 to 5

Tillage will cause a shift over a threshold leading to the 5.1 Annual Crops within the Crop Production State (State 5).

Restoration pathway R4A
State 4 to 3

Long-term prescribed grazing (moderate stocking levels coupled with adequate recovery periods, or other grazing systems such as high-density, low-frequency intended to treat specific species dominance, or periodic light to moderate stocking levels possibly including periodic rest) coupled with prescribed burning may lead this plant community phase over a threshold to the Native/Invaded State (State 3).

Conservation practices

Prescribed Grazing
Integrated Pest Management (IPM)

Transition T4A
State 4 to 5

Tillage will cause a shift over a threshold leading to the 5.1 Annual Crops Community Phase within the Crop Production State (State 5).

Restoration pathway R5A
State 5 to 3

Seeding may lead this Crop Production State (State 5) over a threshold to the Native/Invaded State (State 3).

Restoration pathway R5B
State 5 to 4

Seeding may lead this Crop Production State (State 5) over a threshold to the Invaded State (State 4). Cropping followed by abandonment may lead this plant community phase over a threshold to the Invaded State (State 4) and more specifically to the 4.3 Annual Pioneer-Perennial Pioneer Plant Community Phase.

Additional community tables

Table 9. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
-------	-------------	--------	-----------------	--------------------------------	---------------------

Grass/Grasslike					
1	Tall Warm-season Grasses			690–2300	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	690–2300	–
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	460–1380	–
	switchgrass	PAVI2	<i>Panicum virgatum</i>	230–690	–
	spiked muhly	MUGL3	<i>Muhlenbergia glomerata</i>	0–230	–
	composite dropseed	SPCOC2	<i>Sporobolus compositus</i> var. <i>compositus</i>	0–138	–
2	Mid Warm-season Grasses			230–460	
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	92–460	–
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	46–230	–
	prairie dropseed	SPHE	<i>Sporobolus heterolepis</i>	0–138	–
3	Cool-season Bunchgrasses			230–460	
	green needlegrass	NAVI4	<i>Nassella viridula</i>	92–460	–
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	46–230	–
	porcupinegrass	HESP11	<i>Hesperostipa spartea</i>	0–138	–
4	Wheatgrass			92–460	
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	92–460	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	0–230	–
5	Other Native Grasses			92–230	
	Graminoid (grass or grass-like)	2GRAM	<i>Graminoid (grass or grass-like)</i>	46–230	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	46–92	–
	Scribner's rosette grass	DIOLS	<i>Dichanthelium oligosanthos</i> var. <i>scribnerianum</i>	0–46	–
6	Grass-likes			46–230	
	sedge	CAREX	<i>Carex</i>	46–230	–
	Grass-like (not a true grass)	2GL	<i>Grass-like (not a true grass)</i>	0–92	–
Forb					
7	Forbs			230–460	
	Forb, native	2FN	<i>Forb, native</i>	46–138	–
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	0–138	–
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	46–138	–
	blazing star	LIATR	<i>Liatris</i>	46–92	–
	western yarrow	ACMIO	<i>Achillea millefolium</i> var. <i>occidentalis</i>	46–92	–
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	46–92	–
	Indianhemp	APCA	<i>Apocynum cannabinum</i>	0–92	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	46–92	–
	wavyleaf thistle	CIUN	<i>Cirsium undulatum</i>	46–92	–
	goldenrod	SOLID	<i>Solidago</i>	46–92	–
	white heath aster	SYER	<i>Symphyotrichum ericoides</i>	46–92	–
	American vetch	VIAM	<i>Vicia americana</i>	46–92	–
	cinquefoil	POTEN	<i>Potentilla</i>	46–92	–
	scurfpea	PSORA2	<i>Psoraleidum</i>	46–92	–

	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	46–92	–
	ragwort	SENEC	<i>Senecio</i>	0–46	–
	golden tickseed	COTI3	<i>Coreopsis tinctoria</i>	0–46	–
	Illinois bundleflower	DEIL	<i>Desmanthus illinoensis</i>	0–46	–
	northern bedstraw	GABO2	<i>Galium boreale</i>	0–46	–
	Canadian anemone	ANCA8	<i>Anemone canadensis</i>	0–46	–
	wood lily	LIPH	<i>Lilium philadelphicum</i>	0–46	–
Shrub/Vine					
8	Shrubs			230–460	
	snowberry	SYMPH	<i>Symphoricarpos</i>	46–184	–
	leadplant	AMCA6	<i>Amorpha canescens</i>	46–184	–
	false indigo bush	AMFR	<i>Amorpha fruticosa</i>	0–92	–
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0–92	–
	American plum	PRAM	<i>Prunus americana</i>	0–92	–
	chokecherry	PRVI	<i>Prunus virginiana</i>	0–92	–
	currant	RIBES	<i>Ribes</i>	0–92	–
	rose	ROSA5	<i>Rosa</i>	46–92	–

Table 10. Community 3.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass/Grasslike					
1	Tall Warm-season Grasses			170–850	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	170–680	–
	switchgrass	PAVI2	<i>Panicum virgatum</i>	170–340	–
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	0–340	–
	composite dropseed	SPCOC2	<i>Sporobolus compositus</i> var. <i>compositus</i>	0–170	–
2	Mid Warm-season Grasses			340–850	
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	340–850	–
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	68–272	–
3	Cool-season Bunchgrasses			34–340	
	green needlegrass	NAVI4	<i>Nassella viridula</i>	34–340	–
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	0–68	–
	porcupinegrass	HESP11	<i>Hesperostipa spartea</i>	0–68	–
4	Wheatgrass			0–170	
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	0–170	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	0–102	–
5	Other Native Grasses			34–170	
	Graminoid (grass or grass-like)	2GRAM	<i>Graminoid (grass or grass-like)</i>	0–170	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0–68	–
	Scribner's rosette grass	DIOLS	<i>Dichanthelium oligosanthes</i> var. <i>scribnerianum</i>	0–34	–
6	Grass-likes			34–170	
	codon	CABEV	<i>Cory</i>	34–170	

	Seuge	CAREX	Carex	34-170	-
	Grass-like (not a true grass)	2GL	Grass-like (not a true grass)	0-68	-
7	Non-Native Grasses			340-850	
	Kentucky bluegrass	POPR	<i>Poa pratensis</i>	170-680	-
	smooth brome	BRIN2	<i>Bromus inermis</i>	68-340	-
	quackgrass	ELRE4	<i>Elymus repens</i>	0-170	-
Forb					
8	Forbs			170-510	
	goldenrod	SOLID	<i>Solidago</i>	34-170	-
	white heath aster	SYER	<i>Symphyotrichum ericoides</i>	34-136	-
	Forb, introduced	2FI	<i>Forb, introduced</i>	34-136	-
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	34-136	-
	Forb, native	2FN	<i>Forb, native</i>	0-102	-
	western yarrow	ACMIO	<i>Achillea millefolium var. occidentalis</i>	34-102	-
	Indianhemp	APCA	<i>Apocynum cannabinum</i>	0-102	-
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	0-102	-
	scurfpea	PSORA2	<i>Psoraleidum</i>	34-102	-
	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	0-68	-
	blazing star	LIATR	<i>Liatris</i>	0-68	-
	cinquefoil	POTEN	<i>Potentilla</i>	0-68	-
	common dandelion	TAOF	<i>Taraxacum officinale</i>	0-68	-
	white clover	TRRE3	<i>Trifolium repens</i>	0-68	-
	Canada thistle	CIAR4	<i>Cirsium arvense</i>	0-68	-
	wavyleaf thistle	CIUN	<i>Cirsium undulatum</i>	0-34	-
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	0-34	-
	American vetch	VIAM	<i>Vicia americana</i>	0-34	-
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	0-34	-
	ragwort	SENEC	<i>Senecio</i>	0-34	-
Shrub/Vine					
9	Shrubs			68-340	
	snowberry	SYMPH	<i>Symphoricarpos</i>	34-272	-
	rose	ROSA5	<i>Rosa</i>	34-102	-
	false indigo bush	AMFR	<i>Amorpha fruticosa</i>	0-102	-
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0-68	-
	leadplant	AMCA6	<i>Amorpha canescens</i>	0-34	-
	American plum	PRAM	<i>Prunus americana</i>	0-34	-
	chokecherry	PRVI	<i>Prunus virginiana</i>	0-34	-

Table 11. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass/Grasslike					
1	Tall Warm-season Grasses			0-400	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	0-320	-

	switchgrass	PAVI2	<i>Panicum virgatum</i>	0–320	–
	composite dropseed	SPCOC2	<i>Sporobolus compositus</i> var. <i>compositus</i>	0–170	–
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	0–80	–
2	Mid Warm-season Grasses			0–200	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	0–200	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	0–200	–
3	Cool-season Bunchgrasses			0–320	
	green needlegrass	NAVI4	<i>Nassella viridula</i>	0–320	–
4	Wheatgrass			0–200	
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	0–200	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	0–80	–
5	Other Native Grasses			0–200	
	Graminoid (grass or grass-like)	2GRAM	Graminoid (grass or grass-like)	0–160	–
	Scribner's rosette grass	DIOLS	<i>Dichanthelium oligosanthes</i> var. <i>scribnerianum</i>	0–40	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0–40	–
6	Grass-likes			0–200	
	sedge	CAREX	<i>Carex</i>	0–200	–
	Grass-like (not a true grass)	2GL	Grass-like (not a true grass)	0–40	–
7	Non-Native Grasses			1200–2800	
	smooth brome	BRIN2	<i>Bromus inermis</i>	800–2400	–
	Kentucky bluegrass	POPR	<i>Poa pratensis</i>	400–1400	–
	quackgrass	ELRE4	<i>Elymus repens</i>	0–320	–
Forb					
8	Forbs			200–400	
	goldenrod	SOLID	<i>Solidago</i>	40–120	–
	white heath aster	SYER	<i>Symphyotrichum ericoides</i>	40–120	–
	white clover	TRRE3	<i>Trifolium repens</i>	0–120	–
	Forb, introduced	2FI	Forb, introduced	40–120	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	40–120	–
	Canada thistle	CIAR4	<i>Cirsium arvense</i>	0–120	–
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	0–80	–
	Forb, native	2FN	Forb, native	0–80	–
	western yarrow	ACMIO	<i>Achillea millefolium</i> var. <i>occidentalis</i>	40–80	–
	Indianhemp	APCA	<i>Apocynum cannabinum</i>	0–80	–
	scurfpea	PSORA2	<i>Psoraleidium</i>	40–80	–
	common dandelion	TAOF	<i>Taraxacum officinale</i>	0–80	–
	cinquefoil	POTEN	<i>Potentilla</i>	0–40	–
Shrub/Vine					
9	Shrubs			80–400	
	snowberry	SYMPH	<i>Symphoricarpos</i>	40–320	–
	rose	ROSA5	<i>Rosa</i>	40–120	–

	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0–80	–
	false indigo bush	AMFR	<i>Amorpha fruticosa</i>	0–80	–
	American plum	PRAM	<i>Prunus americana</i>	0–80	–

Table 12. Community 4.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass/Grasslike					
1	Tall Warm-season Grasses			0–130	
	switchgrass	PAVI2	<i>Panicum virgatum</i>	0–78	–
	composite dropseed	SPCOC2	<i>Sporobolus compositus</i> var. <i>compositus</i>	0–78	–
2	Mid Warm-season Grasses			0–78	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	0–78	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	0–78	–
3	Cool-season Bunchgrasses			0–130	
	green needlegrass	NAVI4	<i>Nassella viridula</i>	0–130	–
4	Wheatgrass			0–130	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	0–104	–
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	0–52	–
5	Other Native Grasses			0–78	
	Graminoid (grass or grass-like)	2GRAM	<i>Graminoid (grass or grass-like)</i>	0–78	–
	Scribner's rosette grass	DIOLS	<i>Dichanthelium oligosanthos</i> var. <i>scribnerianum</i>	0–26	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0–26	–
6	Grass-likes			26–208	
	sedge	CAREX	<i>Carex</i>	26–208	–
	Grass-like (not a true grass)	2GL	<i>Grass-like (not a true grass)</i>	0–52	–
7	Non-Native Grasses			910–1950	
	Kentucky bluegrass	POPR	<i>Poa pratensis</i>	650–1690	–
	quackgrass	ELRE4	<i>Elymus repens</i>	130–520	–
	smooth brome	BRIN2	<i>Bromus inermis</i>	0–390	–
Forb					
8	Forbs			130–390	
	Canada thistle	CIAR4	<i>Cirsium arvense</i>	26–208	–
	Forb, introduced	2FI	<i>Forb, introduced</i>	26–182	–
	white clover	TRRE3	<i>Trifolium repens</i>	0–156	–
	common dandelion	TAOF	<i>Taraxacum officinale</i>	0–130	–
	goldenrod	SOLID	<i>Solidago</i>	26–104	–
	western yarrow	ACMIO	<i>Achillea millefolium</i> var. <i>occidentalis</i>	26–104	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	26–78	–
	scurfpea	PSORA2	<i>Psoraleidium</i>	26–78	–
	white heath aster	SYER	<i>Symphyotrichum ericoides</i>	26–78	–
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	0–52	–

	cinquefoil	POTEN	<i>Potentilla</i>	0–26	–
	Indianhemp	APCA	<i>Apocynum cannabinum</i>	0–26	–
	Forb, native	2FN	<i>Forb, native</i>	0–26	–
Shrub/Vine					
9	Shrubs			26–156	
	snowberry	SYMPH	<i>Symphoricarpos</i>	26–130	–
	rose	ROSA5	<i>Rosa</i>	0–52	–
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0–26	–
	American plum	PRAM	<i>Prunus americana</i>	0–26	–

Animal community

Animal Community – Grazing Interpretations The following table lists annual, suggested initial stocking rates with average growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of conservation planning. Often, the current plant composition does not entirely match any particular plant community (as described in this ES description). Because of this a resource inventory is necessary to document plant composition and production. More accurate carrying capacity estimates should eventually be calculated using the following stocking rate information along with animal preference data and actual stocking records, particularly when grazers other than cattle are involved. With consultation of the land manager, more intensive grazing management may result in improved harvest efficiencies and increased carrying capacity.

Big Bluestem/Indiangrass/Switchgrass (1.1 & 3.1)

Average Annual Production (lbs./acre, air-dry): 4600

Stocking Rate* (AUM/acre): 1.26

Little Bluestem/Kentucky Bluegrass/Big Bluestem (3.2)

Average Annual Production (lbs./acre, air-dry): 3400

Stocking Rate* (AUM/acre): 0.93

Smooth Bromeagrass/Kentucky Bluegrass (4.1)

Average Annual Production (lbs./acre, air-dry): 4000

Stocking Rate* (AUM/acre): 1.10

Kentucky Bluegrass/Quackgrass (4.2)

Average Annual Production (lbs./acre, air-dry): 2600

Stocking Rate* (AUM/acre): 0.71

Annual/Pioneer, Non-Native Perennial (4.3)

Average Annual Production (lbs./acre, air-dry): 1200

Stocking Rate* (AUM/acre): 0.33

*Based on 912 lbs./acre (air-dry weight) per Animal Unit Month (AUM), and on 25 percent harvest efficiency (refer to United States Department of Agriculture (USDA) Natural Resources Conservation Service (NRCS), National Range and Pasture Handbook).

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage. During the dormant period, the forage for livestock will likely be lacking protein to meet livestock requirements, and added protein will allow ruminants to better utilize the energy stored in grazed plant materials. A forage quality test (either directly or through fecal sampling) should be used to determine the level of supplementation needed.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B. Infiltration is typically moderate to moderately high and runoff potential for this site varies from low to medium

depending on soil hydrologic group, slope, and ground cover. In many cases, areas with greater than 75 percent ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where shortgrasses form a strong sod and dominate the site. Dominance by Kentucky bluegrass and/or smooth brome grass will result in reduced infiltration and increased runoff. Areas where ground cover is less than 50 percent have the greatest potential to have reduced infiltration and higher runoff (refer to Section 4, NRCS National Engineering Handbook for runoff quantities and hydrologic curves).

Recreational uses

This site provides hunting, hiking, photography, bird watching, and other opportunities. The wide varieties of plants that bloom from spring until fall have an esthetic value that appeals to visitors.

Wood products

No appreciable wood products are typically present on this site.

Other products

Seed harvest of native plant species can provide additional income on this site.

Inventory data references

MLRA 102D was created in 2022 with Agricultural Handbook 296 updated. This area was MLRA 102A prior to this time. Information was copied from MLRA 102A ESDs to create the MLRA 102D ESDs.

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range-trained personnel were also used. Those involved in developing this site include: Stan Boltz, Range Management Specialist, NRCS; and Bruce Kunze, Soil Scientist, NRCS.

Data Source Sample Period State County
NP-ESC-1 (0100746039) 2007 SD Deuel

Other references

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Conterminous United States. USDA Forest Service, General Technical Report WO-76. Washington, DC. 92 pps.

Gilbert, M. C., Whited, P. M., Clairain Jr, E. J., & Smith, R. D. (2006). A Regional Guidebook for Applying the Hydrogeomorphic Approach to Assessing Wetland Functions of Prairie Potholes. Washington DC.

Samson, F. B., & Knopf, F. L. (1996). Prairie Conservation Preserving North America's Most Endangered Ecosystem. Washington D.C.: Island Press.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Official Soil Series Descriptions. Available online. Accessed March 2018.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2003. National Range and Pasture Handbook, Revision 1. Grazing Lands Technology Institute. 214 pps.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. 672pps.

United States Department of Agriculture, Natural Resources Conservation Service. 2022. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture, Agriculture Handbook 296.

USDA, NRCS. National Soil Information System, Information Technology Center, 2150 Centre Avenue, Building A, Fort Collins, CO 80526. (<http://soils.usda.gov/technical/nasis/>)

USDA, NRCS. 2018. The PLANTS Database (<http://plants.usda.gov>, 27 March 2018). National Plant Data Team, Greensboro, NC 27401-4901 USA.

U.S. Environmental Protection Agency [EPA]. 2013. Level III and Level IV Ecoregions of the Continental United States. Corvallis, OR, U.S. EPA, National Health and Environmental Effects Research Laboratory, map scale 1:3,000,000. Available at <http://www.epa.gov/eco-research/level-iii-and-iv-ecoregions-continental-united-states>. (Accessed 1 March 2018).

Contributors

Megan Baxter
Stan Boltz
Lance Howe
Steve Winter

Approval

Suzanne Mayne-Kinney, 8/14/2024

Acknowledgments

Contact for Lead Authors: Natural Resources Conservation Service (USDA-NRCS), Redfield Soil Survey Office Redfield, SD; Lance Howe (Lance.Howe@usda.gov), Soil Survey Office Leader, USDA-NRCS, Redfield, SD; and Steve Winter (Steven.Winter@usda.gov), Soil Scientist, USDA-NRCS, Redfield, SD

Additional Information Acknowledgment: Jason Hermann (Jason.Hermann@usda.gov), Area Rangeland Management Specialist, USDA-NRCS, Redfield, SD.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	David Schmidt, Tim Nordquist, Stan Boltz
Contact for lead author	
Date	12/04/2007
Approved by	Suzanne Mayne-Kinney
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rills should not be present.

2. **Presence of water flow patterns:** Barely observable.

-
3. **Number and height of erosional pedestals or terracettes:** Essentially, non-existent.
-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground less than 5% and less than 2 inches in diameter.
-
5. **Number of gullies and erosion associated with gullies:** Active gullies should not be present.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** None.
-
7. **Amount of litter movement (describe size and distance expected to travel):** Little to no plant litter movement. Plant litter remains in place and is not moved by erosional forces.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Stability class usually 6. Typically high root content, organic matter, and granular structure. Soil surface is very resistant to erosion.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Use soil series description for depth and color of A-horizon.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Healthy, deep rooted native grasses enhance infiltration and reduce runoff.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer should be evident.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Tall warm-season rhizomatous grass >> tall cool-season bunch grass
- Sub-dominant: > mid warm-season bunch grass > short cool-season grass = forb > shrub > tree
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Very little to no evidence of decadence or mortality.

14. **Average percent litter cover (%) and depth (in):** 85-90%, roughly 0.5 inch thick or less. Litter cover is in contact with soil surface.

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 4000 – 5200 lbs./acre air-dry weight, average 4,600 lbs./acre air-dry weight

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Refer to State and Local Noxious Weed List, also Kentucky bluegrass, smooth brome grass

17. **Perennial plant reproductive capability:** All species are capable of reproducing.
