

# Ecological site R102DY036SD Saline Subirrigated

Last updated: 8/14/2024 Accessed: 05/12/2025

#### **General information**

**Provisional**. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

#### **MLRA** notes

Major Land Resource Area (MLRA): 102D-Prairie Coteau

This area makes up about 7,867 square miles (20,375 square kilometers), consisting mostly of nearly level to undulating till plains with potholes and moraines. Elevation ranges from 1,150 to 2,130 feet (350 to 650 meters). The average annual precipitation is 22 to 29 inches (559 to 734 millimeters). The average annual temperature is 42 to 45 degrees F (6 to 7 degrees C). The dominant soil order in this MLRA is Mollisols. The soils in this area dominantly have a frigid temperature regime, and an aquic or udic moisture regime. They are generally very deep and loamy. Soils range from well drained to very poorly drained. Parent materials are dominantly fine-loamy till to clayey material, with smaller amounts of outwash, glaciofluvial deposits, eolian deposits, alluvium, and, to a lesser extent, loess and organic materials.

#### **Classification relationships**

\*Fenneman (1916) Physiographic Regions\* Division - Interior Plains East: Province - Central Lowland Section - Western Lake / Dissected Till Plains (12b/12e)

\*USFS (2007) Ecoregions\* Domain - Humid Temperate Division - Prairie Province - Prairie Parkland (Temperate) Section - North-Central Glaciated Plains (251B)

\*EPA Ecoregions (Omernik 1997)\*
I - Great Plains (9)
II - Temperate Prairies (9.2)
III - Aspen Parkland/Northern Glaciated Plains (9.2.1)

#### **Ecological site concept**

The Saline Subirrigated ecological site typically occurs along the edges of drainageways or closed depressions. Soils are somewhat poorly drained which have a water table within 2 to 5 feet of the soil surface that persists longer than the wettest part of the growing season typically until the month of August. The soils will have visible salts within 16 inches of the soil surface. Dominant vegetation is adapted to the high salinity. Vegetation in the Reference State includes big bluestem, Indiangrass, and Switchgrass. Forbs include Pursh seepweed, goldenrods, cudweed sagewort, and cinquefoil. The site my become degraded due to change in disturbance regime, and vegetation may shift to community dominated by foxtail barley, inland saltgrass, and bareground.

## **Associated sites**

R102DY004SD	Wet Meadow These sites occur in a basin or closed depression. Soils are poorly drained and the site ponds water for 4 to 8 weeks in the spring of the year or after a heavy rain.
R102DY002SD	<b>Linear Meadow</b> These sites occur in drainageways. Soils are poorly and very poorly drained which have a water table within 0 to 2 feet of the soil surface that persists longer than the wettest part of the growing season typically until the month of August.
R102DY007SD	<b>Saline Lowland</b> These sites typically occur in drainageways, but can occur along the edges of larger closed depressions. Soils are poorly and very poorly drained which have a water table within 0 to 2 feet of the soil surface that persists longer than the wettest part of the growing season typically until the month of August. The soils will have visible salts within 16 inches of the soil surface.
R102DY003SD	<b>Subirrigated</b> These sites occur in drainageways. Soils are somewhat poorly drained which have a water table within 2 to 5 feet of the soil surface that persists longer than the wettest part of the growing season typically until the month of August.
R102DY006SD	<b>Limy Subirrigated</b> These sites occur along the edges of drainageways. Soils are somewhat poorly drained which have a water table within 2 to 5 feet of the soil surface that persists longer than the wettest part of the growing season typically until the month of August. Soils will effervesce with acid at or near the surface.

## Similar sites

R102DY006SD	Limy Subirrigated
	The Limy Subirrigated site is in a similar landscape position, but will not have visible salts within 16 inches
	of the soil surface. The Limy Subirrigated site will have less switchgrass and prairie cordgrass, and more
	needlegrasses than the Saline Subirrigated site.

#### Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	<ul><li>(1) Andropogon gerardii</li><li>(2) Sorghastrum nutans</li></ul>

## **Physiographic features**

This site typically occurs along the edges of drainageways or closed depressions.

#### Table 2. Representative physiographic features

Landforms	<ul><li>(1) Drainageway</li><li>(2) Closed depression</li></ul>
Runoff class	Low
Flooding duration	Brief (2 to 7 days)
Flooding frequency	None to frequent
Ponding frequency	None
Elevation	1,000–2,000 ft
Slope	0–2%
Water table depth	18–30 in
Aspect	W, NW, N, NE, E, SE, S, SW

#### **Climatic features**

The average annual precipitation is 22 to 28 inches. Half or more of the precipitation falls during the growing season. Rainfall typically occurs during high-intensity, convective thunderstorms in summer. In the western part of the MLRA, rainfall is less abundant and not always adequate for full maturation of crops. Precipitation in winter is typically snow. The average annual temperature is 42 to 45 degrees F. The freeze-free period averages 143 days and ranges from 131 to 151 days.

Frost-free period (characteristic range)	114-128 days
Freeze-free period (characteristic range)	139-149 days
Precipitation total (characteristic range)	24-27 in
Frost-free period (actual range)	110-131 days
Freeze-free period (actual range)	131-151 days
Precipitation total (actual range)	22-28 in
Frost-free period (average)	122 days
Freeze-free period (average)	143 days
Precipitation total (average)	25 in

#### Table 3. Representative climatic features



Figure 1. Monthly precipitation range



Figure 2. Monthly minimum temperature range



Figure 3. Monthly maximum temperature range



Figure 4. Monthly average minimum and maximum temperature



Figure 5. Annual precipitation pattern



Figure 6. Annual average temperature pattern

### **Climate stations used**

- (1) ROY LAKE [USC00397326], Lake City, SD
- (2) WAUBAY NWR [USC00398980], Waubay, SD
- (3) WEBSTER [USC00399004], Webster, SD

- (4) WATERTOWN 1W [USC00398930], Watertown, SD
- (5) WATERTOWN RGNL AP [USW00014946], Watertown, SD
- (6) CASTLEWOOD [USC00391519], Castlewood, SD
- (7) ARLINGTON 1 W [USC00390281], Arlington, SD
- (8) BROOKINGS 2 NE [USC00391076], Brookings, SD
- (9) ASTORIA 4S [USC00390422], White, SD
- (10) CLEAR LAKE [USC00391777], Clear Lake, SD
- (11) TYLER [USC00218429], Tyler, MN

### Influencing water features

No riparian areas or wetland features are directly associated with this site.

### **Soil features**

Soils are typically formed in loess over till. Surface textures are typically silty clay loam. Soils are somewhat poorly drained. These soils have visible salts within 16 inches of the surface.

Parent material	(1) Loess
Surface texture	(1) Silty clay loam
Drainage class	Somewhat poorly drained
Permeability class	Moderately rapid
Soil depth	80 in
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	4.6–6.6 in
Soil reaction (1:1 water) (0-10in)	7.4–8.4
Subsurface fragment volume <=3" (0-60in)	0–2%
Subsurface fragment volume >3" (0-60in)	0%

#### Table 4. Representative soil features

### **Ecological dynamics**

This site developed under Northern Great Plains climatic conditions, light to severe grazing by bison and other large herbivores, sporadic natural or man-caused wildfire (often of light intensities), fluctuating water tables and flooding events, and other biotic and abiotic factors that typically influence soil/site development. Changes will occur in the plant communities due to short-term weather variations, impacts of native and/or exotic plant and animal species, and management actions. While the following plant community descriptions describe more typical transitions that will occur, severe disturbances, such as periods of well below average precipitation, can cause significant shifts in plant communities and/or species composition that may not be described within this document.

Heavy continuous grazing without adequate recovery periods following each grazing occurrence over several years causes this site to depart from the interpretive plant community. Species such as little bluestem (*Schizachyrium scoparium*) and sedge (Carex) will initially increase. Big bluestem, Indiangrass, and switchgrass will decrease in frequency and production. Heavy, continuous grazing causes inland saltgrass (*Distichlis spicata*) to increase and eventually develop into a sod condition. Extended periods of nonuse and no fire will result in a plant community having high litter levels which favors an increase in species such as spikerush (*Eleocharis palustris*), sedge, foxtail barley (*Hordeum jubatum*), and prairie cordgrass (*Spartina pectinata*). Grazing, especially if adequate recovery periods are not allowed may be more detrimental on this site than haying. Biotic integrity on this site may be maintained more readily through periodic haying than through grazing.

Interpretations are primarily based on the 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase. It has been determined by study of rangeland relic areas, areas protected from excessive disturbance, and areas under long-term rotational grazing regimes. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts also have been used. Plant community phases, states, transitional pathways, and thresholds have been determined through similar studies and experience.

### State and transition model

Ecosystem states



T1 - Heavy continuous grazing

R2 - Long-term prescribed grazing, prescribed burning

#### State 1 submodel, plant communities



1.1A - Heavy continuous grazing

1.2A - Prescribed grazing with recovery periods

#### State 2 submodel, plant communities



2.1A - Heavy continuous grazing

2.2A - Time without disturbance

### State 1 Reference State

The Reference State represents the natural range of variability that dominates the dynamics of this ecological site (ES). This state is dominated by warm-season grasses. In pre-European times, the primary disturbance mechanisms for this site in the reference condition included periods of below and/or above average precipitation, periodic fire, and herbivory by insects and large ungulates. Timing of fires and herbivory coupled with weather events dictated the dynamics that occurred within the natural range of variability. In some locations, this site likely received relatively heavy grazing pressure. Tall warm-season grasses would have declined, and shorter-statured grass and grass-likes would have increased. Today, this state can be found on areas that are properly managed with grazing and/or prescribed burning and sometimes on areas receiving occasional short periods of rest.

#### **Dominant plant species**

- big bluestem (Andropogon gerardii), grass
- little bluestem (Schizachyrium scoparium), grass
- slender wheatgrass (Elymus trachycaulus), grass
- switchgrass (Panicum virgatum), grass
- sedge (Carex), grass
- white sagebrush (Artemisia ludoviciana), other herbaceous
- Maximilian sunflower (Helianthus maximiliani), other herbaceous
- goldenrod (Solidago), other herbaceous
- white heath aster (Symphyotrichum ericoides), other herbaceous

### Community 1.1 Big Bluestem-Indiangrass-Switchgrass

Interpretations are based primarily on the 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase (this is also considered to be climax). The potential vegetation is about 85 percent grasses or grass-like plants and 15 percent forbs. The community is dominated by warm-season grasses. The major grasses include big bluestem, little bluestem, Indiangrass, and switchgrass. Other grass or grass-like species include prairie cordgrass (*Spartina pectinata*), slender wheatgrass (*Elymus trachycaulus*), western wheatgrass (*Pascopyrum smithii*), sideoats grama (*Bouteloua curtipendula*), alkali sacaton (*Sporobolus airoides*), plains bluegrass (*Poa arida*), and sedge. This plant community is resilient and well adapted to the Northern Great Plains climatic conditions. The diversity in plant species allows for high drought tolerance. This is a sustainable plant community in regards to site/soil stability, watershed function, and biologic integrity.

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	3210	3960	4410
Forb	190	440	790
Total	3400	4400	5200

Table 5. Annual production by plant type



Figure 8. Plant community growth curve (percent production by month). SD0210, Rolling Till Prairie, lowland warm-season dominant.. Warm-season dominant, lowland..

### Community 1.2 Little Bluestem-Western Wheatgrass-Foxtail Barley-Inland Saltgrass

This plant community evolves under heavy continuous grazing or from over utilization during extended drought periods. The potential plant community is made up of approximately 90 percent grasses and grass-like species and 10 percent forbs. Dominant grass and grass-like species include little bluestem, western wheatgrass, foxtail barley, inland saltgrass, and slender wheatgrass. Grass and grass-like species of secondary importance include big bluestem, sedge, spikerush, plains bluegrass, prairie cordgrass, and switchgrass. Forbs commonly found in this plant community include Pursh seepweed (*Suaeda calceoliformis*), goldenrod (Solidago), cudweed sagewort (*Artemisia ludoviciana*), silverleaf cinquefoil (*Potentilla argentea*), alkali plantain (*Plantago eriopoda*), western ragweed (*Ambrosia psilostachya*), and annual marshelder (*Iva annua*). When compared to the 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase, slender wheatgrass, western wheatgrass, foxtail barley, inland

saltgrass, sedge, and grass-like species increase. Production of tall warm-season grasses is reduced. This plant community is moderately resistant to change. The herbaceous species present are well adapted to grazing; however, species composition can be altered through long-term overgrazing. If the herbaceous component is intact, it tends to be resilient if the disturbance is not long-term. Most of the components of the ecological processes will be functioning at optimum levels. However, the vigor and reproductive capability of the tall warm-season grasses will be reduced due to grazing pressure or a combination of stressors. A reduction of this dominant functional group allows for an increase in shorter-statured (and shallower rooted) species.

Table 6. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	2540	3330	3980
Forb	160	270	420
Total	2700	3600	4400



Figure 10. Plant community growth curve (percent production by month). SD0208, Rolling Till Prairie, lowland cool-season/warm-season codominant.. Cool-season, warm-season codominant, lowland..

### Pathway 1.1A Community 1.1 to 1.2

Heavy continuous grazing which includes herbivory at moderate to heavy levels at the same time of year each year without adequate recovery periods, or during periods of below normal precipitation when grazing frequency and intensity increases on these sites due to limited forage availability on adjacent upland sites will shift this community to the 1.2 Little Bluestem-Western Wheatgrass-Foxtail Barley-Inland Saltgrass Plant Community Phase.

## Pathway 1.2A Community 1.2 to 1.1

Prescribed grazing (alternating season of use and providing adequate recovery periods) or periodic light to moderate grazing possibly including periodic rest will convert this plant community to the 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase.

#### **Conservation practices**

Prescribed Grazing

### State 2 Degraded State

This state is characterized by the dominance of the shorter-statured, more saline tolerant species such as foxtail barley and inland saltgrass, the increase in bare ground, and the increased presence of salt accumulations on the soil surface. Infiltration is reduced, which allows the moisture and the salts carried by the moisture to be wicked up to the soil surface. The short-statured and shallow rooted species are more capable of withstanding the higher concentrations of salts in the soil surface. As the disturbance level increases, plant density decreases even more,

giving way to annual species and invasive perennial species, as well as, a further increase in bare ground.

#### **Dominant plant species**

- saltgrass (Distichlis spicata), grass
- foxtail barley (Hordeum jubatum), grass
- western wheatgrass (Pascopyrum smithii), grass
- spikerush (Eleocharis), grass
- prairie cordgrass (Spartina pectinata), grass
- Pursh seepweed (Suaeda calceoliformis), other herbaceous
- annual marsh elder (Iva annua), other herbaceous
- redwool plantain (Plantago eriopoda), other herbaceous

### Community 2.1 Foxtail Barley-Inland Saltgrass/Bareground

This plant community developed with heavy continuous season-long grazing where adequate recovery periods between grazing events were not allowed. Patches of inland saltgrass sod are typical and foxtail barley is well distributed throughout the community. Tall warm-season grasses are nearly absent, and little bluestem, slender wheatgrass, and western wheatgrass have been greatly reduced and may persist in remnant amounts, reduced in vigor. Bare ground may develop in micro lows where salt concentrations are highest. A white salt crust may form on the soil surface. The forb component is comprised of salt tolerant species such as Pursh seepweed and silverleaf cinquefoil. This plant community is resistant to change due to the grazing tolerance of inland saltgrass and increased surface salts. A significant amount of production and diversity has been lost when compared to 1.1 Big Bluestem-Indiangrass-Switchgrass Plant Community Phase. Loss of key warm-season grasses and increased bare ground has negatively impacted energy flow and nutrient cycling. Water infiltration is reduced significantly due to the shallow rooting depth of inland saltgrass and increased bare ground.

Table 7. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	1365	1880	2375
Forb	35	120	225
Total	1400	2000	2600



Figure 12. Plant community growth curve (percent production by month). SD0208, Rolling Till Prairie, lowland cool-season/warm-season codominant.. Cool-season, warm-season codominant, lowland..

### Community 2.2 Annual Pioneer-Pioneer Perennial

This plant community developed under continuous heavy grazing or other excessive disturbances. The potential plant community is made up of approximately 40 to 80 percent grasses and grass-like species and 20 to 60 percent forbs. The species present in this phase are highly variable but often include nonnative invasive and/or early seral species. Plant diversity is low (plant richness may be high but areas are often dominated by a few species). The ecological processes are difficult to restore because of the loss of plant diversity and overall soil disturbance. Soil

erosion is potentially very high because of the bare ground and shallow rooted herbaceous plant community. Water runoff will increase and infiltration will decrease due to animal related soil compaction and loss of root mass due to low plant diversity and vigor. This plant community will require significant economic inputs and time to move towards another plant community. This movement is highly variable in its succession. This is due to the loss of diversity (including the loss of the seed bank), within the existing plant community, and the plant communities on adjacent sites.

## Pathway 2.1A Community 2.1 to 2.2

Heavy continuous grazing which includes herbivory at moderate to heavy levels at the same time of year each year without adequate recovery periods, or during periods of below normal precipitation when grazing frequency and intensity increases on these sites due to limited forage availability on adjacent upland sites will shift this community to the 2.2 Annual Pioneer-Perennial Pioneer Plant Community Phase.

## Pathway 2.2A Community 2.2 to 2.1

This community pathway occurs with the passage of time as successional processes take place and perennial plants gradually begin to establish on the site again. This pathway will lead to the 2.1 Foxtail Barley-Inland Saltgrass/Bareground Plant Community Phase.

## Transition T1 State 1 to 2

Heavy continuous grazing (stocking levels well above carrying capacity for extended portions of the growing season and often at the same time of year each year), will likely lead this 1.2 Little Bluestem-Western Wheatgrass-Foxtail Barley-Inland Saltgrass Plant Community Phase within the Reference State (State 1) over a threshold leading to the 2.1 Foxtail Barley-Inland Saltgrass-Bareground Plant Community Phase within the Degraded State (State 2). Grazing repeatedly in the early growing season can expedite this shift by causing mechanical disturbance due to trampling.

### Restoration pathway R2 State 2 to 1

Long-term prescribed grazing (moderate stocking levels coupled with adequate recovery periods, or other grazing systems such as high-density, low-frequency intended to treat specific species dominance, or periodic light to moderate stocking levels possibly including periodic rest) coupled with prescribed burning may lead this plant community phase over a threshold to the Reference State (State 1). Wetland restoration techniques may be necessary to restore biotic integrity and plant diversity and productivity.

### **Conservation practices**

Prescribed Grazing

### Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)		
Grass	Grass/Grasslike						
1	Tall Warm-season Grasses			1100–2420			
	big bluestem	ANGE	Andropogon gerardii	440–1320	-		
	switchgrass	PAVI2	Panicum virgatum	220–660	-		
	Indiangrass	SONU2	Sorghastrum nutans	220–660	-		
	prairie cordgrass	SPPE	Spartina pectinata	44–440	_		

	I' ~	I .	, , , , , , , , , , , , , , , , , , ,	1 1	
2	Mid Warm-season Grasse	S		660–1320	
	little bluestem	SCSC	Schizachyrium scoparium	660–1320	_
	alkali sacaton	SPAI	Sporobolus airoides	0–220	_
	sideoats grama	BOCU	Bouteloua curtipendula	0–220	_
3	Cool-season Grasses	<u>.</u>		220–440	
	slender wheatgrass	ELTR7	Elymus trachycaulus	44–440	_
	western wheatgrass	PASM	Pascopyrum smithii	44–440	_
	plains bluegrass	POAR3	Poa arida	44–220	-
	foxtail barley	HOJU	Hordeum jubatum	0–44	-
4	Short Warm-season Grass	ses		44–88	
	saltgrass	DISP	Distichlis spicata	44–88	_
	mat muhly	MURI	Muhlenbergia richardsonis	0–44	-
5	Other Native Grasses			44–220	
	Graminoid (grass or grass- like)	2GRAM	Graminoid (grass or grass-like)	0–220	-
	prairie Junegrass	KOMA	Koeleria macrantha	44–132	-
	Scribner's rosette grass	DIOLS	Dichanthelium oligosanthes var. scribnerianum	0-44	_
6	Grass-likes			88–352	
	sedge	CAREX	Carex	44–352	_
	spikerush	ELEOC	Eleocharis	0–132	_
	Grass-like (not a true grass)	2GL	Grass-like (not a true grass)	0–88	_
Forb		1			
7	Forbs			220–660	
	Forb, native	2FN	Forb, native	44–220	_
	white sagebrush	ARLU	Artemisia ludoviciana	44–132	-
	Maximilian sunflower	HEMA2	Helianthus maximiliani	44–132	-
	goldenrod	SOLID	Solidago	44–132	_
	white heath aster	SYER	Symphyotrichum ericoides	44–88	_
	silver cinquefoil	POAR8	Potentilla argentea	44–88	_
	upright prairie coneflower	RACO3	Ratibida columnifera	44–88	_
	Indianhemp	APCA	Apocynum cannabinum	44–88	_
	tall blazing star	LIAS	Liatris aspera	44–88	_
	western marbleseed	ONBEO	Onosmodium bejariense var. occidentale	44–88	-
	Flodman's thistle	CIFL	Cirsium flodmanii	44–88	-
	western yarrow	ACMIO	Achillea millefolium var. occidentalis	44–88	_
	Cuman ragweed	AMPS	Ambrosia psilostachya	44–88	_
	Canadian anemone	ANCA8	Anemone canadensis	0–44	_
	smooth horsetail	EQLA	Equisetum laevigatum	0–44	_
	bluebell bellflower	CARO2	Campanula rotundifolia	0–44	_
	redwool plantain	PLER	Plantago eriopoda	0-44	
	palespike lobelia	LOSP	Lobelia spicata	0–44	
	rough bugleweed	LYAS	Lycopus asper	0–44	

annual marsh elder	IVAN2	Iva annua	0–44	-
western dock	RUAQ	Rumex aquaticus	0–44	-
Norwegian cinquefoil	PONO3	Potentilla norvegica	0–44	-
prairie violet	VIPE2	Viola pedatifida	0–44	-
meadow zizia	ZIAP	Zizia aptera	0–44	-
Pursh seepweed	SUCA2	Suaeda calceoliformis	0–44	-

#### Table 9. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass	/Grasslike	•	•		
1	Tall Warm-season Grasses			72–540	
	big bluestem	ANGE	Andropogon gerardii	36–360	_
	prairie cordgrass	SPPE	Spartina pectinata	0–252	_
	switchgrass	PAVI2	Panicum virgatum	0–180	_
	Indiangrass	SONU2	Sorghastrum nutans	0–108	_
2	Mid Warm-season Grasses		180–1260		
	little bluestem	SCSC	Schizachyrium scoparium	180–1260	-
	alkali sacaton	SPAI	Sporobolus airoides	0–108	-
3	Cool-season Grasses		360–1080		
	western wheatgrass	PASM	Pascopyrum smithii	180–720	-
	slender wheatgrass	ELTR7	Elymus trachycaulus	72–540	-
	foxtail barley	HOJU	Hordeum jubatum	36–360	-
	plains bluegrass	POAR3	Poa arida	0–144	-
4	Short Warm-season Grasses		180–540		
	saltgrass	DISP	Distichlis spicata	180–540	-
	mat muhly	MURI	Muhlenbergia richardsonis	0–108	-
5	Other Native Grasses		0–180		
	Graminoid (grass or grass- like)	2GRAM	Graminoid (grass or grass-like)	0–144	_
	prairie Junegrass	KOMA	Koeleria macrantha	0–36	_
6	Grass-likes			180–720	
	sedge	CAREX	Carex	108–540	_
	spikerush	ELEOC	Eleocharis	36–360	_
	Grass-like (not a true grass)	2GL	Grass-like (not a true grass)	0–108	-
Forb					
7	Forbs			40–200	
	Pursh seepweed	SUCA2	Suaeda calceoliformis	36–108	_
	Forb, native	2FN	Forb, native	36–108	-
	western yarrow	ACMIO	Achillea millefolium var. occidentalis	0–108	-
	Cuman ragweed	AMPS	Ambrosia psilostachya	36–72	-
	white sagebrush	ARLU	Artemisia ludoviciana	0–72	-
	redwool plantain	PLER	Plantago eriopoda	0–72	
	silver cinquefoil	POAR8	Potentilla argentea	36–72	
	annual march aldar	I\/AN/2	בווחתב בעו	∩_72	_

I			iva annua	v− <i>i</i> ∠	_
	Forb, introduced	2FI	Forb, introduced	0–72	-
	goldenrod	SOLID	Solidago	0–72	
	tall blazing star	LIAS	Liatris aspera	0–36	-
	western marbleseed	ONBEO	Onosmodium bejariense var. occidentale	0–36	-
	Norwegian cinquefoil	PONO3	Potentilla norvegica	0–36	-
	upright prairie coneflower	RACO3	Ratibida columnifera	0–36	-
	western dock	RUAQ	Rumex aquaticus	0–36	-
	Flodman's thistle	CIFL	Cirsium flodmanii	0–36	-
	smooth horsetail	EQLA	Equisetum laevigatum	0–36	-
	American licorice	GLLE3	Glycyrrhiza lepidota	0–36	-
	Maximilian sunflower	HEMA2	Helianthus maximiliani	0–36	-
	Indianhemp	APCA	Apocynum cannabinum	0–36	-
	white heath aster	SYER	Symphyotrichum ericoides	0–36	_

Table 10. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass	/Grasslike	•			
1	Tall Warm-season Grasses			0–100	
	prairie cordgrass	SPPE	Spartina pectinata	0–100	
2	Mid Warm-season Grasses	•		0–100	
	little bluestem	SCSC	Schizachyrium scoparium	0–100	_
	alkali sacaton	SPAI	Sporobolus airoides	0–40	_
3	Cool-season Grasses		200–700		
	foxtail barley	HOJU	Hordeum jubatum	200–700	_
	western wheatgrass	PASM	Pascopyrum smithii	0–160	_
	slender wheatgrass	ELTR7	Elymus trachycaulus	0–100	_
	plains bluegrass	POAR3	Poa arida	0–60	_
4	Short Warm-season Grasse	s		600–1100	
	saltgrass	DISP	Distichlis spicata	500–1000	_
	mat muhly	MURI	Muhlenbergia richardsonis	20–160	_
5	Other Native Grasses		•	0–40	
	Graminoid (grass or grass- like)	2GRAM	Graminoid (grass or grass-like)	0–40	_
6	Grass-likes			20–160	
	spikerush	ELEOC	Eleocharis	20–140	_
	sedge	CAREX	Carex	0–100	-
	Grass-like (not a true grass)	2GL	Grass-like (not a true grass)	0–40	_
Forb					
7	Forbs			40–200	
	Pursh seepweed	SUCA2	Suaeda calceoliformis	20–100	-
	Forb, introduced	2FI	Forb, introduced	0–60	_
	Forb, native	2FN	Forb, native	20–60	-
	annual marsh elder	IVAN2	lva annua	0–60	_
	redwool plantain	PLER	Plantago eriopoda	20–60	-
	silver cinquefoil	POAR8	Potentilla argentea	20–60	-
	Norwegian cinquefoil	PONO3	Potentilla norvegica	0–20	_
	goldenrod	SOLID	Solidago	0–20	-
	western yarrow	ACMIO	Achillea millefolium var. occidentalis	0–20	_
	Cuman ragweed	AMPS	Ambrosia psilostachya	0–20	_
	Flodman's thistle	CIFL	Cirsium flodmanii	0–20	
	smooth horsetail	EQLA	Equisetum laevigatum	0–20	_

## **Animal community**

#### Animal Community – Grazing Interpretations

The following table lists annual, suggested initial stocking rates with average growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of conservation planning. Often, the current plant composition does not entirely match any particular plant community (as described in this ES description). Because of this, a resource inventory is necessary to document plant composition and production. More accurate carrying capacity estimates should eventually be calculated using the following stocking rate

information along with animal preference data and actual stocking records, particularly when grazers other than cattle are involved. With consultation of the land manager, more intensive grazing management may result in improved harvest efficiencies and increased carrying capacity.

Bluestem/Indiangrass/Switchgrass (1.1) Average Annual Production (lbs./acre, air-dry): 4400 Stocking Rate\* (AUM/acre): 1.21

Little Bluestem/Wheatgrass/Foxtail Barley/Inland Saltgrass (1.2) Average Annual Production (lbs./acre, air-dry): 3600 Stocking Rate\* (AUM/acre): 0.99

Foxtail Barley/Inland Saltgrass, Bare Ground (2.1) Average Annual Production (lbs./acre, air-dry): 2000 Stocking Rate\* (AUM/acre): 0.55

Annual/Pioneer, Non-Native Perennial, Bare Ground (2.2) Average Annual Production (lbs./acre, air-dry): 1200 Stocking Rate\* (AUM/acre): 0.33

\*Based on 912 lbs./acre (air-dry weight) per Animal Unit Month (AUM), and on 25 percent harvest efficiency (refer to United States Department of Agriculture (USDA) Natural Resources Conservation Service (NRCS), National Range and Pasture Handbook).

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage. During the dormant period, the forage for livestock will likely be lacking protein to meet livestock requirements and added protein will allow ruminants to better utilize the energy stored in grazed plant materials. A forage quality test (either directly or through fecal sampling) should be used to determine the level of supplementation needed.

### Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic groups C and D. Infiltration is typically moderate to moderately slow and runoff potential for this site varies from negligible to low depending on soil hydrologic group, slope, and ground cover. In many cases, areas with greater than 75 percent ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where shortgrasses form a strong sod and dominate the site. Areas where ground cover is less than 50 percent have the greatest potential to have reduced infiltration and higher runoff (refer to Section 4, NRCS National Engineering Handbook for runoff quantities and hydrologic curves).

#### **Recreational uses**

This site provides hunting, hiking, photography, bird watching, and other opportunities. The wide varieties of plants that bloom from spring until fall have an esthetic value that appeals to visitors.

#### Wood products

No appreciable wood products are typically present on this site.

#### **Other products**

Seed harvest of native plant species can provide additional income on this site.

#### Inventory data references

MLRA 102D was created in 2022 with Agricultural Handbook 296 updated. This area was MLRA 102A prior to this time. Information was copied from MLRA 102A ESDs to create the MLRA 102D ESDs.

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range-trained personnel were also used. Those involved in developing this site include: Stan Boltz, Range Management Specialist, NRCS; and Bruce Kunze, Soil Scientist, NRCS.

### **Other references**

High Plains Regional Climate Center, University of Nebraska, 830728 Chase Hall, Lincoln, NE 68583-0728. (http://www.hprcc.unl.edu/)

United States Department of Agriculture, Natural Resources Conservation Service. 2022. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture, Agriculture Handbook 296.

USDA, NRCS. National Water and Climate Center, 101 SW Main, Suite 1600, Portland, OR 97204-3224. (http://www.wcc.nrcs.usda.gov)

USDA, NRCS. National Range and Pasture Handbook, September 1997

USDA, NRCS. National Soil Information System, Information Technology Center, 2150 Centre Avenue, Building A, Fort Collins, CO 80526. (http://soils.usda.gov/technical/nasis/)

USDA, NRCS. 2001. The PLANTS Database, Version 3.1 (http://plants.usda.gov). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

### Contributors

Megan Baxter Stan Boltz

### Approval

Suzanne Mayne-Kinney, 8/14/2024

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	David Schmidt, Tim Nordquist, Stan Boltz
Contact for lead author	
Date	12/04/2007
Approved by	Suzanne Mayne-Kinney
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

#### Indicators

1. Number and extent of rills: Rills should not be present.

- 2. Presence of water flow patterns: Barely observable.
- 3. Number and height of erosional pedestals or terracettes: Essentially, non-existent.
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): Bare ground less than 5% and less than 2 inches in diameter.
- 5. Number of gullies and erosion associated with gullies: Active gullies should not be present.
- 6. Extent of wind scoured, blowouts and/or depositional areas: None.
- 7. Amount of litter movement (describe size and distance expected to travel): Little to no plant litter movement. Plant litter remains in place and is not moved by erosional forces.
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values): Stability class usually 6. Typically high root content, organic matter, and granular structure. Soil surface is very resistant to erosion.
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Use soil series description for depth and color of A-horizon.
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Healthy, deep rooted native grasses enhance infiltration and reduce runoff.
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): No compaction layer should be evident.
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Mid warm-season bunch grass >> tall warm-season rhizomatous grass

Sub-dominant: > mid cool-season rhizomatous grass > short warm-season rhizomatous grass = short cool-season grass = forb

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Very little to no evidence of decadence or mortality.
- 14. Average percent litter cover (%) and depth ( in): 85-90%, roughly 0.5 inch thick or less. Litter cover is in contact with soil surface.
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction): 3800 – 5000 lbs./acre air-dry weight, average 4,400 lbs./acre air-dry weight
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Refer to State and Local Noxious Weed List
- 17. Perennial plant reproductive capability: All species are capable of reproducing.