

## **Ecological site F108XA005IL Sandstone Woodland**

Last updated: 11/05/2024  
Accessed: 05/11/2025

---

### **General information**

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### **MLRA notes**

Major Land Resource Area (MLRA): 108X–Illinois and Iowa Deep Loess and Drift

The Illinois and Iowa Deep Loess and Drift, Eastern Part (MLRA 108A) encompasses the Grand Prairie physiographic division (Schewman et al. 1973). It spans two states – Illinois (97 percent) and Indiana (3 percent) – comprising about 11,145 square miles (Figure 1). The elevation ranges from 985 feet above sea level (ASL) in the northern part to 660 feet above sea level in the southern part. Local relief varies from 3 to 10 feet on most of the area which is on broad flat uplands. The maximum relief is about 160 feet along major streams. The northern part of this area is underlain by Ordovician and Silurian limestone and the southern part is underlain by Pennsylvanian shale, siltstone, and limestone. Except for some areas along streams where bedrock is exposed, glacial drift covers all the MLRA. The glacial drift consists of till and stratified outwash and is of Wisconsinan age. A moderately thin to thick layer of loess covers the entire area (USDA-NRCS 2006).

The vegetation in the MLRA has undergone drastic changes over time. At the end of the last glacial episode – the Wisconsinan glaciation – the evolution of vegetation began with the development of tundra habitats, followed by a phase of spruce and fir forests, and eventually spruce-pine forests. Not until approximately 9,000 years ago did the climate undergo a warming trend which prompted the development of deciduous forests dominated by oak and hickory. As the climate continued to warm and dry, prairies began to develop approximately 8,300 years ago. Another shift in climate that resulted in an increase in moisture prompted the emergence of savanna-like habitats from 8,000 to 5,000 years before present. Moisture continued to increase in the southernmost region 5,000 years ago, resulting in an increase of forested systems (Taft et al. 2009). Fire, droughts, and grazing by native mammals helped to maintain the prairies and savannas until the arrival of European settlers, and the forests were maintained by droughts, wind, lightning, and occasional fire (Taft et al. 2009; NatureServe 2018).

### **Classification relationships**

USFS Subregions: Central Till Plains and Grand Prairies (251D) and Central Till Plains-Beech-Maple Sections; Northern Grand Prairie (251Dc), Eastern Grand Prairie (251Dd), Southern Grand Prairie (251De), and Entrenched Valleys (222Hf) Subsections (Cleland et al. 2007)

U.S. EPA Level IV Ecoregion: Illinois/Indiana Prairies (54a) and Glaciated Wabash Lowlands (72b) (USEPA 2013)

National Vegetation Classification – Ecological Systems: North-Central Interior Dry Oak Forest and Woodland (CES202.047) (NatureServe 2018)

National Vegetation Classification – Plant Associations: *Quercus stellata* – *Quercus velutina*/Schizachyrium scoparium Woodland (CEGL005281) (Nature Serve 2018)

Biophysical Settings: North-Central Dry Oak Forest and Woodland (BpS 4913110) (LANDFIRE 2009)

Illinois Natural Areas Inventory: Dry barren (White and Madany 1978)

Ecological site concept

Sandstone Woodlands are located within the blue areas on the map (Figure 1). They occur on uplands. The soils are Alfisols and Entisols that are well to excessively-drained and moderately deep, formed in loess over residuum weathered from sandstone.

The historic pre-European settlement vegetation on this ecological site was dominated by an open oak canopy and sparse herbaceous layer. Black oak (*Quercus velutina* Lam.) and blackjack oak (*Quercus marilandica* Münchh.) are the dominant species on the site, and little bluestem (*Schizachyrium scoparium* (Michx.) Nash.) is the dominant grass. The moderately deep, acidic soils over bedrock result in a less productive plant community with smaller, stunted trees and an herbaceous layer that is generally less than 3 feet tall (White and Madany 1978). Fire is the primary disturbance that maintains this ecological site, and drought, storm damage, and periodic pest influences are secondary disturbances (LANDFIRE 2009).

Associated sites

R108XA004IL	<b>Sandstone Prairie</b> Drift over sandstone parent material including Channahon and Hesch soils
-------------	--

Similar sites

F108XA003IL	<b>Shale Woodland</b> Shale Woodlands are in a similar landscape position, but the soils are shallow to shale bedrock
-------------	--

Table 1. Dominant plant species

Tree	(1) <i>Quercus velutina</i> (2) <i>Quercus marilandica</i>
Shrub	Not specified
Herbaceous	(1) <i>Schizachyrium scoparium</i>

Physiographic features

Sandstone Woodlands occur on uplands in fire-protected landscapes. They are situated on elevations ranging from approximately 470 to 1020 feet. The site does not experience flooding but rather generates runoff to adjacent, downslope ecological sites (Table 1).




Figure 1. Figure 1. Location of Sandstone Woodland ecological site within MLRA 108A.

Table 2. Representative physiographic features

Slope shape across	(1) Convex
Slope shape up-down	(1) Convex
Landforms	(1) Upland > Hillslope (2) Terrace (3) Outwash plain
Runoff class	Low to high
Flooding frequency	None
Ponding frequency	None
Elevation	470–1,020 ft
Slope	2–9%
Water table depth	80 in
Aspect	Aspect is not a significant factor

## Climatic features


The Illinois and Iowa Deep Loess and Drift, Eastern Part falls into the hot-summer humid continental climate (Dfa) and the humid subtropical continental climate (Cfa) Köppen-Geiger climate classifications (Peel et al. 2007). The two main factors that drive the climate of the MLRA are latitude and weather systems. Latitude, and the subsequent reflection of solar input, determines air temperatures and seasonal variations. Solar energy varies across the seasons, with summer receiving three to four times as much energy as opposed to winter. Weather systems (air masses and cyclonic storms) are responsible for daily fluctuations of weather conditions. High-pressure systems are responsible for settled weather patterns where sun and clear skies dominate. In fall, winter, and spring, the polar jet stream is responsible for the creation and movement of low-pressure systems. The clouds, winds, and precipitation associated with a low-pressure system regularly follow high-pressure systems every few days (Angel n.d.).

The soil temperature regime of MLRA 108A is classified as mesic, where the mean annual soil temperature is between 46 and 59°F (USDA-NRCS 2006). Temperature and precipitation occur along a north-south gradient, where temperature and precipitation increase the further south one travels. The average freeze-free period of this ecological site is about 141 days, while the frost-free period is about 179 days (Table 2). The majority of the precipitation occurs as rainfall in the form of convective thunderstorms during the growing season. Average annual precipitation is approximately 37 inches, which includes rainfall plus the water equivalent from snowfall (Table 3). The average annual low and high temperatures are 40 and 60°F, respectively.


Climate data and analyses are derived from 30-year averages gathered from two National Oceanic and Atmospheric Administration (NOAA) weather stations contained within the range of this ecological site (Table 4).

**Table 3. Representative climatic features**


Frost-free period (characteristic range)	139-144 days
Freeze-free period (characteristic range)	173-185 days
Precipitation total (characteristic range)	36-37 in
Frost-free period (actual range)	137-146 days
Freeze-free period (actual range)	170-188 days
Precipitation total (actual range)	36-38 in
Frost-free period (average)	142 days
Freeze-free period (average)	179 days
Precipitation total (average)	37 in




**Figure 2. Monthly precipitation range**




**Figure 3. Monthly minimum temperature range**




**Figure 4. Monthly maximum temperature range**



**Figure 5. Monthly average minimum and maximum temperature**



**Figure 6. Annual precipitation pattern**




**Figure 7. Annual average temperature pattern**

## Climate stations used

- (1) OTTAWA 5SW [USC00116526], Ottawa, IL
- (2) PERU [USC00116753], La Salle, IL

## Influencing water features

Sandstone Woodlands are not influenced by wetland or riparian water features. Precipitation is the main source of water for this ecological site. Infiltration is high or low (Hydrologic Groups A and C), and surface runoff is low to high. Surface runoff contributes some water to downslope ecological sites (Figure 4).




**Figure 8. Figure 4. Hydrologic cycling in Sandstone Woodland ecological site.**

## Soil features

Soils of Sandstone Woodlands are in the Alfisols and Entisols orders, further classified as Typic Hapludalfs and Typic Quartzipsamments with slow to high infiltration and low to high runoff potential. The soil series associated

with this site includes Boone, Gale, and High Gap (Figure 5). The parent material is loess over residuum weathered from sandstone, and the soils are well to excessively-drained and moderately deep. Soil pH classes are very strongly acid to neutral. A paralithic contact is noted as a rooting restriction for the soils of this ecological site (Table 5).



**Figure 9. Figure 5. Profile sketches of soil series associated with Sandstone Woodland.**

**Table 4. Representative soil features**

Parent material	(1) Loess
Surface texture	(1) Loamy fine sand (2) Silt loam
Family particle size	(1) Fine-silty over sandy or sandy-skeletal (2) Fine-loamy (3) Not used
Drainage class	Well drained to excessively drained
Permeability class	Slow to moderately slow
Depth to restrictive layer	20–40 in
Soil depth	20–40 in
Surface fragment cover <=3"	0–8%
Surface fragment cover >3"	0%
Available water capacity (Depth not specified)	6.1–7.3 in
Soil reaction (1:1 water) (Depth not specified)	4.5–7.3
Subsurface fragment volume <=3" (Depth not specified)	3–12%
Subsurface fragment volume >3" (Depth not specified)	0–2%

## Ecological dynamics

The information in this Ecological Site Description, including the state-and-transition model (STM), was developed based on historical data, current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

The MLRA lies within the tallgrass prairie ecosystem of the Midwest. The heterogeneous topography of the area results in variable microclimates and fuel matrices that in turn support prairies, savannas, and forests. Sandstone

Woodlands form an aspect of this vegetative continuum. This ecological site occurs on uplands on well to excessively-drained soils that are moderately deep to sandstone bedrock. Species characteristic of this ecological site consist of a scrubby, open oak canopy and sparse herbaceous vegetation.


Fire is a critical factor that maintains Sandstone Woodlands. Fire typically consisted of low-severity surface fires every 10 years (LANDFIRE 2009). Ignition sources included summertime lightning strikes from convective storms and bimodal, human ignitions during the spring and fall seasons. Native Americans regularly set fires to improve sight lines for hunting, drive large game, improve grazing and browsing habitat, agricultural clearing, and enhance vital ethnobotanical plants (Barrett 1980; LANDFIRE 2009).

Drought and storm damage have also played a role in shaping this ecological site. The periodic episodes of reduced soil moisture in conjunction with the well to excessively-drained acidic soils have favored the proliferation of plant species tolerant of such conditions. Drought can also slow the growth of plants and result in dieback of certain species. Damage to trees from storms and pest outbreaks can vary from minor, patchy effects of individual trees to stand effects that temporarily affect community structure and species richness and diversity (Irland 2000; Peterson 2000). When coupled with fire, periods of drought and catastrophic storm damage can greatly delay the establishment and maturation of woody vegetation (Pyne et al. 1996).

Today, Sandstone Woodlands have been reduced from their pre-settlement extent. Low to moderate slopes have been converted to cropland, while steeper slopes have been converted to forage land. Remnants that do exist have experienced long-term fire suppression and overbrowsing resulting in significant changes to the forest structure. A return to the historic plant community may not be possible following extensive land modification, but long-term conservation agriculture or forest reconstruction efforts can help to restore some biotic diversity and ecological function. The state-and-transition model that follows provides a detailed description of each state, community phase, pathway, and transition. This model is based on available experimental research, field observations, literature reviews, professional consensus, and interpretations.

## **State and transition model**

## F108AY005IL SANDSTONE WOODLAND



Code	Process
T1A, T3A, T4A, T5A	Long-term fire suppression and/or land abandonment
1.1A	Fire return interval greater than 25 years
1.2A	Mixed-severity fire
T1B, T2A, T4B, T5B	Cultural treatments are implemented to increase forage quality and yield
3.1A	Mechanical harvesting is replaced with domestic livestock and continuous grazing
3.1B	Mechanical harvesting is replaced with domestic livestock and rest-rotational grazing
3.2A, 3.3B	Domestic livestock grazing is replaced by mechanical harvesting
3.2B	Implementation of rest-rotational grazing
3.3A	Implementation of continuous grazing
T1C, T2B, T3B, T5C	Agricultural conversion via tillage, seeding, and non-selective herbicide
4.1A	Less tillage, residue management
4.1B	Less tillage, residue management, and implementation of cover cropping
4.2B	Implementation of cover cropping
4.2A, 4.3B	Intensive tillage, remove residue, and reinitiate monoculture row cropping
4.3A	Remove cover cropping
R2A, R3A, R4A	Site preparation, tree planting, non-native species control, and native seeding
5.1A	Invasive species control and implementation of disturbance regimes
5.2A	Drought or improper timing/use of management actions

### State 1

#### Reference State

The reference plant community is categorized as an open woodland community, dominated by scrubby oaks and sparse herbaceous vegetation. The two community phases within the reference state are dependent on periodic



fire. The severity and intensity of fire alters species composition, cover, and extent. Drought, storm damage, and periodic pest outbreaks have more localized impacts in the reference phases, but do contribute to overall species composition, diversity, cover, and productivity.

## **Community 1.1**

### **Black Oak - Blackjack Oak/Little Bluestem**

Sites in this reference community phase are an open canopy woodland. Black oak and blackjack oak are the dominant trees on the site. Trees are large (21 to 33-inch DBH), and cover is 11 to 60 percent (LANDFIRE 2009). The herbaceous layer can be sparse with grasses less than 3 feet tall, including little bluestem, poverty oatgrass (*Danthonia spicata*(L.) P. Beauv. Ex Roem. & Schult.), prairie Junegrass (*Koeleria macrantha*(Ledeb.) Schult.), and Pennsylvania sedge (*Carex pensylvanica*Lam.) (White and Madany 1978). Surface fires every 10 years will maintain this community phase, but an extended fire return interval will shift the community to phase 1.2 (LANDFIRE 2009).

#### **Dominant plant species**

- black oak (*Quercus velutina*), tree
- blackjack oak (*Quercus marilandica*), tree
- little bluestem (*Schizachyrium scoparium*), grass

## **Community 1.2**

### **Black Oak - Blackjack Oak/Shagbark Hickory/Little Bluestem**

This reference community phase represents a fire return interval greater than 25 years. Black oak and blackjack oak remain the canopy dominants, but shagbark hickory (*Carya ovata* (Mill.) K. Koch) can begin to become prominent in the subcanopy. Tree size class remains large, but the canopy closes to 61 to 80 percent. Surface fires every 10 years will maintain this community phase, but a mixed-severity fire will shift the community back to phase 1.1 (LANDFIRE 2009).

#### **Dominant plant species**

- black oak (*Quercus velutina*), tree
- blackjack oak (*Quercus marilandica*), tree
- shagbark hickory (*Carya ovata*), shrub
- little bluestem (*Schizachyrium scoparium*), grass

## **Pathway 1.1A**

### **Community 1.1 to 1.2**

Fire return interval greater than 25 years.

## **Pathway 1.2A**

### **Community 1.2 to 1.1**

Mixed-severity fire event.

## **State 2**

### **Fire-Suppressed Forest State**

Fire suppression can transition the reference plant community from an oak forest to an oak-maple mesophytic forest. As the natural fire regime is removed from the landscape, encroachment and dominance by shade-tolerant, fire-intolerant species ensues. This results in a positive feedback loop of mesophication whereby plant community succession continuously creates cool, damp shaded conditions that perpetuate a closed canopy ecosystem (Nowacki and Abrams 2008). Succession to this forested state can occur in as little as 80 years from the last fire (LANDFIRE 2009).

## **Community 2.1**

## **Black Oak - Sugar Maple/Honeysuckle - Common Buckthorn/Pennsylvania Sedge**

This community phase represents the early stages of long-term fire suppression. Mature black and blackjack oaks and shagbark hickory are still present, but the more shade tolerant sugar maple (*Acer saccharum* Marshall) begins to co-dominate. The tree canopy closes to 100 percent cover and basal area increases (LANDFIRE 2009). Non-native shrubs, such as honeysuckle (*Lonicera*L.) and common buckthorn (*Rhamnus cathartica*L.), can rapidly colonize. The herbaceous layer shifts to shade-tolerant species, and diversity is reduced.

### **Dominant plant species**

- black oak (*Quercus velutina*), tree
- sugar maple (*Acer saccharum*), tree
- honeysuckle (*Lonicera*), shrub
- common buckthorn (*Rhamnus cathartica*), shrub
- Pennsylvania sedge (*Carex pensylvanica*), other herbaceous

## **State 3**

### **Forage State**

The forage state occurs when the reference state is converted to a farming system that emphasizes domestic livestock production known as grassland agriculture. Fire suppression, periodic cultural treatments (e.g., clipping, drainage, soil amendment applications, planting new species and/or cultivars, mechanical harvesting) and grazing by domesticated livestock transition and maintain this state (USDA-NRCS 2003). Early settlers seeded non-native species, such as smooth brome (*Bromus inermis* Leyss.) and Kentucky bluegrass (*Poa pratensis* L.), to help extend the grazing season. Over time, as lands were continuously harvested or grazed by herds of cattle, the non-native species were able to spread and expand across the landscape, reducing the native species diversity and ecological function.

## **Community 3.1**

### **Hayfield**

Sites in this community phase consist of forage plants that are planted and mechanically harvested. Mechanical harvesting removes much of the aboveground biomass and nutrients that feed the soil microorganisms (Franzluebbers et al. 2000; USDA-NRCS 2003). As a result, soil biology is reduced leading to decreases in nutrient uptake by plants, soil organic matter, and soil aggregation. Frequent biomass removal can also reduce the site's carbon sequestration capacity (Skinner 2008).

## **Community 3.2**

### **Continuous Pastured Grazing System**

This community phase is characterized by continuous grazing where domestic livestock graze a pasture for the entire season. Depending on stocking density, this can result in lower forage quality and productivity, weed invasions, and uneven pasture use. Continuous grazing can also increase the amount of bare ground and erosion and reduce soil organic matter, cation exchange capacity, water-holding capacity, and nutrient availability and retention (Bharati et al. 2002; Leake et al. 2004; Teague et al. 2011). Smooth brome, Kentucky bluegrass, and white clover (*Trifolium repens* L.) are common pasture species used in this phase. Their tolerance to continuous grazing has allowed these species to dominate, sometimes completely excluding the native vegetation.

## **Community 3.3**

### **Rest-Rotation Pastured Grazing System**

This community phase is characterized by rotational grazing where the pasture has been subdivided into several smaller paddocks. Through the development of a grazing plan, livestock utilize one or a few paddocks, while the remaining area is rested allowing plants to restore vigor and energy reserves, deepen root systems, develop seeds, as well as allow seedling establishment (Undersander et al. 2002; USDA-NRCS 2003). Rest-rotation pastured grazing systems include deferred rotation, rest rotation, high intensity – low frequency, and short duration methods. Vegetation is generally more diverse and can include orchardgrass (*Dactylis glomerata* L.), timothy (*Phleum pratense* L.), red clover (*Trifolium pratense* L.), and alfalfa (*Medicago sativa* L.). The addition of native prairie

species can further bolster plant diversity and, in turn, soil function. This community phase promotes numerous ecosystem benefits including increasing biodiversity, preventing soil erosion, maintaining and enhancing soil quality, sequestering atmospheric carbon, and improving water yield and quality (USDA-NRCS 2003).

### **Pathway 3.1A** **Community 3.1 to 3.2**

Mechanical harvesting is replaced with domestic livestock utilizing continuous grazing.

### **Pathway 3.1B** **Community 3.1 to 3.3**

Mechanical harvesting is replaced with domestic livestock utilizing rotational grazing.

### **Pathway 3.2A** **Community 3.2 to 3.1**

Domestic livestock are removed, and mechanical harvesting is implemented.

### **Pathway 3.2B** **Community 3.2 to 3.3**

Rotational grazing replaces continuous grazing.

### **Pathway 3.3B** **Community 3.3 to 3.1**

Domestic livestock are removed, and mechanical harvesting is implemented.

### **Pathway 3.3A** **Community 3.3 to 3.2**

Continuous grazing replaces rotational grazing.

## **State 4** **Cropland State**

The continuous use of tillage, row-crop planting, and chemicals (i.e., herbicides, fertilizers, etc.) has effectively eliminated the reference community and many of its natural ecological functions in favor of crop production. Corn and soybeans are the dominant crops for the site, and oats (*Avena L.*) and alfalfa (*Medicago sativa* L.) may be rotated periodically. These areas are likely to remain in crop production for the foreseeable future.

### **Community 4.1** **Conventional Tillage Field**

Sites in this community phase typically consist of monoculture row-cropping maintained by conventional tillage practices. They are cropped in either continuous corn or corn-soybean rotations. The frequent use of deep tillage, low crop diversity, and bare soil conditions during the non-growing season negatively impacts soil health. Under these practices, soil aggregation is reduced or destroyed, soil organic matter is reduced, erosion and runoff are increased, and infiltration is decreased, which can ultimately lead to undesirable changes in the hydrology of the watershed (Tomer et al. 2005).

### **Community 4.2** **Conservation Tillage Field**

This community phase is characterized by rotational crop production that utilizes various conservation tillage methods to promote soil health and reduce erosion. Conservation tillage methods include strip-till, ridge-till, vertical-

till, or no-till planting systems. Strip-till keeps seedbed preparation to narrow bands less than one-third the width of the row where crop residue and soil consolidation are left undisturbed in-between seedbed areas. Strip-till planting may be completed in the fall and nutrient application either occurs simultaneously or at the time of planting. Ridge-till uses specialized equipment to create ridges in the seedbed and vegetative residue is left on the surface in between the ridges. Weeds are controlled with herbicides and/or cultivation, seedbed ridges are rebuilt during cultivation, and soils are left undisturbed from harvest to planting. Vertical-till systems employ machinery that lightly tills the soil and cuts up crop residue, mixing some of the residue into the top few inches of the soil while leaving a large portion on the surface. No-till management is the most conservative, disturbing soils only at the time of planting and fertilizer application. Compared to conventional tillage systems, conservation tillage methods can improve soil ecosystem function by reducing soil erosion, increasing organic matter and water availability, improving water quality, and reducing soil compaction.

### **Community 4.3**

#### **Conservation Tillage Field/Alternative Crop Field**

This community phase applies conservation tillage methods as described above as well as adds cover crop practices. Cover crops typically include nitrogen-fixing species (e.g., legumes), small grains (e.g., rye, wheat, oats), or forage covers (e.g., turnips, radishes, rapeseed). The addition of cover crops not only adds plant diversity but also promotes soil health by reducing soil erosion, limiting nitrogen leaching, suppressing weeds, increasing soil organic matter, and improving the overall soil ecosystem. In the case of small grain cover crops, surface cover and water infiltration are increased, while forage covers can be used to graze livestock or support local wildlife. Of the three community phases for this state, this phase promotes the greatest soil sustainability and improves ecological functioning within a cropland system.

#### **Pathway 4.1A**

##### **Community 4.1 to 4.2**

Tillage operations are greatly reduced, crop rotation occurs on a regular interval, and crop residue remains on the soil surface.

#### **Pathway 4.1B**

##### **Community 4.1 to 4.3**

Tillage operations are greatly reduced or eliminated, crop rotation occurs on a regular interval, crop residue remains on the soil surface, and cover crops are planted following crop harvest.

#### **Pathway 4.2A**

##### **Community 4.2 to 4.1**

Intensive tillage is utilized, and monoculture row-cropping is established.

#### **Pathway 4.2B**

##### **Community 4.2 to 4.3**

Cover crops are implemented to minimize soil erosion.

#### **Pathway 4.3B**

##### **Community 4.3 to 4.1**

Intensive tillage is utilized, cover crop practices are abandoned, monoculture row-cropping is established, and crop rotation is reduced or eliminated.

#### **Pathway 4.3A**

##### **Community 4.3 to 4.2**

Cover crop practices are abandoned.

## **State 5**

### **Reconstructed Bedrock Woodland State**

The combination of natural and anthropogenic disturbances occurring today has resulted in numerous forest health issues, and restoration back to the historic reference condition may not be possible. Woodlands are being stressed by non-native diseases and pests, habitat fragmentation, changes in soil conditions, and overabundant deer populations on top of naturally-occurring disturbances (severe weather and native pests) (IFDC 2018). However, these habitats provide multiple ecosystem services including carbon sequestration; clean air and water; soil conservation; biodiversity support; wildlife habitat; timber, fiber, and fuel products; as well as a variety of cultural activities (e.g., hiking, camping, hunting) (Millennium Ecosystem Assessment 2005; IFDC 2018). Therefore, conservation of woodlands should still be pursued. Woodland reconstructions are an important tool for repairing natural ecological functioning and providing habitat protection for numerous species associated with Sandstone Woodlands. Therefore, ecological restoration should aim to aid the recovery of degraded, damaged, or destroyed ecosystems. A successful restoration will have the ability to structurally and functionally sustain itself, demonstrate resilience to the ranges of stress and disturbance, and create and maintain positive biotic and abiotic interactions (SER 2002). The reconstructed bedrock woodland state is the result of a long-term commitment involving a multi-step, adaptive management process.

## **Community 5.1**

### **Early Successional Reconstructed Woodland**

This community phase represents the early community assembly from woodland reconstruction. It is highly dependent on the current condition of the site based on past and current land management actions, invasive species, and proximity to land populated with non-native pests and diseases. Therefore, no two sites will have the same early successional composition. Technical forestry assistance should be sought to develop suitable conservation management plans.

## **Community 5.2**

### **Late Successional Reconstructed Woodland**

Appropriately timed management practices (e.g., prescribed fire, hazardous fuels management, forest stand improvement, continuing integrated pest management) applied to the early successional community phase can help increase the stand maturity, pushing the site into a late successional community phase over time. A late successional reconstructed woodland will have an uneven-aged canopy with a sparse, short herbaceous understory.

## **Pathway 5.1A**

### **Community 5.1 to 5.2**

Application of stand improvement practices in line with a developed management plan.

## **Pathway 5.2A**

### **Community 5.2 to 5.1**

Reconstruction experiences a setback from extreme weather event or improper timing of management actions.

## **Transition T1A**

### **State 1 to 2**

Long-term fire suppression in excess of 80 years transitions the site to the fire-suppressed forest state (2).

## **Transition T1B**

### **State 1 to 3**

Cultural treatments to enhance forage quality and yield transitions the site to the forage state (3).

## **Transition T1C**

### **State 1 to 4**

Tillage, seeding of agricultural crops, and non-selective herbicide transition this site to the cropland state (4).

### **Transition T2A**

#### **State 2 to 3**

Cultural treatments to enhance forage quality and yield transitions the site to the forage state (3).

### **Transition T2B**

#### **State 2 to 4**

Tillage, seeding of agricultural crops, and non-selective herbicide transition this site to the cropland state (4).

### **Restoration pathway R2A**

#### **State 2 to 5**

Site preparation, tree planting, invasive species control, seeding native species, and deer management transition this site to the reconstructed bedrock woodland state (5).

### **Transition T3A**

#### **State 3 to 2**

Land abandonment transitions the site to the fire-suppressed forest state (2).

### **Transition T3B**

#### **State 3 to 4**

Tillage, seeding of agricultural crops, and non-selective herbicide transition this site to the cropland state (4).

### **Restoration pathway R3A**

#### **State 3 to 5**

Site preparation, tree planting, invasive species control, and seeding native species transition this site to the reconstructed bedrock woodland state (5).

### **Transition T4A**

#### **State 4 to 2**

Land abandonment transitions the site to the fire-suppressed forest state (2).

### **Transition T4B**

#### **State 4 to 3**

Cultural treatments to enhance forage quality and yield transitions the site to the forage state (3).

### **Restoration pathway R4A**

#### **State 4 to 5**

Site preparation, tree planting, invasive species control, and seeding native species transition this site to the reconstructed bedrock woodland state (5).

### **Transition T5A**

#### **State 5 to 2**

Fire suppression and removal of active management transitions this site to the fire-suppressed forest state (2).

## **Transition T5B**

### **State 5 to 3**

Cultural treatments to enhance forage quality and yield transition the site to the forage state (3).

## **Transition T5C**

### **State 5 to 4**

Tillage, seeding of agricultural crops, and non-selective herbicide transition this site to the cropland state (4).

## **Additional community tables**

### **Inventory data references**

No field plots were available for this site. A review of the scientific literature and professional experience were used to approximate the plant communities for this provisional ecological site. Information for the state-and-transition model was obtained from the same sources. All community phases are considered provisional based on these plots and the sources identified in this ecological site description.

### **Other references**

Angel, J. No date. Climate of Illinois Narrative. Illinois State Water Survey, Prairie Research Institute, University of Illinois at Urbana-Champaign. Available at <https://www.isws.illinois.edu/statecli/General/Illinois-climate-narrative.htm>. Accessed 8 November 2018.

Barrett, S.W. 1984. Indians and fire. *Western Wildlands*. Spring: 17-20.

Bharati, L., K.-H. Lee, T.M. Isenhardt, and R.C. Schultz. 2002. Soil-water infiltration under crops, pasture, and established riparian buffer in Midwestern USA. *Agroforestry Systems* 56: 249-257.

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Conterminous United States. USDA Forest Service, General Technical Report WO-76. Washington, DC. 92 pps.

Franzluebbers, A.J., J.A. Stuedemann, H.H. Schomberg, and S.R. Wilkinson. 2000. Soil organic C and N pools under long-term pasture management in the Southern Piedmont USA. *Soil Biology and Biochemistry* 32:469-478.

Illinois Forestry Development Council (IFDC). 2018. Illinois Forest Action Plan: A Statewide Forest Resource Assessment and Strategy, Version 4.1. Illinois Forestry Development Council and Illinois Department of Natural Resources. 80 pps.

Ireland, L.C. 2000. Ice storms and forest impacts. *The Science of the Total Environment* 262:231-242.

LANDFIRE. 2009. Biophysical Setting 4913110 North-Central Interior Dry Oak Forest and Woodland. In: LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior. Washington, DC.

Leake, J., D. Johnson, D. Donnelly, G. Muckle, L. Boddy, and D. Read. 2004. Networks of power and influence: the role of mycorrhizal mycelium in controlling plant communities and agroecosystem functioning. *Canadian Journal of Botany* 82: 1016-1045.

Millennium Ecosystem Assessment. 2005. Ecosystems and Human Well-Being: Current States and Trends. World Resources Institute. Island Press, Washington, D.C. 948 pages.

NatureServe. 2018. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1 NatureServe, Arlington, VA. Available at <http://explorer.natureserve.org>. (Accessed 18 April 2019).

Peel, M.C., B.L. Finlayson, and T.A. McMahon. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11: 1633-1644.

- Peterson, C.J. 2000. Catastrophic wind damage to North American forests and the potential impact of climate change. *The Science of the Total Environment* 262: 287-311.
- Pyne, S.J., P.L. Andrews, and R.D. Laven. 1996. *Introduction to Wildland Fire*, Second Edition. John Wiley and Sons, Inc. New York, New York. 808 pps.
- Schwegman, J.E., G.B. Fell, M. Hutchinson, G. Paulson, W.M. Shepherd, and J. White. 1973. *Comprehensive Plan for the Illinois Nature Preserves System, Part 2 The Natural Divisions of Illinois*. Illinois Nature Preserves Commission, Rockford, IL. 32 pps.
- Skinner, R.H. 2008. High biomass removal limits carbon sequestration potential of mature temperate pastures. *Journal for Environmental Quality* 37: 1319-1326.
- Society for Ecological Restoration [SER] Science & Policy Working Group. 2002. *The SER Primer on Ecological Restoration*. Available at: <http://www.ser.org/>. (Accessed 28 February 2017).
- Taft, J.B., R.C. Anderson, L.R. Iverson, and W.C. Handel. 2009. Chapter 4: Vegetation ecology and change in terrestrial ecosystems. In: C.A. Taylor, J.B. Taft, and C.E. Warwick (eds.). *Canaries in the Catbird Seat: The Past, Present, and Future of Biological Resources in a Changing Environment*. Illinois Natural Heritage Survey Special Publication 30, Prairie Research Institute, University of Illinois at Urbana-Champaign. 306 pps.
- Teague, W.R., S.L. Dowhower, S.A. Baker, N. Haile, P.B. DeLaune, and D.M. Conover. 2011. Grazing management impacts on vegetation, soil biota and soil chemical, physical and hydrological properties in tall grass prairie. *Agriculture, Ecosystems and Environment* 141: 310-322.
- Tomer, M.D., D.W. Meek, and L.A. Kramer. 2005. Agricultural practices influence flow regimes of headwater streams in western Iowa. *Journal of Environmental Quality* 34:1547-1558.
- Undersander, D., B. Albert, D. Cosgrove, D. Johnson, and P. Peterson. 2002. *Pastures for Profit: A Guide to Rotational Grazing (A3529)*. University of Wisconsin-Extension and University of Minnesota Extension Service. 43 pps.
- United States Department of Agriculture – Natural Resources Conservation Service (USDA-NRCS). 2003. *National Range and Pasture Handbook, Revision 1*. Grazing Lands Technology Institute. 214 pps.
- United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2006. *Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin*. U.S. Department of Agriculture Handbook 296. 682 pps.
- U.S. Environmental Protection Agency [EPA]. 2013. *Level III and Level IV Ecoregions of the Continental United States*. Corvallis, OR, U.S. EPA, National Health and Environmental Effects Research Laboratory, map scale 1:3,000,000. Available at <http://www.epa.gov/eco-research/level-iii-andiv-ecoregions-continental-united-states>. (Accessed 1 March 2017).
- White, J. and M.H. Madany. 1978. Classification of natural communities in Illinois. In: J. White. *Illinois Natural Areas Inventory Technical Report*. Illinois Natural Areas Inventory, Department of Landscape Architecture, University of Illinois at Urbana/Champaign. 426 pps.

## Contributors

Lisa Kluesner  
 Kristine Ryan  
 Sarah Smith  
 Tiffany Justus

## Approval



Acknowledgments

This project could not have been completed without the dedication and commitment from a variety of staff members (Table 6). Team members supported the project by serving on the technical team, assisting with the development of state and community phases of the state-and-transition model, providing peer review and technical editing, and conducting quality control and quality assurance reviews.

Table 6. List of primary contributors and reviewers.

Organization	Name	Title	Location
Natural Resources Conservation Service:			
	Scott Brady,	Acting Regional Ecological Site Specialist,	Havre, MT
	Stacey Clark,	Regional Ecological Site Specialist,	St. Paul, MN
	Tonie Endres,	Senior Regional Soil Scientist,	Indianapolis, IN
	Tiffany Justus,	Soil Scientist,	Aurora, IL
	Lisa Kluesner,	Ecological Site Specialist,	Waverly, IA
	Kevin Norwood,	Soil Survey Regional Director,	Indianapolis, IN
	Kristine Ryan,	MLRA Soil Survey Leader,	Aurora, IL
	Sarah Smith,	Soil Scientist,	Aurora, IL

This site was originally approved by Chris Tecklenburg, 5/01/2020.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	01/26/2023
Approved by	Suzanne Mayne-Kinney
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:  

---
2. Presence of water flow patterns:  

---
3. Number and height of erosional pedestals or terracettes:  

---
4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not

bare ground):

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

- 
16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 

17. **Perennial plant reproductive capability:**
-