

# Ecological site F120CY008IN Loamy Skeletal Uplands

Last updated: 10/01/2024 Accessed: 05/11/2025

### **General information**

**Provisional**. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

#### **MLRA** notes

Major Land Resource Area (MLRA): 120C–Kentucky and Indiana Sandstone and Shale Hills and Valleys, Northeastern Part

120C—Kentucky and Indiana Sandstone and Shale Hills and Valleys, Northeastern Part

This area is entirely in Indiana and makes up about 1,050 square miles (2,725 square kilometers). Physiography: This area is in the Highland Rim Section of the Interior Low Plateaus Province of the Interior Plains. Both large and small tributaries of the Ohio River and the East Fork of the White River dissect the nearly level to very steep uplands in the area. The major streams and rivers have well defined valleys with broad flood plains and numerous stream terraces. The flood plains along the smaller streams are narrow. Summits are narrow and are nearly level to gently sloping. Geology: The geologic materials in this area are of Early and Middle Pennsylvanian and Late Mississippian age. The rocks consist mainly of flat-lying, interbedded sandstone, shale, coal, and siltstone with minor areas of limestone. Bedrock outcrops are common on the bluffs along the Ohio River and its major tributaries. The surficial geologic materials consist mainly of a layer of loess, typically less than 3.5 feet (1 meter) thick, on the less eroded parts of the landscape and stratified sediments of Pleistocene age along the Ohio River and its tributaries. Unconsolidated alluvium is deposited in the river valleys.

Soils: The dominant soil orders in this MLRA are Alfisols, Ultisols, and Inceptisols. The soils in the area have a mesic soil temperature regime, an udic or aquic soil moisture regime, and dominantly mixed mineralogy. They formed dominantly in loess and in residuum derived from siltstone and shale. They range from moderately deep to very deep and from somewhat poorly drained to well drained and are loamy, silty, or clayey. Fragiudults (Spickert and Tilsit series) and Hapludults (Wrays series) are the dominant soils on ridgetops and the upper parts of hills and knobs. Halpudalfs (Kurtz series), Hapludults (Gilwood and Gnawbone series), and Dystrudepts (Brownstown series) are on moderately sloping to very steep side slopes. Hapludalfs (Coolville, Rarden, Stonehead, and Wellrock series) are on the gently sloping to moderately steep lower parts of side slopes. Hapludalfs (Elkinsville series), Fragiudalfs (Pekin series), and Fragiaqualfs (Bartle series) are on stream terraces. Dystrudepts (Beanblossom, Cuba, and Steff series) and Endoaquepts (Stendal series) are on flood plains.

### **Classification relationships**

NatureServe Ecological System. Scientific Name: Southern Interior Low Plateau Dry-Mesic Oak Forest Unique Identifier: CES202.898

#### **Ecological site concept**

The Loamy Skeletal Uplands ecological site is found on hillsides and ridges. Representative soils include: Berks, Brownstone, Trevlac.

The communities described in this provisional document reflect plant communities that are likely to be found on these soils and have not been field verified. This PES describes hypotheses based on available data of many different scales and sources and has not been developed utilizing site-specific ecological field monitoring. This PES does not encompass the entire complexity or diversity of these sites. Field studies would be required to develop a

comprehensive and science-based restoration plan for these sites.

The following information is a compilation of data acquired from the Hoosier National Forest, Purdue University, Nature Conservancy, NatureServe Explorer, Vegbank.org, and USDA plants database.

Forest (phases influences by aspect and topography):

State 1, Phase 1.1. chestnut oak (Quercus prinus)- post oak (*Quercus stellata*)/ blueberry (Vaccinium spp.) – greenbrier (Smilax spp.) / poison ivy (*Toxicodendron radicans*) - sedges (Carex spp.).

Common trees in this community are white oak, chestnut oak, black oak, post oak and pignut hickory. Shrub species include members of the Vaccinium species including V. arboreum and V. pallidum.

Sapling tree species may include black oak, white oak, pignut hickory, persimmon, black cherry, eastern red cedar, sassafras and the more shade tolerant species such as white ash and sugar maple.

Subcanopy tree species will include flowering dogwood, winged elm, ironwood, serviceberry, eastern redbud, and southern blackhaw.

Understory species are variable depending on canopy cover, aspect, soil depth and micro-topography. Characteristic dry site species may include *Orbexilum pedunculatum* (Sampson's snakeroot), *Scleria oligantha* (little headed nutrush) and *Antennaria plantaginifolia* (pussytoes).

In areas of canopy openings, Andropogon scoparius (little bluestem) and Liatris spp. (blazing stars) may occur.

On the north and east slopes of these soils the community will exhibit less post oak and more white oak, chestnut oak, black oak, and hickories. Understory plants will transition to more dry-mesic species and increase in species numbers and diversity.

Topography, aspect and soil depth influence the plant communities on these sites. Sites may exhibit a gradient between these two communities but on the same mapunits.

Pasture:

State 2, Phase 2.1: *Schedonorus arundinaceus* (tall fescue). Species present would be dependent upon seeding and management.

Transitional (abandoned) field.

State 3, Phase 3.1: eastern red cedar (*Juniperus virginiana*) / greenbrier (Smilax spp.) – blueberries (Vaccinium spp.) / broomsedge bluestem (*Andropogon virginicus*)

This phase is best described as an old field habitat with a mixture of native and introduced grasses and a variety of native and non-native herbs, forbs, seedlings, and saplings. Species present would depend on adjacent seed sources and the presence of on-going disturbances such as grazing, mowing, etc.

Due to low available water and high slope percentages, cropland was not included as a major state in this ecological model.

Restoration of states to the reference community would require long-term, intensive management inputs.

### Similar sites

F120CY009IN Shallow Loamy Skeletal Uplands Shallow Loamy Skeletal Uplands

Table 1. Dominant plant species

Tree	(1) Quercus prinus
	(2) Quercus stellata

Shrub	(1) Vaccinium (2) Smilax
Herbaceous	<ul><li>(1) Toxicodendron radicans</li><li>(2) Carex</li></ul>

# **Physiographic features**

These sites are found on hillsides and ridges.

#### Table 2. Representative physiographic features

Landforms	(1) Hill (2) Ridge
Runoff class	High
Elevation	450–1,250 ft
Slope	18–75%
Water table depth	72 in
Aspect	Aspect is not a significant factor

# **Climatic features**

Climate: The average annual precipitation in most of this area is 41 to 47 inches (1,040 to 1,195 millimeters). About 60 percent of the precipitation falls during the freeze-free period. Most of the rainfall occurs as high-intensity, convective thunderstorms in summer. Snowfall is common in winter. The average annual temperature is 52 to 56 degrees F (11 to 14 degrees C). The freeze-free period averages 205 days and ranges from 190 to 220 days. The longer freeze-free periods occur along the Ohio River. (Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. United States Department of Agriculture. Natural Resources Conservation Service. United States Department of Agriculture Handbook 296. Issued 2006.)

Frost-free period (characteristic range)	169 days
Freeze-free period (characteristic range)	195 days
Precipitation total (characteristic range)	47 in
Frost-free period (actual range)	169 days
Freeze-free period (actual range)	195 days
Precipitation total (actual range)	47 in
Frost-free period (average)	169 days
Freeze-free period (average)	195 days
Precipitation total (average)	47 in

#### Table 3. Representative climatic features



Figure 1. Monthly precipitation range







Figure 3. Monthly maximum temperature range



Figure 4. Monthly average minimum and maximum temperature



Figure 5. Annual precipitation pattern



Figure 6. Annual average temperature pattern

# **Climate stations used**

• (1) BLOOMINGTON IN UNIV [USC00120784], Bloomington, IN

### Influencing water features

There are no influencing water features for this group.

# **Soil features**

Soils in this initial PES group are loamy skeletal. presetative soils include: Berks, Brownstone, Trevlac.

#### Table 4. Representative soil features

Parent material	(1) Residuum–siltstone	
Surface texture	(1) Channery silt loam (2) Very fine sandy loam	
Family particle size	(1) Loamy	
Drainage class	Well drained	
Soil depth	20–40 in	
Surface fragment cover <=3"	0%	
Surface fragment cover >3"	0–5%	
Available water capacity (0-40in)	4–6 in	
Soil reaction (1:1 water) (0-40in)	3.5–7.3	
Subsurface fragment volume <=3" (Depth not specified)	0–25%	

# **Ecological dynamics**

The communities described in this provisional document reflect plant communities that are likely to be found on these soils and have not been field verified. This PES describes hypotheses based on available data of many different scales and sources and has not been developed utilizing site-specific ecological field monitoring. This PES does not encompass the entire complexity or diversity of these sites. Field studies would be required to develop a comprehensive and science-based restoration plan for these sites.

The following information is a compilation of data acquired from the Hoosier National Forest, Purdue University, Nature Conservancy, NatureServe Explorer, Vegbank.org, and USDA plants database.

Forest (phases influences by topography and aspect):

State 1, Phase 1.1. chestnut oak (Quercus prinus)- post oak (*Quercus stellata*)/ blueberry (Vaccinium spp.) – greenbrier (Smilax spp.) / poison ivy (*Toxicodendron radicans*) - sedges (Carex spp.).

Common trees in this community are chestnut oak, black oak, white oak, post oak and pignut hickory. Shrub species include members of the Vaccinium species including V. arboreum and V. pallidum.

Sapling tree species may include black oak, white oak, pignut hickory, persimmon, black cherry, eastern red cedar, sassafras and the more shade tolerant species such as white ash and sugar maple.

Subcanopy tree species will include flowering dogwood, winged elm, ironwood, serviceberry, eastern redbud, and southern blackhaw.

Understory species are variable depending on canopy cover, aspect, soil depth and micro-topography. Characteristic dry site species may include *Orbexilum pedunculatum* (Sampson's snakeroot), *Scleria oligantha* (little headed nutrush) and *Antennaria plantaginifolia* (pussytoes).

In areas of canopy openings, Andropogon scoparius (little bluestem) and Liatris spp. (blazing stars) may occur.

On the north and east slopes of these soils the community will exhibit less post oak and more white oak, chestnut oak, black oak, and hickories. Understory plants will transition to more dry-mesic species and increase in species numbers and diversity.

Topography, aspect and soil depth influence the plant communities on these sites. Sites may exhibit a gradient between these two communities but on the same mapunits.

Pasture:

State 2, Phase 2.1: *Schedonorus arundinaceus* (tall fescue). Species present would be dependent upon seeding and management.

Transitional (abandoned) field.

State 3, Phase 3.1: eastern red cedar (*Juniperus virginiana*) / greenbrier (Smilax spp.) – blueberries (Vaccinium spp.) / broomsedge bluestem (*Andropogon virginicus*)

This phase is best described as an old field habitat with a mixture of native and introduced grasses and a variety of native and non-native herbs, forbs, seedlings, and saplings. Species present would depend on adjacent seed sources and the presence of on-going disturbances such as grazing, mowing, etc.

Due to low available water and high slope percentages, cropland was not included as a major state in this ecological model.

Restoration of states to the reference community would require long-term, intensive management inputs.



#### Provisional Ecological Site F120CY008IN – Loamy Skeletal Uplands

T1A: Pasture establishment.

T1B: Tree removal -- no post harvest management inputs. Fescue present only if on adjacent sites.

T3A: Pasture re-establishment. Inputs may include brush/tree removal, weed control, seeding, etc.

T2A: Natural transition in absence of management inputs.

R2A, R3A: Long term management inputs required to successfully restore a reference community.

Figure 7. 120C

### Inventory data references

Site Development and Testing Plan

Future work is needed, as described in a future project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

# **Other references**

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C.A. Carpenter, and W.H.McNab. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. [Map. presentation scale 1:3,500,000, colored; A.M. Sloan, cartographer] Gen. Tech. Report WO-76D. U.S. Department of Agriculture, Forest Service, Washington, DC. (https://www.fs.fed.us/research/publications/misc/73326-wo-gtr-76d-cleland2007.pdf)

Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, and K. Snow. 2003. Ecological Systems of the United States: A Working Classification of US Terrestrial Systems. NatureServe, Arlington, VA. (

https://www.natureserve.org/sites/default/files/pcom\_2003\_ecol\_systems\_us.pdf ).

NatureServe 2020. NatureServe Explorer: An Online Encyclopedia of Life [web application]. NatureServe, Arlington,

VA. ( http://explorer.natureserve.org )

Soil Survey Staff-USDA-NRCS [United States Department of Agriculture, Natural Resources Conservation Service] 2016. National Soils Information Service (NASIS Data Model Version 7.3.4) Lincoln, NE. ( https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/soils/survey/tools/?cid=nrcs142p2\_053552).

USDA-NRCS [United States Department of Agriculture, Natural Resources Conservation Service]. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

(https://www.nrcs.usda.gov/Internet/FSE\_DOCUMENTS/nrcs142p2\_051845.pdf).

USNVC [United States National Vegetation Classification]. 2019. United States National Vegetation Classification Database, V2.03. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC. ( http://usnvc.org ).

# Approval

Greg Schmidt, 10/01/2024

# Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/11/2025
Approved by	Greg Schmidt
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
- 5. Number of gullies and erosion associated with gullies:

- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values):
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth ( in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state

# 17. Perennial plant reproductive capability: