

Ecological site F121XY018KY Moderately Well Drained Fragipan Terrace

Last updated: 10/01/2024 Accessed: 05/10/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 121X-Kentucky Bluegrass

General: MLRA 121 is in Kentucky (83 percent), southern Ohio (11 percent), and southern Indiana (6 percent). It makes up about 10,680 square miles (27,670 square kilometers). The cities of Cincinnati, Ohio, and Louisville, Frankfort, and Lexington, Kentucky, are in this area.

Physiography: This area is primarily in the Lexington Plain Section of the Interior Low Plateaus Province of the Interior Plains.

Soils: The dominant soil orders in MLRA 121 are Alfisols, Inceptisols, and Mollisols. The soils in the area dominantly have a mesic soil temperature regime, an udic soil moisture regime, and mixed mineralogy. They are shallow to very deep, generally well-drained, and loamy or clayey. Hapludalfs formed in residuum on hills and ridges (Beasley, Cynthiana, Eden, Faywood, Lowell, and McAfee series) and in loess over residuum on hills and ridges (Carmel and Shelbyville series). Paleudalfs (Crider and Maury series) formed in loess or other silty sediments over residuum on hills and ridges. Fragiudalfs (Nicholson series) formed in loess over residuum on ridges. Hapludolls formed in residuum on hills and ridges (Fairmount series) and in alluvium on floodplains (Huntington series). Eutrudepts (Nolin series) formed in alluvium on flood plains.

Geology: Most of this area has an Ordovician-age limestone that has been brought to the surface in the Jessamine Dome, a high part of a much larger structure called the Cincinnati Arch. The strata of limestone have a propensity to form caves and karst topography. Younger units of thin-bedded shale, siltstone, and limestone occur at the eastern and western edges of the area.

The area has no coal-bearing units. Pleistocene-age loess deposits cover most of the bedrock units in this MLRA, and some glacial lake sediments are at the surface in the northwest corner of the area. Unconsolidated alluvium is deposited in the river valleys.

Classification relationships

Interior Highlands Mesic Hardwood Forest. (Plant Communities of the Midwest).

Deep soil Mesophytic Forest (Kentucky State Nature Preserves Commission).

Calcareous Mesophytic Forest ((Kentucky State Nature Preserves Commission).

Ecological site concept

Th Moderately Well Drained Fragipan Terrace ecological site includes soils that are moderately well drained with a fragipan. Representative soils include: Captina, Monogahela, Otwell, Otwood, Pekin, Sciotoville.

These sites are generally found on terraces within MLRA 121. Sites included in this provisional grouping are on a

number of differing aspects, slope shapes, profile positions and geomorphic components. Future field work is needed to refine this grouping into potentially multiple ESDs.

The majority of these sites in MLRA 121 are now pastureland, cropland, urban development or poor-quality stands of highly disturbed hardwoods.

State 1. (Reference): Provisional Ecological Site (PES) State 1, Phase 1.1: Plant species dominants: Quercus alba-Liriodendron tulipifera/Sanicula odorata-Agrimonia (white oak – tulip poplar / / common black snakeroot – agrimony)

Narrative: These sites are found on moderately well drained terraces with fragipans. These sites are now mostly agricultural but historically were likely dominated by a tall and closed tree canopy, a well-developed shrub layer and a diverse understory. Some sites may be subject to short-duration flooding.

Common tree species may include: Acer saccharum (sugar maple), Fagus grandifolia (beech), Fraxinus spp. (ash), Juglans nigra (black walnut), Tilia americana, and Carya cordiformis (bitternut hickory). Community composition will vary depending on site conditions, disturbances, and micro-topography. Other tree species may include: Quercus spp., Carya laciniosa, Acer nigrum, Acer negundo, Gymnocladus dioicus, Ulmus americana, Liquidambar styraciflua, Aesculus glabra, Magnolia acuminata, and Nyssa sylvatica. Shrub may include Asimina triloba, Lindera benzoin and Arundinaria gigantea.

State 2, Phase 2.1: Managed Pasture. Plant species dominant: Schedonorus arundinaceus (tall fescue)

State 2, Phase 2.2: Minimally Managed Pasture. Plant species dominants: Rosa multiflora- Rubus spp. /Schedonorus arundinaceus

State 2, Phase 2.3: Warm-season Pasture. Plant species dominants depend on landowner objectives and site characteristics, but may include: switchgrass (Panicum virgatum), little bluestem (Andropogon scoparius), indianagrass (Sorghastrum nutans), big bluestem (Andropogon gerardii), and eastern gamagrass (Tripsacum dactyloides).

State 3, Phases 3.1: Plant species dominants: Juniperus virginiana-Liriodendron tulipifera/ Rubus spp. - Rosa multiflora/ Vernonia gigantea -Schedonorus arundinaceus. (Eastern red cedar- tulip poplar/ berries-multiflora rose/ ironweed-tall fescue)

State 4, Phase 4.1: Plant species dominants: Acer saccharum – Liriodendron tulipifera / Lonicera maackii.

State 5, Phase 5.1: Plant species dominants: dependent upon seeding and management. Most common crops are corn and soybeans.

Associated sites

F121XY017KY	Somewhat Poorly Drained Fragipan Terrace
	Somewhat Poorly Drained Fragipan Terrace

Table 1. Dominant plant species

Tree	(1) Quercus alba (2) Liriodendron tulipifera
Shrub	Not specified
Herbaceous	(1) Sanicula (2) Agrimonia

Physiographic features

Site: Moderately well drained terrace soils that have a fragipan located within MLRA 121.

Geology(NASIS data): Parent Material Kind: Alluvium, Colluvium, Lacustrine deposits, Loess Parent Material Origin: sedimentary rock, shale and siltstone

Landform: terrace

Landscape: valley

Soil mapunits included in this provisional ES include: Captina, Monongahela, Otwell, Otwood, Pekin, and Sciotoville

Future field work may determine that this grouping be modified or split into multiple ESDs.

Landforms	(1) Terrace(2) Stream terrace
Runoff class	Low to very high
Flooding duration	Extremely brief (0.1 to 4 hours) to brief (2 to 7 days)
Flooding frequency	None to occasional
Ponding duration	Very brief (4 to 48 hours)
Ponding frequency	None to frequent
Elevation	410–1,250 ft
Slope	0–12%
Ponding depth	0–15 in
Water table depth	15–33 in
Aspect	Aspect is not a significant factor

Table 2. Representative physiographic features

Climatic features

These ecological sites are located in MLRA 121 and are at the northern periphery of the humid subtropical climate zone. Generally characterized by hot, humid summers and cold winter, the area has four distinct seasons. The expected annual precipitation for sites included in this ecological site description is generally in the range of 40 to 50 inches. The majority of precipitations falls during the freeze-free months, and thunderstorms with heavy rainfall are common during the spring and summer months. The freeze-free period varies somewhat based on localized topography and longitude.

MLRA climate summary: The average annual precipitation in most of this area is 41 to 45 inches. It is 45 to 52 inches along the southern edge of the area. About one-half of the precipitation falls during the growing season. Most of the rainfall occurs as high-intensity, convective thunderstorms. The annual snowfall averages about 14 inches (370 millimeters). The average annual temperature is 51 to 57 degrees F (10 to 14 degrees C). From: Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin (U.S. Department of Agriculture Handbook 296, 2006)

Frost-free period (characteristic range)	160-178 days
Freeze-free period (characteristic range)	186-199 days
Precipitation total (characteristic range)	43-45 in
Frost-free period (actual range)	155-183 days
Freeze-free period (actual range)	186-205 days

Table 3. Representative climatic features

Precipitation total (actual range)	43-45 in
Frost-free period (average)	169 days
Freeze-free period (average)	193 days
Precipitation total (average)	44 in







Figure 2. Monthly minimum temperature range



Figure 3. Monthly maximum temperature range



Figure 4. Monthly average minimum and maximum temperature



Figure 5. Annual precipitation pattern



Figure 6. Annual average temperature pattern

Climate stations used

- (1) CINCINNATI NORTHERN KY AP [USW00093814], Burlington, KY
- (2) LEXINGTON BLUEGRASS AP [USW00093820], Lexington, KY
- (3) LOUISVILLE INTL AP [USW00093821], Louisville, KY

Influencing water features

These sites may incur extremely brief to brief episodes of flooding or ponding.

Soil features

Very deep, moderately well drained soils with a fragipan located on terraces in MLRA 121. Representative soils include: Captina, Monogahela, Otwell, Otwood, Pekin, Sciotoville.

Table 4. Representative soil features

Parent material	(1) Alluvium
-----------------	--------------

Surface texture	(1) Silt loam	
Family particle size	(1) Loamy	
Drainage class	Moderately well drained	
Permeability class	Very slow	
Soil depth	18–31 in	
Surface fragment cover <=3"	0%	
Surface fragment cover >3"	0%	
Available water capacity (0-40in)	3–6 in	
Soil reaction (1:1 water) (0-40in)	3.5–8.4	
Subsurface fragment volume <=3" (Depth not specified)	0–10%	
Subsurface fragment volume >3" (Depth not specified)	0–9%	

Ecological dynamics

State and transition model





2.18 and 2.28 Development of warm season grass pasture.

Inventory data references

Site Development and Testing Plan

Future work is needed, as described in a future project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

Other references

Abrams, M.D. and G.J.Nowacki. 2008. Native Americans as active and passive promoters of mast and fruit trees in the eastern USA. The Holocene 18.7. pp. 1123-1137.

Barbour, M.G., J.H. Burk, W.D. Pitts, F.S. Gilliam, and M.W. Schwartz. 1999. Terrestrial Plant Ecology (ed. 3). Benjamin/Cummings, Inc., Menlo Park, California.

Braun, E.L. 1950. Deciduous forests of Eastern North America. Blakinston Co, Pennsylvania. Reprinted in 2001 by Blackburn Press, Caldwell, New Jersey.

Carey, Andrew B. 1983. Cavities in trees in hardwood forests. In: Davis, Jerry W.; Goodwin, Gregory A.; Ockenfeis, Richard A., technical coordinators. Snag habitat management: proceedings of the symposium; 1983 June 7-9; Flagstaff, AZ. Gen. Tech. Rep. RM-99. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 167-184. [17833]

Clark, R.C. and T.J. Weckman. 2008. Annotated catalog and atlas of Kentucky woody plants. Castanea, Occasional Paper in Eastern Botany No. 3: 1–114.

Cleland, D. T., J. A. Freeouf, J. E. Keys, Jr., G. J. Nowacki, C. A. Carpenter, and W. H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Conterminous United States. GTR-WO-76C-1. http://fsgeodata.fs.fed.us/other_resources/ecosubregions.html.

Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, K. Snow, and J. Teague. 2003. Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems. NatureServe, Arlington, Virginia.

DeGraaf, Richard M; Shigo, Alex L. 1985. Managing cavity trees for wildlife in the Northeast. Gen. Tech. Rep. NE-101. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station.

Delcourt, P.A. and H.R Delcourt. 1998. The influence of prehistoric human-set fires on oak- chestnut forests in the southern Appalachians. Castanea 63:337-345.

Evans, M., and G. Abernathy. 2008. Presettlement land cover of Kentucky. Kentucky State Nature Preserves Commission, Frankfort, Kentucky, USA.

Fenneman, N.M. 1917. Physiographic subdivisions of the United States. Proceedings of the National Academy of Sciences of the United States of America. Vol. 3(1). pp. 17 -22.

Fenneman, N.M. 1938. Physiography of Eastern United States. McGraw-Hill Book Co., New York.

Guyette, R.P. and D.C. Dey. 2000. Humans, topography, and wildland fire: the ingredients for long-term patterns in ecosystems. Pp. 28-35 in D.A. Yaussy (ed.). Proceedings of the workshop on fire, people, and the central hardwoods landscape. General Technical Report NE-274.

U.S. Department of Agriculture, Forest Service, Northeastern Forest Experimentation Station. Radnor, Pennsylvania.

Guyette, R.P., M.C. Stambaugh, D.C. Dey and R. Muzika. 2011. Predicting fire frequency with chemistry and climate. Ecosystems Published online: DOI: 10.1007/s10021-011-9512-0.

Halls, Lowell K. 1977. Southern fruit-producing woody plants used by wildlife. USDA Forest Service, General Technical Report SO-16. Southern Forest Experiment Station, New Orleans, LA.

Hardin, Kimberly I.; Evans, Keith E. 1977. Cavity nesting bird habitat in the oak-hickory forests--a review. Gen. Tech. Rep. NC-30. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 23 p.

Jennings, M.D., Faber-Langedoen, D., Loucks, O.L., Peet, R.K. and Roberts, D. 2009. Standards for associations and alliances of the U.S. National Vegetation Classification. Ecological Monographs, 79(2), 2009, pp. 173–199.

Johnson, Paul S. 1992. Oak overstory/reproduction relations in two xeric ecosystems in Michigan. Forest Ecology and Management. 48: 233-248.

Kartesz, J.T., The Biota of North America Program (BONAP). 2011. North American Plant Atlas (http://www.bonap.org/MapSwitchboard.html). Chapel Hill, N.C. [maps generated from Kartesz, J.T. 2010. Floristic Synthesis of North America, Version 1.0. Biota of North America Program (BONAP). (in press)].

Kentucky Division of Geographic Information. 2004. Kentucky 2001 Anderson level III land cover. Kentucky Division of Geographic Information, Frankfort, Kentucky, USA.

Keever, C. 1978. A study of the mixed mesophytic, western mesophytic, and oak chestnut regions of the eastern deciduous forest including a review of the vegetation and sites recommended as potential natural landmarks. Millersville State College, Pennsylvania.

Kentucky Geological Survey, Geospatial Analysis Section, Digital Mapping Team. 2004. Geologic formations. Kentucky Geological Survey, Lexington, Kentucky, USA.

Kentucky State Nature Preserves Commission. 2009. Natural communities of Kentucky. Frankfort, KY

Kentucky State Nature Preserves Commission. 2011. Kentucky natural areas inventory dataset. Frankfort, KY.

Kentucky State Nature Preserves Commission. 2012. Kentucky natural heritage database. Frankfort, KY.

Lawless, P. J., Baskin, J. M. and C. C. Baskin. 2006. Xeric Limestone Prairies of Eastern United States: Review and Synthesis. The Botanical Review 73(4): 303–325. The New York Botanical Garden.

NatureServe. 2006. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA USA

NatureServe. 2014. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available http://www.natureserve.org/explorer.

McNab, W.H. and P.E. Avers. 1994. Ecological subregions of the United States. U.S. Forest Service. Prepared in cooperation with Regional Compilers and the ECOMAP Team of the Forest Service.

McNab, W.H, D. T. Cleland, J. A. Freeouf, J. E. Keys, Jr., G. J. Nowacki, and C. A. Carpenter. 1997. Description of "Ecological Subregions: Sections of the Conterminous United States".

Noss, R. F. 1983. A regional landscape approach to maintain biodiversity. BioScience 33(11): 700-706.

Pickett, S.T.A. and P.S. White. 1985. Patch dynamics: a synthesis. In: S.T.A. Pickett and P.S. White. The ecology of natural disturbance and patch dynamics. New York: Academic Press: 371-384.

Pyne, S.J. 1982. Fire in America: a cultural history of wildland and rural fire. Princeton University Press, Princeton, New Jersey.

Quarterman, E. and R.L. Powell. 1978. Potential ecological/geological natural landmarks on the Interior Low

Plateaus. pp. 7-73. U.S. Department of the Interior, Washington, D.C. Quarterman,

Rooney, T.P., S.M. Wiegmann, D.A. Rogers and D.M. Waller. 2004. Biotic impoverishment and homogenization in unfragmented forest understory communities. Conservation Biology (in press).

Slone, T. and Wethington, T. 2001. Kentucky's Threatened and Endangered Species. Kentucky Department of Fish and Wildlife Resources, Frankfort, KY.

Smalley, Glendon W. 1990. Carya glabra (Mill.) Sweet pignut hickory. Silvics of North America. Vol. 2. Hardwoods. Agric. Handbook 654. Washington, DC: U.S. Department of Agriculture, Forest Service: 198-204.

Stambaugh, M.C. and R.P. Guyette. 2008. Predicting spatio-temporal variability in fire return intervals with a topographic roughness index. Forest Ecology and Management 254:463-473.

Stritch, L.R. 1990. Landscape-scale restoration of barrens-woodland within the oak-hickory forest mosaic. Restoration & Management Notes 8: 73-77.

Sweeney, J.M., ed. 1990. Management of dynamic ecosystems. North Cent. Sect., The Wildl. Soc., West Lafayette, Ind.

Sole, Jeffery. 1999. Distribution and Habitat of Appalachian Cottontails in Kentucky. Proceedings of the Annual Conference of Southeastern Association Fish and Wildlife Agencies 53:444-448.

United States Department of Agriculture (USDA), Natural Resources Conservation Service. Soil survey of Bullitt and Spencer Counties, KY.

United States Department of Agriculture (USDA), Natural Resources Conservation Service. Soil survey of Casey County, KY.

United States Department of Agriculture (USDA), Natural Resources Conservation Service. Soil survey of Garrard and Lincoln Counties, KY.

United States Department of Agriculture (USDA), Natural Resources Conservation Service. Soil survey of Hardin and Larue Counties, KY.

United States Department of Agriculture (USDA), Natural Resources Conservation Service. Soil survey of Jefferson County, KY.

United States Department of Agriculture (USDA), Natural Resources Conservation Service. Soil survey of Marion County, KY.

United States Department of Agriculture (USDA), Natural Resources Conservation Service. Soil survey of Nelson County, KY.

United States Department of Agriculture-Forest Service, Agriculture Handbook 654, Silvics of North America.

Sork, Victoria L.; Stacey, Peter; Averett, John E. 1983. Utilization of red oak acorns in non-bumper crop year. Oecologia. 59: 49-53.

Woods, A.J., Omernik, J.M., Martin, W.H., Pond, G.J., Andrews, W.M., Call, S.M, Comstock, J.A., and Taylor, D.D., 2002, Ecoregions of Kentucky (color poster with map, descriptive text, summary tables, and photographs): Reston, VA., U.S. Geological Survey (map scale 1:1,000,000).

Zollner, D., M.H. MacRoberts, B.R. MacRoberts, & D. Ladd. 2005. Endemic vascular plants of the Interior Highlands, U.S.A. Sida 21:1781-1791.

Contributors

Anita Arends

Approval

Greg Schmidt, 10/01/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/10/2025
Approved by	Greg Schmidt
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values):

- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 17. Perennial plant reproductive capability: