

## Ecological site F126XY005OH Wet Floodplain

Last updated: 9/27/2024  
Accessed: 05/13/2025

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 126X—Central Allegheny Plateau

This site contains poorly and very poorly drained on alluvial land deposited during the Quaternary period. ecosite is found in alluvial plain, and river valleys within hills, mountains, plateau, in MLRA 126

### Classification relationships

USDA-NRCS (USDA 2006):

Land Resource Region (LRR): N—East and Central Farming and Forest Region

Major Land Resource Area (MLRA): 126—Central Allegheny Plateau

USDA-FS (Cleland et al. 2007)

Province: 221 - Eastern Broadleaf Province

Section: 221E - Southern Unglaciaded Allegheny Plateau

Subsection: 221Ea - Pittsburgh Low Plateau

221Eb - Teays Plateau

221Ec - Ohio Valley Lowland

221Ed - East Hocking Plateau

This site crosswalks to Landfire biophysical setting (BpS) Central Interior and Appalachian Floodplain Systems.

NatureServe (2007) describes this as an aggregated system including the following standard ecological systems (for MLRA 127):

- Central Appalachian River Floodplain (CES202.608)

Component Associations (all not applicable to MLRA 127)

Association Unique ID Association Name

CEGL002014 *Fraxinus pennsylvanica* - *Ulmus* spp. - *Celtis occidentalis* Forest

CEGL002086 *Betula nigra* - *Platanus occidentalis* Forest

CEGL002191 *Cephalanthus occidentalis* / *Carex* spp. - *Lemna* spp. Southern Shrubland

CEGL002586 *Acer saccharinum* - *Fraxinus pennsylvanica* - *Ulmus americana* Forest

CEGL003896 *Platanus occidentalis* - *Betula nigra* - *Salix* (*caroliniana*, *nigra*) Woodland

CEGL004073 *Platanus occidentalis* - *Acer negundo* - *Juglans nigra* / *Asimina triloba* / *Mertensia virginica* Forest

CEGL004103 *Carex torta* Herbaceous Vegetation

CEGL004286 *Justicia americana* Herbaceous Vegetation

CEGL005014 *Fagus grandifolia* - *Quercus* spp. - *Acer rubrum* - *Juglans nigra* Forest

CEGL005033 *Acer negundo* Ruderal Floodplain Forest

CEGL005082 *Alnus serrulata* Swamp Shrubland

CEGL005449 *Calamagrostis canadensis* North-Central Wet Meadow

CEGL006001 *Acer saccharinum* - *Ulmus americana* / *Onoclea sensibilis* Forest

CEGL006036 *Platanus occidentalis* - *Fraxinus pennsylvanica* Forest

CEGL006042 *Acer saccharinum* - *Ulmus americana* / *Physocarpus opulifolius* Forest

CEGL006069 *Cephalanthus occidentalis* - *Decodon verticillatus* Shrubland

CEGL006114 *Acer saccharum* - *Fraxinus* spp. - *Tilia americana* / *Matteuccia struthiopteris* - *Ageratina altissima* Forest

CEGL006147 *Acer saccharinum* - (*Populus deltoides*) / *Matteuccia struthiopteris* - *Laportea canadensis* Forest

CEGL006176 *Acer saccharinum* / *Onoclea sensibilis* - *Boehmeria cylindrica* Forest

CEGL006184 *Betula nigra* - *Platanus occidentalis* / *Impatiens capensis* Forest

CEGL006185 *Quercus palustris* - *Acer rubrum* / *Carex grayi* - *Geum canadense* Forest

CEGL006217 *Acer saccharinum* - *Acer negundo* / *Ageratina altissima* - *Laportea canadensis* - (*Elymus virginicus*) Forest

CEGL006218 *Quercus bicolor* - *Fraxinus pennsylvanica* - (*Platanus occidentalis*) / *Chasmanthium latifolium* - *Dichanthelium clandestinum* - *Zizia aurea* Woodland

CEGL006244 *Peltandra virginica* - *Polygonum amphibium* var. *emersum* - *Carex stricta* - *Impatiens capensis* Herbaceous Vegetation

CEGL006251 *Alnus serrulata* - *Physocarpus opulifolius* Shrubland

CEGL006255 *Liriodendron tulipifera* - *Platanus occidentalis* - *Betula lenta* / *Lindera benzoin* / *Circaea lutetiana* ssp. *canadensis* Forest

CEGL006283 *Andropogon gerardii* - *Panicum virgatum* - *Baptisia australis* Herbaceous Vegetation

CEGL006305 *Salix sericea* Shrubland

CEGL006314 *Liriodendron tulipifera* - *Fraxinus* spp. / *Lindera benzoin* - *Viburnum prunifolium* / *Podophyllum peltatum* Forest

CEGL006386 *Quercus bicolor* - *Acer rubrum* / *Carpinus caroliniana* Forest

CEGL006405 *Tilia americana* - *Acer saccharum* - *Acer nigrum* / *Laportea canadensis* Forest

CEGL006445 *Carya cordiformis* - *Prunus serotina* / *Ageratina altissima* Forest

CEGL006447 *Carex trichocarpa* Herbaceous Vegetation

CEGL006459 *Acer saccharum* - *Fraxinus americana* / *Carpinus caroliniana* / *Podophyllum peltatum* Forest

CEGL006463 *Salix nigra* - *Betula nigra* / *Schoenoplectus pungens* Wooded Herbaceous Vegetation

CEGL006466 *Platanus occidentalis* / *Aesculus flava* Forest

CEGL006473 *Acer saccharum* - *Liriodendron tulipifera* / *Galium concinnum* - *Carex laxiculmis* Forest

CEGL006476 *Platanus occidentalis* - *Acer saccharinum* - *Betula nigra* - *Fraxinus pennsylvanica* / *Boehmeria cylindrica* - *Carex emoryi* Woodland

CEGL006478 *Fraxinus americana* / *Andropogon gerardii* - *Sorghastrum nutans* - *Schizachyrium scoparium* - *Pycnanthemum tenuifolium* Herbaceous Vegetation

CEGL006480 *Verbesina alternifolia* - *Elymus riparius* - *Solidago gigantea* - (*Teucrium canadense*) Herbaceous Vegetation

CEGL006481 *Eupatorium serotinum* - *Polygonum* (*lapathifolium*, *punctatum*, *pensylvanicum*) Herbaceous Vegetation

CEGL006483 *Eragrostis hypnoides* - *Ludwigia palustris* - *Lindernia dubia* - *Cyperus squarrosus* Herbaceous Vegetation

CEGL006484 *Carpinus caroliniana* - *Ilex decidua* Shrubland

CEGL006491 (*Hypericum prolificum*, *Leucothoe racemosa*) / *Schizachyrium scoparium* - *Solidago simplex* var. *racemosa* - *Ionactis linariifolius* Sparse Vegetation

CEGL006495 *Quercus rubra* - *Quercus shumardii* / *Cercis canadensis* Temporarily Flooded Forest

CEGL006497 *Quercus palustris* - *Quercus bicolor* / *Carex tribuloides* - *Carex radiata* - (*Carex squarrosa*) Forest

CEGL006518 *Prunus pumila* / *Andropogon gerardii* - *Sorghastrum nutans* Herbaceous Vegetation

CEGL006536 *Carex torta* - *Apocynum cannabinum* - *Cyperus* spp. Herbaceous Vegetation

CEGL006546 *Alnus incana* - *Viburnum recognitum* / *Calamagrostis canadensis* Shrubland

CEGL006548 *Acer* (*rubrum*, *saccharinum*) - *Fraxinus pennsylvanica* - *Ulmus americana* / *Boehmeria cylindrica* Forest

CEGL006575 *Fraxinus pennsylvanica* - (*Juglans nigra*, *Platanus occidentalis*) Forest

CEGL006595 *Spiraea alba* Shrubland

CEGL007696 *Peltandra virginica* - *Saururus cernuus* - *Boehmeria cylindrica* / *Climacium americanum* Herbaceous Vegetation

CEGL007879 *Juglans nigra* / *Verbesina alternifolia* Ruderal Forest

CEGL008405 *Liriodendron tulipifera* - *Pinus strobus* - (*Tsuga canadensis*) / *Carpinus caroliniana* / *Amphicarpaea bracteata* Forest

CEGL008449 *Pinus virginiana* - *Juniperus virginiana* var. *virginiana* - *Quercus stellata* / *Amelanchier stolonifera* / *Danthonia spicata* / *Leucobryum glaucum* Woodland

• North-Central Interior Floodplain (CES202.694)

Component Associations (all not applicable to MLRA 127)

Association Unique ID Association Name

CEGL002018 *Populus deltoides* - *Salix nigra* - *Acer saccharinum* Forest

CEGL002026 *Schoenoplectus tabernaemontani* - *Typha* spp. - (*Sparganium* spp., *Juncus* spp.) Herbaceous Vegetation

CEGL002086 *Betula nigra* - *Platanus occidentalis* Forest

CEGL002087 *Carya illinoensis* - *Celtis laevigata* Forest

CEGL002099 *Quercus michauxii* - *Quercus shumardii* - *Liquidambar styraciflua* / *Arundinaria gigantea* Forest

CEGL002101 *Quercus palustris* - (*Quercus stellata*) - *Quercus pagoda* / *Isoetes* spp. Forest

CEGL002102 *Quercus phellos* - (*Quercus lyrata*) / *Carex* spp. - *Leersia* spp. Forest

CEGL002103 *Salix nigra* Forest

CEGL002191 *Cephalanthus occidentalis* / *Carex* spp. - *Lemna* spp. Southern Shrubland

CEGL002314 River Mudflats Sparse Vegetation

CEGL002386 *Nuphar advena* - *Nymphaea odorata* Herbaceous Vegetation

CEGL002410 *Fraxinus pennsylvanica* - *Celtis* spp. - *Quercus* spp. - *Platanus occidentalis* Bottomland Forest

CEGL002417 *Quercus stellata* - *Quercus marilandica* - *Quercus falcata* / *Schizachyrium scoparium* Sand Woodland

CEGL002420 *Taxodium distichum* / *Lemna minor* Forest

CEGL002427 *Fraxinus pennsylvanica* - *Ulmus americana* - *Celtis laevigata* / *Ilex decidua* Forest

CEGL002430 *Polygonum* spp. - Mixed Forbs Herbaceous Vegetation

CEGL002431 *Acer saccharinum* - *Celtis laevigata* - *Carya illinoensis* Forest

CEGL002586 *Acer saccharinum* - *Fraxinus pennsylvanica* - *Ulmus americana* Forest

CEGL003725 *Platanus occidentalis* - *Betula nigra* / *Cornus amomum* / (*Andropogon gerardii*, *Chasmanthium latifolium*) Woodland

CEGL003836 *Arundinaria gigantea* ssp. *gigantea* Shrubland

CEGL003895 *Alnus serrulata* - *Xanthorrhiza simplicissima* Shrubland

CEGL003910 (*Diospyros virginiana*, *Platanus occidentalis*) / *Eupatorium serotinum* - *Diodia virginiana* Ruderal Herbaceous Vegetation

CEGL004049 River Valley Impoundment Mudflat Sparse Vegetation

CEGL004103 *Carex torta* Herbaceous Vegetation

CEGL004286 *Justicia americana* Herbaceous Vegetation

CEGL004420 *Acer rubrum* var. *trilobum* - *Fraxinus pennsylvanica* / *Carex crinita* - *Peltandra virginica* Forest

CEGL004544 *Quercus macrocarpa* - *Quercus shumardii* - *Carya cordiformis* / *Chasmanthium latifolium* Forest

CEGL004618 *Fraxinus pennsylvanica* - *Ulmus crassifolia* - *Celtis laevigata* Forest

CEGL004691 *Platanus occidentalis* - *Liriodendron tulipifera* - *Betula* (*alleghaniensis*, *lenta*) / *Alnus serrulata* - *Leucothoe fontanesiana* Forest

CEGL004979 *Quercus nigra* - *Quercus* (*alba*, *phellos*) Forest

CEGL005014 *Fagus grandifolia* - *Quercus* spp. - *Acer rubrum* - *Juglans nigra* Forest

CEGL005033 *Acer negundo* Ruderal Floodplain Forest

CEGL005035 *Acer saccharum* - *Carya cordiformis* / *Asimina triloba* Floodplain Forest

CEGL005057 *Quercus stellata* / (*Danthonia spicata*, *Croton willdenowii*) Woodland

CEGL005451 *Lemna* spp. Eastern North American Aquatic Vegetation

CEGL006458 *Platanus occidentalis* - *Fraxinus pennsylvanica* / *Carpinus caroliniana* / *Verbesina alternifolia* Forest

CEGL006462 *Quercus* (*rubra*, *velutina*, *alba*) / *Carpinus caroliniana* - (*Halesia tetraptera*) / *Maianthemum racemosum* Forest

CEGL006466 *Platanus occidentalis* / *Aesculus flava* Forest

CEGL006480 *Verbesina alternifolia* - *Elymus riparius* - *Solidago gigantea* - (*Teucrium canadense*) Herbaceous Vegetation

CEGL006620 *Tsuga canadensis* - *Quercus rubra* - (*Platanus occidentalis*, *Betula nigra*) / *Rhododendron maximum* / *Anemone quinquefolia* Forest

CEGL007064 *Salix caroliniana* Temporarily Flooded Ozark Shrubland

CEGL007334 *Platanus occidentalis* - *Acer saccharinum* - *Juglans nigra* - *Ulmus rubra* Forest

CEGL007339 *Platanus occidentalis* - *Fraxinus pennsylvanica* - *Quercus imbricaria* Forest

CEGL007399 *Quercus palustris* - (*Fraxinus nigra*) / *Lindera benzoin* / *Carex bromoides* Forest

CEGL007410 *Salix nigra* Large River Floodplain Forest

CEGL007702 *Liquidambar styraciflua* - *Quercus michauxii* - *Carya laciniata* / *Fagus grandifolia* - (*Aesculus flava*)

## Forest

CEGL007810 *Acer saccharinum* - *Betula nigra* / *Cephalanthus occidentalis* Forest

CEGL007879 *Juglans nigra* / *Verbesina alternifolia* Ruderal Forest

CEGL007880 *Liquidambar styraciflua* - *Liriodendron tulipifera* - (*Platanus occidentalis*) / *Carpinus caroliniana* - *Halesia tetraptera* / *Amphicarpaea bracteata* Forest

CEGL007999 *Platanus occidentalis* - *Betula nigra* - *Celtis laevigata* - *Fraxinus pennsylvanica* / *Arundinaria gigantea* Temporarily Flooded Forest

CEGL008404 *Osmunda regalis* var. *spectabilis* Seepage Scour Herbaceous Vegetation

CEGL008495 *Hypericum densiflorum* - *Alnus serrulata* / *Tripsacum dactyloides* Shrubland

CEGL008562 *Salix interior* Temporarily Flooded Shrubland

## Ecological site concept

From NatureServe (2007): This system encompasses floodplains of medium to large rivers in Atlantic drainages from southern New England to Virginia. This system can include a complex of wetland and upland vegetation on deep alluvial deposits and scoured vegetation on depositional bars and on bedrock where rivers cut through resistant geology. This complex includes floodplain forests in which *Acer saccharinum*, *Populus deltoides*, and *Platanus occidentalis* are characteristic, as well as herbaceous sloughs, shrub wetlands, riverside prairies and woodlands. Microtopography and soil texture determine how long the various habitats are inundated. Depositional and erosional features may both be present depending on the particular floodplain.

From Landfire <http://www.landfire.gov/index.php>:

This systems group encompasses large-river floodplains over much of the eastern US (NatureServe 2007), but differs from the Coastal Plain model in several ways. First it is floristically different in that it lacks cypress and tupelo except in its lowest elevation where it transitions to the Coastal Plain. Permanent standing water is lacking except in the areas closest to sea level where oxbow lakes may exist. Hydroperiods are shorter and fluvial features such as river terraces, oxbows, alluvial flats, point bars, streamside levees and other fine-scale alluvial floodplain features are abundant. NatureServe (2007) notes some will include herbaceous sloughs and shrub wetlands, particularly in abandoned channels

The substrate is primarily alluvium. The generally fertile soils are usually sandy to loamy but include local clayey and gravelly areas (NatureServe 2007).

Synonyms for this BpS and its variations include eastern riverfront forest, bottomland hardwood forest, and alluvial forest. Fire and beaver activity create a mosaic whose elements include canebrake, beaver ponds, and grass-sedge meadows in abandoned beaver clearings, as well as the swamps and bottomland hardwood forests that make up more than 95% of the cover that exists today.

Standard Ecological Systems noted by NatureServe (2007) that make up this aggregated system include:

- Central Appalachian River Floodplain (CES202.608)
- North-Central Interior Floodplain (CES202.694)
- South-Central Interior Large Floodplain (CES202.705)
- Southern Piedmont Large Floodplain Forest (CES202.324)

Issues or Problems:

Federal reservoirs have had a serious and negative effect, along with agriculture that has converted much floodplain acreage to drained agricultural land. In the remaining less altered floodplains, flood duration varies according to the river's gradient (NatureServe 2007).

## Associated sites

|             |                                                           |
|-------------|-----------------------------------------------------------|
| F126XY006OH | <b>Well Drained Floodplain</b><br>Well-drained floodplain |
|-------------|-----------------------------------------------------------|

Table 1. Dominant plant species

|       |                                  |
|-------|----------------------------------|
| Tree  | (1) <i>Platanus occidentalis</i> |
| Shrub | (1) <i>Acer negundo</i>          |

## Physiographic features

The ecological site are flood plains found in alluvial plain, and river valleys within hills, mountains, plateau, in MLRA 126

**Table 2. Representative physiographic features**

|                    |                    |
|--------------------|--------------------|
| Landforms          | (1) Flood plain    |
| Runoff class       | Very low to medium |
| Flooding frequency | Rare to frequent   |
| Ponding frequency  | Rare to frequent   |
| Elevation          | 32–422 m           |
| Slope              | 2–3%               |
| Water table depth  | 0–53 cm            |


## Climatic features

The regional climate of the unglaciated Central Allegheny Plateau is predominately a humid continental climate grading at the extreme southwestern corner a to humid temperate climate with hot summers and cool winters (Beck et al., 2018; Bailey, 2014). However, the local climate is highly influenced by the dissected terrain, where climatic variations may be greater at the local scale, e.g., cooler temperatures and shorter growing season at higher elevations and more northerly latitudes. High-intensity, convective thunderstorms are common in summer. Winter precipitation is mostly snow.


Climate change is occurring, and the resiliency of any ecological site will depend upon the direct and indirect effects upon component species and shifting atmospheric and soil conditions.

**Table 3. Representative climatic features**


|                                            |                |
|--------------------------------------------|----------------|
| Frost-free period (characteristic range)   | 139-168 days   |
| Freeze-free period (characteristic range)  | 167-198 days   |
| Precipitation total (characteristic range) | 1,016-1,092 mm |
| Frost-free period (actual range)           | 131-171 days   |
| Freeze-free period (actual range)          | 156-201 days   |
| Precipitation total (actual range)         | 991-1,143 mm   |
| Frost-free period (average)                | 151 days       |
| Freeze-free period (average)               | 184 days       |
| Precipitation total (average)              | 1,067 mm       |




**Figure 1. Monthly precipitation range**




**Figure 2. Monthly minimum temperature range**




**Figure 3. Monthly maximum temperature range**



**Figure 4. Monthly average minimum and maximum temperature**



**Figure 5. Annual precipitation pattern**



**Figure 6. Annual average temperature pattern**

## Climate stations used

- (1) WHEELING [USC00469482], Wheeling, WV
- (2) SENECAVILLE LAKE [USC00337559], Senecaville, OH
- (3) CALDWELL 3 SE [USC00331175], Caldwell, OH
- (4) HANNIBAL L&D [USC00333500], New Martinsville, OH
- (5) WAYNESBURG 1 E [USC00369367], Waynesburg, PA
- (6) WEST UNION 2 [USC00469458], West Union, WV
- (7) PARKERSBURG [USW00013867], Parkersburg, WV
- (8) WINFIELD LOCKS [USC00469683], Red House, WV
- (9) PEA RIDGE PSD [USC00466912], Huntington, WV
- (10) HUNTINGTON SEWAGE PLT [USC00464397], Kenova, WV
- (11) LOYALHANNA LAKE [USC00365212], New Alexandria, PA
- (12) PITTSBURGH INTL AP [USW00094823], Coraopolis, PA

## Influencing water features

Wet Floodplain and Drainageway ecological sites are wetlands located along river systems of varying size. River flooding and ponding are variable, ranging from rare to frequent. Flood events may be seasonal, most common in the spring and early summer.

## Wetland description

In the National Wetland Classification System (Cowardin et al., 1979), the Wet Floodplain and Drainageway ecological site may be considered in the Palustrine system, with a vegetated class, such as emergent, scrub-shrub, or forested, and modified by a non-tidal water regime ranging from seasonally flooded to seasonally-flooded/saturated.

## Soil features

The ecological site contains alluvial, poorly drained to very poorly drained soils.

**Table 4. Representative soil features**

|                                                          |                   |
|----------------------------------------------------------|-------------------|
| Parent material                                          | (1) Alluvium      |
| Surface texture                                          | (1) Silt loam     |
| Permeability class                                       | Very slow to slow |
| Depth to restrictive layer                               | 76–183 cm         |
| Surface fragment cover <=3"                              | 0%                |
| Surface fragment cover >3"                               | 0%                |
| Available water capacity<br>(Depth not specified)        | 10.16–22.86 cm    |
| Soil reaction (1:1 water)<br>(Depth not specified)       | 4.5–7.8           |
| Subsurface fragment volume <=3"<br>(Depth not specified) | 2–28%             |
| Subsurface fragment volume >3"<br>(Depth not specified)  | 1–5%              |

## Ecological dynamics

Ecological Dynamics: Information contained in this section was adapted from several sources. The information presented is representative of very complex vegetation communities. Key indicator plants, animals and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and aspect. The reference plant community is not necessarily the management goal. The species lists are representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.

State Correlation: This site will be correlated in: OH,PA,WV

From Landfire <http://www.landfire.gov/index.php>:

Most of the system is forest vegetation. The canopy is usually dominated by a mix of characteristic alluvial and bottomland species (depending on the region) such as sycamore (*Platanus occidentalis*), river birch (*Betula nigra*), box elder (*Acer negundo*), eastern cottonwood (*Populus deltoides*), sugarberry (*Celtis laevigata*), green ash (*Fraxinus pennsylvanica*), sweetgum (*Liquidambar styraciflua*), and red maple (*Acer rubrum*). NatureServe (2007) also notes characteristic trees include silver maple (*Acer saccharinum*), willows, especially black willow (*Salix nigra*) in the wettest areas, American elm (*Ulmus Americana*), swamp chestnut oak (*Quercus michauxii*), cherrybark oak (*Quercus pagoda*), and (at least in the Midwest) Bur oak (*Quercus macrocarpa*) in more well-drained areas. The particular mix of tree species will vary across the geographic range of this systems group, with some trees absent over parts of the range. Successional areas are often dominated by sweetgum (*Liquidambar styraciflua*), or tulip tree (*Liriodendron tulipifera*).

The driest and most fire sheltered sites support species such as pignut hickory (*Carya glabra*), shagbark hickory (*Carya ovata*), beech (*Fagus grandifolia*), and other fire sensitive species.

Subcanopy species included American holly (*Ilex opaca*), deciduous holly (*Ilex decidua* and *Ilex ambigua*), red mulberry (*Morus rubra*), ironwood (*Carpinus caroliniana*) and hop hornbeam (*Ostrya virginiana*). Shrubs such as spicebush (*Lindera benzoin*), beautyberry (*Callicarpa americana*), yellowroot (*Xanthorhiza simplicissima*), common buttonbush (*Cephalanthus occidentalis*), roughleaf dogwood (*Cornus drummondii*), and pawpaw (*Asimina triloba*); sedges (*Carex* spp.); and grasses including eastern bottlebrush grass (*Elymus hystrix*), Canada wildrye (*Elymus Canadensis*), and Indian woodoats (*Chasmanthium latifolium*), and false nettle (*Boehmeria cylindrica*) may be present.

Because of the particular mix and diversity of tree species will vary across the geographic range of this systems group, the list of BpS dominant and indicator species is only a rudimentary list.

State and transition model




Figure 7. State and Transition




Figure 8. Legend

**State 1  
Reference Pre-Contact Forest**

The reference state can be represented by several communities within Central Appalachian River Floodplain (CES202.608) (Natureserve, 2007). Forest overstory canopies are sycamore dominated and generally closed canopy.

## **Community 1.1**

### **Tree/Shrub/Herb**

Sycamore and River Birch Forest best captures the nature of this vegetation state. The dominate overstory canopy consist of Sycamore and river birch with minor canopy coverage of red maple. The shrub layer contains box elder, hornbeam and poison ivy. The herb layer will contain flowering forbes, ferns,grasses and sedges.

## **State 2**

### **Post Settlemet Forest**

This forest vegetation community is the result poor logging techniques (high grading).

## **Community 2.1**

### **Tree/Shrub/Herb**

Sycamore and River Birch Forest best captures the nature of this vegetation state. The dominate overstory canopy consist of Sycamore and river birch with minor canopy coverage of red maple. The shrub layer contains box elder, hornbeam and poison ivy. The herb layer will contain flowering forbes, ferns,grasses and sedges.

## **State 3**

### **Agricultural Land**

Land managed for agricultural production of crops and livestock.

## **Community 3.1**

### **Pasture (Grasses, Forbes) or Cropland (Grasses, Forbes)**

This community phase may contain a wide variety of plants depending on the level of management. In pasture circumstances that are managed tall fescue, bluegrass and white clover will dominate the vegetation canopy. Without management such as prescribed grazing, nutrient management and weed control, less desirable forage species and weeds will invade.

## **Transition T1A**

### **State 1 to 2**

The site is logged and managed for agricultural land.

## **Transition T1B**

### **State 1 to 3**

The site is logged and high graded, lees desirable species dominate succession.

## **Restoration pathway R3A**

### **State 2 to 1**

Remove undesirable species using herbicides, cutting or prescribed fire. Plant desired species if absent from the site.

## **Transition T3A**

### **State 2 to 2**

The site is logged and managed for agricultural land.

## **Transition T2A**

### **State 3 to 3**

The site agricultural management is abandoned and forest regrowth occurs through natural succession or tree

planting.

## **Additional community tables**

### **Inventory data references**

#### **Site Development and Testing Plan**

Future work is needed, as described in a future project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

### **Other references**

Anderson, D. M. 1982. Plant Communities of Ohio: A Preliminary Classification. Division of Natural Areas and Preserves, Ohio Department of Natural Resources, Columbus, OH.  
([https://www.lm.doe.gov/cercla/documents/fernald\\_docs/cat/112509.pdf](https://www.lm.doe.gov/cercla/documents/fernald_docs/cat/112509.pdf)).

Apsley, D., and B.C. McCarthy. 2004. White-tailed deer herbivory on forest regeneration following fire and thinning treatments in southern Ohio mixed oak forests. P. 461–471. In: Yaussy, D.A., D.M. Hix, R.P. Long, and P.C. Goebel (eds.) Proceedings, 14th Central Hardwood Forest Conference, Wooster, OH. 16-19 March 2004. Gen. Tech. Rep. NE-316. USDA Forest Service, Northeastern Research Station, Newtown Square, PA.

Bailey, R. 2014. Ecoregions: the ecosystem geography of the oceans and continents. 2nd ed. New York, NY: Springer-Verlag.

Beck, H.E., N.E. Zimmermann, T.R. McVicar, N. Vergopolan, A. Berg, E.F. Wood. 2018. Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Scientific Data* 5(1):1-12.

Butler, P.R., L. Iverson, F.R. Thompson, L. Brandt, S. Handler, M. Janowiak, P.D. Shannon, C. Swanston, K. Karriker, J. Bartig, and S. Connolly. 2015. Central Appalachians Forest Ecosystem Vulnerability Assessment and Synthesis: a Report From The Central Appalachians Climate Change Response Framework Project. Gen. Tech. Rep. NRS-146, US Department of Agriculture, Forest Service, Northern Research Station, Newtown Square, PA.

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C.A. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. [Map. presentation scale 1:3,500,000, colored; A.M. Sloan, cartographer] Gen. Tech. Report WO-76D. U.S. Department of Agriculture, Forest Service, Washington, DC. (<https://www.fs.fed.us/research/publications/misc/73326-wo-gtr-76d-cleland2007.pdf>)

Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, and K. Snow. 2003. Ecological Systems of the United States: A Working Classification of US Terrestrial Systems. NatureServe, Arlington, VA.  
([https://www.natureserve.org/sites/default/files/pcom\\_2003\\_ecol\\_systems\\_us.pdf](https://www.natureserve.org/sites/default/files/pcom_2003_ecol_systems_us.pdf)).

FGDC (Federal Geographic Data Committee). 2008. National Vegetation Classification Standard, Version 2. VGDC-STD-005-2008 (Version 2). FGDC Vegetation Subcommittee, Reston, Virginia.  
([https://www.fgdc.gov/standards/projects/vegetation/NVCS\\_V2\\_FINAL\\_2008-02.pdf](https://www.fgdc.gov/standards/projects/vegetation/NVCS_V2_FINAL_2008-02.pdf)).

Lafon, C.W., A.T. Naito, H.D. Grissino-Mayer, S.P. Horn, and T.A. Waldrop. 2017. Fire History of the Appalachian Region: a Review and Synthesis. Gen. Tech. Rep. SRS-219., U.S. Department of Agriculture, Forest Service, Southern Research Station, Asheville, NC.

NatureServe 2007. NatureServe Explorer: An Online Encyclopedia of Life [web application]. NatureServe, Arlington, VA. Available: <http://explorer.natureserve.org> (Accessed: April 2020).

Nowacki, G.J. and M.D. Abrams. 2008. The demise of fire and “mesophication” of forests in the eastern United States. *Bioscience* 58(2):123–138.

Ohio Division of Wildlife. 2015. Ohio's State Wildlife Action Plan. Columbus, Ohio, USA.  
([https://ohiodnr.gov/static/documents/wildlife/wildlife-management/OH\\_SWAP\\_2015.pdf](https://ohiodnr.gov/static/documents/wildlife/wildlife-management/OH_SWAP_2015.pdf)).

Soil Survey Staff-USDA-NRCS [United States Department of Agriculture, Natural Resources Conservation Service] 2016. National Soils Information Service (NASIS Data Model Version 7.3.4) Lincoln, NE. Available description: [https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/soils/survey/tools/?cid=nrcs142p2\\_053552](https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/soils/survey/tools/?cid=nrcs142p2_053552) (Accessed January 2020).

USDA-NRCS [United States Department of Agriculture, Natural Resources Conservation Service]. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.  
([https://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcs142p2\\_051845.pdf](https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_051845.pdf)).

USNVC [United States National Vegetation Classification]. 2019. United States National Vegetation Classification Database, V2.03. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC. <http://usnvc.org> (accessed April 2020).

Zimmerman, E., T. Davis, G. Podniesinski, M. Furedi, J. McPherson, S. Seymour, B. Eichelberger, N. Dewar, J. Wagner, and J. Fike (editors). 2012. Terrestrial and Palustrine Plant Communities of Pennsylvania, 2nd Edition. Pennsylvania Natural Heritage Program, Pennsylvania Department of Conservation and Natural Resources, Harrisburg, PA.

## Contributors

Jason Teets

## Approval

Greg Schmidt, 9/27/2024

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

|                                             |                   |
|---------------------------------------------|-------------------|
| Author(s)/participant(s)                    |                   |
| Contact for lead author                     |                   |
| Date                                        | 05/13/2025        |
| Approved by                                 | Greg Schmidt      |
| Approval date                               |                   |
| Composition (Indicators 10 and 12) based on | Annual Production |

## Indicators

### 1. Number and extent of rills:

---

### 2. Presence of water flow patterns:

- 
3. **Number and height of erosional pedestals or terracettes:**
- 
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
- 
5. **Number of gullies and erosion associated with gullies:**
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:**
- 
7. **Amount of litter movement (describe size and distance expected to travel):**
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**

---