

# **Ecological site F128XY503WV** **Thermic High And Intermediate Stream Terrace Alluvium**

Accessed: 05/14/2025

## **General information**

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

## **MLRA notes**

Major Land Resource Area (MLRA): 128X–Southern Appalachian Ridges and Valleys

MLRA 128, partially shown as the gray shaded area on the accompanying figure, falls into the East and Central Farming and Forest Region. This MLRA is in Tennessee (36 percent), Alabama (27 percent), Virginia (25 percent), and Georgia (12 percent). It makes up about 21,095 square miles (54,660 square kilometers).

Most of this MLRA is in the Tennessee Section of the Valley and Ridge Province of the Appalachian Highlands. The thin stringers in the western part of the area are mostly in the Cumberland Plateau Section of the Appalachian Plateaus Province of the Appalachian Highlands. A separate area of the MLRA in northern Alabama is in the Highland Rim Section of the Interior Low Plateaus Province of the Interior Plains. The western side of the area is dominantly hilly to very steep and is rougher and much steeper than the eastern side, much of which is rolling and hilly. Elevation ranges from 660 feet (200 meters) near the southern end of the area to more than 2,400 feet (730 meters) in the part of the area in the western tip of Virginia. Some isolated linear mountain ridges rise to nearly 4,920 feet (1,500 meters) above sea level.

The MLRA is highly diversified. It has many parallel ridges, narrow intervening valleys, and large areas of low, irregular hills. The bedrock in this area consists of alternating beds of limestone, dolomite, shale, and sandstone of early Paleozoic age. Ridgetops are capped with more resistant carbonate and sandstone layers, and valleys have been eroded into the less resistant shale beds. These folded and faulted layers are at the southernmost extent of the Appalachian Mountains. The narrow river valleys are filled with unconsolidated deposits of clay, silt, sand, and gravel.

## **Ecological site concept**

This site occurs primarily in alluvium on high and intermediate stream terraces associated with major rivers and streams in the Ridge and Valley, MLRA 128. This site is of large extent. Most of this area has been cleared. Where forests remain, mixed oak/pine is a reasonable classification. Longleaf pine may be important on this site toward the south in some places.

**Table 1. Dominant plant species**

Tree	(1) <i>Quercus prinus</i> (2) <i>Pinus virginiana</i>
Shrub	(1) <i>Vaccinium pallidum</i>
Herbaceous	Not specified

## **Physiographic features**

This site occurs primarily in alluvium on high and intermediate stream terraces associated with major rivers and streams in the Ridge and Valley, MLRA 128. This site is of large extent.




Figure 1. Block Diagram

Table 2. Representative physiographic features


Landforms	(1) Stream terrace (2) Terrace (3) Interfluve
Flooding duration	Brief (2 to 7 days)
Flooding frequency	None to occasional
Elevation	174–570 m
Slope	0–45%
Water table depth	46–168 cm
Aspect	Aspect is not a significant factor

### Climatic features


This area falls under the humid, mesothermal climate classification (Thornwaite, 1948). Precipitation is fairly evenly distributed throughout the year, with little or no water deficiency during any season. The average annual precipitation in most of this area is 45 to 55 inches. It increases to the south. Maximum precipitation occurs in midwinter and midsummer, and the minimum occurs in autumn. Most rainfall occurs as high-intensity, convective thunderstorms. Snowfall may occur in winter. Average annual temperatures range from 46 to 70 degrees F, increasing to the south. The freeze-free period averages 205 days and is longest in the southern part of the area and shortest at higher elevations to the north. The growing season corresponds to climate. Local climate can be variable and microclimates factor into the distribution of plants. In general, topographic features such as slope aspect, landform, steepness, and position of the ridges and valleys are important site variables in the distribution of vegetation across the landscape (Martin, 1989).

Table 3. Representative climatic features


Frost-free period (average)	192 days
Freeze-free period (average)	218 days
Precipitation total (average)	1,346 mm



**Figure 2. Monthly precipitation range**



**Figure 3. Monthly average minimum and maximum temperature**



**Figure 4. Annual precipitation pattern**

## Climate stations used

- (1) TALLADEGA [USC00018024], Talladega, AL
- (2) ROME [USC00097600], Rome, GA
- (3) KNOXVILLE MCGHEE TYSON AP [USW00013891], Alcoa, TN
- (4) MORRISTOWN RADIO WCRK [USC00406271], Morristown, TN
- (5) HUNTSVILLE INTL AP [USW00003856], Madison, AL
- (6) CHATTANOOGA LOVELL AP [USW00013882], Chattanooga, TN

## Influencing water features

The influence of water on the vegetation of this site may be significant but that is currently unknown.

## Soil features

These soils formed primarily in alluvium on high and intermediate stream terraces associated with major rivers and streams. The slopes range from 0 to 45 percent. They are very deep (more than 60 inches) to bedrock, and are moderately well or well drained. The available water capacity of these soils is moderate or high. The depth to a

seasonal high water table ranges from 1.5 to more than 6 feet. Flooding frequency ranges from none to occasional. They are not subject to ponding. The soil reaction ranges from extremely acid to neutral (pH from 3.5 to 7.3).

The soil series associated with this site are: Anniston, Armour, Cumberland, Etowah, Holston, Humphreys, Leesburg, McQueen, Nolichucky, Waynesboro

Parent Material Kind: alluvium, colluvium, residuum

Parent Material Origin: interbedded sedimentary; limestone and shale; limestone, cherty; limestone, sandstone and shale; limestone, unspecified; quartzite; sandstone and shale; sandstone, unspecified; sedimentary, unspecified

**Table 4. Representative soil features**

Parent material	(1) Alluvium–limestone and shale (2) Colluvium–cherty limestone (3) Residuum–limestone, sandstone, and shale
Surface texture	(1) Cobbly clay loam (2) Gravelly fine sandy loam (3) Stony loam
Drainage class	Moderately well drained to well drained
Permeability class	Slow to moderately rapid
Soil depth	0–152 cm
Surface fragment cover <=3"	0–5%
Surface fragment cover >3"	0–5%
Available water capacity (0-101.6cm)	9.65–19.3 cm
Soil reaction (1:1 water) (0-101.6cm)	5–5.6
Subsurface fragment volume <=3" (Depth not specified)	0–25%
Subsurface fragment volume >3" (Depth not specified)	0–15%

## Ecological dynamics

This site is now mostly cleared and used for general farm crops including small grains, tobacco, cotton, hay, and pasture. Originally, forests were mixed hardwoods and pines, chiefly oak species, hickory, dogwood, yellow-poplar, elm, beech, and shortleaf, loblolly, and Virginia pines. Steep slopes are in mixed hardwood and pine forests. Where pastures have reverted to forests, dominant tree species consist of mostly pine.

VegBank lists three usable NatureServe plot classifying it as *Pinus virginiana* - *Pinus* (rigida, echinata) - (*Quercus prinus*) / *Vaccinium pallidum* Forest, *Pinus palustris* - *Pinus echinata* - (*Pinus virginiana*) / *Quercus marilandica* - (*Quercus prinus*) / *Vaccinium pallidum* Woodland, and *Quercus alba* - *Quercus rubra* - *Quercus muehlenbergii* / *Cercis canadensis* Forest. The oak dominated community and the Virginia pine dominated community are on eroded phases of Anniston and Allen gravelly loams. The longleaf pine dominated community is mapped Anniston and Allen stony loams, 10 - 25% slope. Longleaf may be a good reference community for this PES in the southern extent of the PES and should be investigated for possible separation based on prominent vegetation in the future. Some of the differences in plant community may be due to microclimate. Before a true reference community can be determined, field investigation is needed. Mixed oak/pine is a generalized classification for the time being.

## State and transition model

### Other references

DeSelm, Hal. 1989 – 2009. Natural Terrestrial Vegetation of Tennessee (Vegetation Plot Data). Unpublished raw

data. <http://treeimprovement.utk.edu/DeSelmData/DataDSC.htm>

Griffith, G.E., Omernik, J.M., and Azevedo, S.H., 1997, Ecoregions of Tennessee: Corvallis, Oregon, U.S. Environmental Protection Agency EPA/600R-97/022, 51 p.

Martin, William H. 1989. Forest patterns in the Great Valley of Tennessee. Journal of the Tennessee Academy of Science 64(3): 137 – 143.

Natureserve. 2016. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.0. NatureServe, Arlington, VA. U.S.A. Available <http://explorer.natureserve.org>.

Thornthwaite, Charles W. 1948. An approach toward a rational classification of climate. Geographical Review 38(1): 55-94.

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

Vegetation plot data. 2015. Retrieved from: <http://vegbank.org/vegbank/index.jsp>

Vegetation community description. 2015. Retrieved from: <http://www.basic.ncsu.edu/segap/>

## Contributors

Belinda Esham Ferro

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

### 1. Number and extent of rills:

---

### 2. Presence of water flow patterns:

---

### 3. Number and height of erosional pedestals or terracettes:

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
- 
5. **Number of gullies and erosion associated with gullies:**
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:**
- 
7. **Amount of litter movement (describe size and distance expected to travel):**
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

- 
16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 

17. **Perennial plant reproductive capability:**
-