

## Ecological site F131DY001AR Loamy Flood Plain

Last updated: 9/22/2023  
Accessed: 05/11/2025

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

### MLRA notes

Major Land Resource Area (MLRA): 131D–Southern Mississippi River Terraces

Major Land Resource Area (MLRA) 131D, Southern Mississippi River Terraces, is located in Arkansas (88 percent) and Louisiana (12 percent). It is in two separate areas that together make up about 2,945 square miles. The towns of Lonoke and Stuttgart, Arkansas, are in the northeastern part of the MLRA, which is called the Grand Prairie. Bastrop, Louisiana, is in the southwestern part of the MLRA.

### Classification relationships

USDA-Natural Resources Conservation Service, 2006.  
-Major Land Resource Area (MLRA) 131D

### Ecological site concept

The Loamy Floodplain ecological site has loamy-textured, very deep, somewhat poorly to well drained soils that are prone to flooding. Sites can flood for prolonged periods, especially during the winter and early spring.

### Associated sites

F131DY002AR	<b>Clayey Flood Plain</b> Landforms are similar, but have clayey-textured soils.
F131DY005AR	<b>Loamy Prairie Terrace</b> Sites have similar textures, but are found on a higher terrace landform.

## Similar sites

F131CY002LA	<b>Loamy Flood Plain</b> Sites are similar, but in a different MLRA.
F131BY003AR	<b>Loamy Flood Plain</b> Sites are similar, but in a different MLRA.

**Table 1. Dominant plant species**

Tree	(1) <i>Quercus nigra</i> (2) <i>Quercus phellos</i>
Shrub	Not specified
Herbaceous	Not specified

## Physiographic features

These soils are on nearly level to gently sloping natural levees on floodplains. Slope is dominantly 0 to 3, but is typically less than 1 percent.

**Table 2. Representative physiographic features**

Landforms	(1) Alluvial plain > Flood plain (2) Alluvial plain > Natural levee
Runoff class	Negligible to low
Flooding duration	Brief (2 to 7 days)
Flooding frequency	None to frequent
Ponding frequency	None
Elevation	43–76 m
Slope	0–3%
Water table depth	0–203 cm
Aspect	Aspect is not a significant factor

## Climatic features

The average annual precipitation is 53 inches, which increases from north to south. Most of the rainfall occurs as frontal storms in spring and early summer. Some high-intensity, convective thunderstorms occur in summer. The average seasonal snowfall is five inches. Temperatures range from highs in the low 90's during the summer to lows in the low 30's during the winter. The frost-free period averages 221 days, while the freeze-free period averages 246 days.

**Table 3. Representative climatic features**

Frost-free period (average)	221 days
Freeze-free period (average)	246 days
Precipitation total (average)	1,346 mm

## Climate stations used

- (1) ARKANSAS POST [USC00030240], Gillett, AR
- (2) DES ARC [USC00031968], Des Arc, AR
- (3) CROSSETT 2 SSE [USC00031730], Crossett, AR
- (4) SAINT CHARLES [USC00036376], Clarendon, AR
- (5) STUTTGART 9 ESE [USC00036920], Stuttgart, AR

## Influencing water features

The ecological site is on floodplains and is prone to annual flooding.

## Wetland description

Most of the soils correlated to this site are considered hydric, but onsite delineations are needed to confirm wetland status according to the United State Army Corps of Engineers.

## Soil features

The soils consist of very deep, well drained, moderately permeable soils that formed in loamy alluvial sediments or mixed alluvial and loess deposits over loamy Arkansas River alluvium. The Goodwill, Idee and Tichnor series are correlated to this site. The taxonomic classification for the Goodwill series is fine-silty, mixed, active, thermic Ultic Hapludalfs and the taxonomic classification for the Idee and Tichnor series is fine-silty, mixed, active, thermic Typic Endoaqualfs.

**Table 4. Representative soil features**

Parent material	(1) Alluvium–igneous and sedimentary rock
Surface texture	(1) Silt loam
Family particle size	(1) Loamy
Drainage class	Very poorly drained to well drained
Permeability class	Slow to very slow
Soil depth	203 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	12.7–17.78 cm
Calcium carbonate equivalent (0-101.6cm)	0%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	4.5–6.5
Subsurface fragment volume <=3" (Depth not specified)	0%
Subsurface fragment volume >3" (Depth not specified)	0%

## Ecological dynamics

The information in this ecological site description (ESD), including the state-and-transition model (STM), was

developed using archeological and historical data, professional experience, and scientific studies. The information is representative of a complex set of plant communities. Not all scenarios or plants are included. Key indicator plants, animals, and ecological processes are described to inform land management decisions.

**Introduction** - The Southern Mississippi River Terrace (MLRA 131D) region is on smooth terraces and floodplains along the Mississippi River and its major tributaries south of its confluence with the Ohio River. The geologic material in the region consists of very thick deposits of sandy to clayey alluvium of Pleistocene to Holocene age. This material was deposited by the rivers with local relief is typically less than 15 feet and elevation ranging from 50 to 250 feet.

**Geology** - Bedrock in this area consists of Tertiary and Cretaceous sands formed as beach deposits during the retreat of the Cretaceous ocean from the midsection of the United States. Alluvial deposits from flooding and lateral migration of the rivers crossing this area typically lie above the bedrock. These sediments form Pleistocene-age alluvial terraces. Silty alluvium underlies most of the area. Clayey sediments are in old channel scars. The Pleistocene terraces are part of the Prairie Terrace complex. A minor portion of the area is in the Deweyville and Montgomery terrace formation. These terraces have a base of red alluvium capped by one to several feet of brownish alluvium.

**Biological Resources** - This area as a whole, supports hardwoods and pines. The Grand Prairie originally supported tall prairie grasses interlaced with hardwood timber. Cherrybark oak (*Quercus pagoda*) and Shumard oak (*Quercus shumardii*) are widely distributed. Tuliptree (*Liriodendron tulipifera*), white ash (*Fraxinus americana*), eastern cottonwood (*Populus deltoides*), and black walnut (*Juglans nigra*) are important species on the flood plains. Loblolly pine (*Pinus taeda*) and shortleaf pine (*Pinus echinata*) are on a wide variety of sites, mainly the eroded soils on uplands and ridges. Other hardwood species that commonly grow in this area are white oak (*Quercus alba*), basswood (*Tilia* sp.), sweetgum (*Liquidambar styraciflua*), water oak (*Quercus nigra*), American elm (*Ulmus americana*), blackgum (*Nyssa sylvatica*), sycamore (*Plantanus occidentalis*), sassafras (*Sassafras albidum*), southern red oak (*Quercus falcata*), chinkapin oak (*Quercus muehlenbergii*), American beech (*Fagus grandifolia*), and hickory (*Carya* sp.).

**Land Use** - Land use varies throughout the MLRA consisting of 42 percent cropland, 4 percent grassland, 47 percent forest, 3 percent urban development, 3 percent water, and 1 percent other. Scattered tracts of forests and farms make up nearly all of this area. Rice, soybeans, and wheat are the main crops. In most areas furrow or flood irrigation is used throughout the growing season. Hardwood timber is harvested on some forested wetlands, and most forested areas are managed for wildlife. Bait fish are produced commercially in ponds that are contained by levees. The area is in a major migratory flight path and hunting waterfowl is a popular recreation activity.

**Conservation** - The major soil resource concerns are management of soil moisture, erosion control, and maintenance of the content of organic matter and productivity of the soils. Depletion of ground water through excessive pumping is a major concern in the Grand Prairie area. Conservation practices on cropland generally include nutrient management, crop residue management, and alternative tillage systems, especially no-till systems that reduce the need for tillage. In many areas land leveling or shaping optimizes the control of surface water. Other major cropland management practices are control of competing vegetation and insects through aerial or ground spraying of herbicides and insecticides and fertility management programs that make use of chemical fertilizers.

## **State and transition model**




Figure 6. STM

## State 1 Forest

The overall state has a high overstory cover of bottomland hardwood species. The dominant overstory species are water oak and willow oak (*Quercus phellos*). Flooding is common, varying from brief durations to long durations depending on micro-relief, size of precipitation events, and current saturation of the soil. This microrelief has significant impact on the plant communities as well. Sites receiving more water will have more hydrophytic vegetation than sites that are higher, or near the peripheries. The most common disturbance is treefall due to windthrow. The rooting systems in the bottoms are oftentimes shallow. In combination with some mortality due to

prolonged flooding, downed trees and upright snags are common. A canopy-clearing disturbance, such as hurricanes or tornadoes, can be inhabited by light-seeded species. If advanced oak reproduction is present at time of disturbance the stand will retain its oak dominance. Oaks will sprout, grow, die-back, and regrow for many years. Otherwise, green ash and sweetgum will colonize the canopy due to their rapid growth and ability to grow into the crown early.

## **Community 1.1**

### **Water Oak/Willow Oak Forest**

Besides the co-dominants water oak and willow oak, associate species may include: sweetgum, Nuttall oak (*Quercus texana*), overcup oak (*Quercus lyrata*), hackberry (*Celtis laevigata*), ash, and red maple (*Acer rubrum*). Understory shrub species include: swamp dogwood (*Cornus foemina*), green hawthorn (*Crataegus viridis*), and possumhaw (*Ilex decidua*). Common understory herbaceous species include: peppervine (*Ampelopsis arborea*), trumpet-creeper (*Campsis radicans*), and poison ivy (*Toxicodendron radicans*). Sedges (*Carex* sp.) and other herbaceous vegetation adapted to seasonally prolonged flooding inhabit the forest understory.

## **State 2**

### **Invasion**

Chinese tallow (*Triadica sebifera*) is an undesired, invasive species brought to the United States in 1776 (Randall & Marinelli, 1996). Rapid expansion along the gulf coastal states has allowed the species to invade many ecosystems and consequently reduce diversity. Tallow trees are known to cause gastrointestinal upset, contact dermatitis, and toxicity in livestock and humans. Mechanical and chemicals options exist as a means to control the trees.

## **Community 2.1**

### **Exotic Thicket**

Chinese tallow invade the ecological site via flooding events as nearby waterways transport seeds. Once settled, the seeds produce saplings viable to reproduce seeds in as little as three years. The rapid establishment immediately blocks sunlight to understory species and reduces diversity. Unabated growth quickly allows the saplings to grow into the overstory, thus changing the ecological state entirely. Reductions in size and number of all vegetative species are seen in all canopy tiers.

## **State 3**

### **Pasture and Cropland**

The Pasture and Cropland State is a result of conversion activities. The landowner has maximized agriculture production by planting a monoculture of introduced grass species or agricultural row crops.

## **Community 3.1**

### **Planted Pasture and Row Crop**

Typical perennial warm-season grasses include Bermudagrass, bahiagrass, dallisgrass, and Johnsongrass. Spring and fall forages may include legumes such as clover. The grasses are grown for livestock production through direct grazing or baling hay for later use. Agricultural row crops are grown for food and fiber production. Typical crops include cotton, soybeans, milo, corn, rice, and sugarcane. Many farmers use herbicides to reduce unwanted plant competition which yields a plant community unrepresentative of State 1 or subsequent vegetative states.

## **Transition T1A**

### **State 1 to 2**

The transition from State 1 to State 2 is a result of occupancy by invasive species or other noxious weeds. Invasive plants outcompete, and eventually choke out, all other native species.

## **Transition T1B**

### **State 1 to 3**

The transition is due to the land manager maximizing agricultural production. If present, merchantable timber is harvested by clearcut, then the site is prepared and planted to either a tame grass or row crop.

## **Restoration pathway R2A**

### **State 2 to 1**

The driver for restoration is control of Chinese tallow. Although an option, mechanical removal of the trees is difficult because they readily regrow from roots and seeds. Several chemicals methods are available, including glyphosate for cut-stump treatments, triclopyr for cut-stump and foliar treatments, imazamox for broad spectrum application, and imazapyr as a foliar spray. Many aquatic herbicides have water use restrictions and can potentially kill hardwoods, so labels and restrictions should be read carefully prior to application.

## **Transition T2A**

### **State 2 to 3**

The transition is due to the land manager maximizing agricultural production. Merchantable timber is harvested by clearcut, then the site is prepared and planted to either a tame grass or row crop.

## **Restoration pathway R3A**

### **State 3 to 1**

This restoration pathway may be accomplished by restoring bottomland hardwoods. Restoration efforts for bottomland hardwood forests have proven difficult and much research has been done on these ecosystems. Many times restoring the function of the ecosystem is the most difficult obstacle. Evapotranspiration and hydrioperiod are closely linked and may never fully be restored until a forested condition exists again (Stanturf et al., 2001). Local tree availability may limit the possibilities of species composition. Careful planning of available species, site design, and further management actions should be conversed with a knowledgeable restoration source. With this in mind, oftentimes late summer and early fall are the best times to begin due to possibly wet conditions during the late fall to early spring. Many detailed guides have been written to assist with restoration, and suggested readings include, "A Guide to Bottomland Hardwood Restoration" (Allen et al., 2001).

## **Transition T3A**

### **State 3 to 2**

The transition is due to the land manager not managing the invasion of exotic weeds. Without proper management, the crops and pastures can become an exotic thicket of invasive species that becomes increasingly harder to control.

## **Additional community tables**

### **Inventory data references**

This site description was developed as part of the provisional ecological site initiative using historic soil survey manuscripts, available range site descriptions, and low intensity field sampling.

### **Other references**

Allen, J. A., B. D. Keeland, J. A. Stanturf, and A. F. Kennedy Jr. 2001. A guide to bottomland hardwood restoration. Technical report, USGS/BRD/ITR-2000-0011.

Louisiana Natural Heritage Program. 2009. The Natural Communities of Louisiana. Baton Rouge, LA, U.S.A. Data current as of August 2009.

NatureServe. 2013. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA, U.S.A. Data current as of 12 July 2013.

Randall, J. M., and J. Marinelli. 1996. Invasive plants: weeds of the global garden. Volume 149. Brooklyn Botanic

Garden, Brooklyn, NY.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Survey Geographic (SSURGO) Database.

Stanturf, J. A., S. H. Schoenholtz, C. J. Schweitzer, and J. P. Shepard. 2001. Achieving restoration success: Myths in bottomland hardwood forests. *Restoration Ecology*, 9:189-200.

Stringham, T. K., W. C. Krueger, and P. L. Shaver. 2003. State and transition modeling: An ecological process approach. *Journal of Range Management* 56:106-113.

U.S. Army Corps of Engineers. 2010. Regional supplement to the Corps of Engineers Wetland Delineation Manual: Atlantic and Gulf Coastal Plain Region (Version 2.0). U.S. Army Corps of Engineers, Engineer Research and Development Center, Environmental Laboratory ERDC/EL TR-10-20.

USDA-NRCS Ag Handbook 296 (2006).

## Contributors

Marc Bordelon  
Tyson Hart

## Approval

Bryan Christensen, 9/22/2023

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	11/15/2021
Approved by	Bryan Christensen
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

### 1. Number and extent of rills:

---

### 2. Presence of water flow patterns:

---

### 3. Number and height of erosional pedestals or terracettes:

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
- 
5. **Number of gullies and erosion associated with gullies:**
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:**
- 
7. **Amount of litter movement (describe size and distance expected to travel):**
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

- 
16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 

17. **Perennial plant reproductive capability:**
-