

# **Ecological site F143XY220ME**

## **Semi-Acidic Peat Wetland Complex**

Last updated: 10/07/2024  
Accessed: 05/10/2025

---

### **General information**

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### **MLRA notes**

Major Land Resource Area (MLRA): 143X–Northeastern Mountains

MLRA 143 is in Maine (51 percent), New York (27 percent), Vermont (13 percent), New Hampshire (7 percent), and Massachusetts (2 percent). It makes up about 34,409 square miles (89,118 square kilometers). The MLRA consists of rolling hills and mountains covered by Wisconsin till. It is in three parts separated by other MLRAs. The western part is in New York (primarily the Adirondack Mountains). The central part is mainly in the Green Mountains in Vermont and the Berkshires in Massachusetts. The eastern part is in New Hampshire and most of northern Maine. The MLRA is used mainly for forestry and recreational purposes. The western part of MLRA 143 in the Adirondack Mountains has a distinct boundary with the physiographical dissimilar Saint Lawrence-Champlain Plain. The middle part that encompasses the Green Mountains has a diffuse boundary as it blends into the northern part of the New England and Eastern New York Uplands on the foothills of the Green Mountains. The southern boundary of the easternmost part of MLRA 143 has the same diffuse boundary. The northern boundary of the MLRA is the Canadian border.

The westernmost part of this MLRA is primarily in the Adirondack province of the Appalachian Highlands. A small area in the southern end of the western part is in the Mohawk section of the Appalachian Plateaus province of the same division. The easternmost part, primarily in northern Maine, is in the New England Upland section of the New England province of the Appalachian Highlands. The southwestern half of this part is in the White Mountain section of the same province and division, and the middle part of the MLRA is in the Green Mountain section. The mountains and foothills in this MLRA are commonly rounded. They are underlain by bedrock and typically covered with thin deposits of till. The more rugged mountain areas are separated by high-gradient streams coursing through steep areas of colluvium or talus-laden valleys. Many glacially broadened valleys are filled with glacial outwash and have numerous swamps and lakes. The mountains and foothills are moderately steep to very steep, and the valleys are nearly level to sloping.

As the northernmost MLRA in the region with the coldest temperatures and shortest growing season, the Northeastern Mountains have less overall tree diversity, fewer pine and oak trees, and more abundant spruce and fir trees than neighboring MLRAs. The variability in microtopography on this site results in a patchy mosaic of plant communities. Silver maple is the most common overstory species, with diverse grasses and herbs indicating differences in soil wetness throughout the site due to slight variability in elevation above the water table. This site is subject to ice scour and flooding, but the most extensive disturbance is cultivation. These broad, flat landforms are nutrient rich with high water-holding capacity. These factors along with their adjacency to rivers made them ideal farming locations for early settlers, much of which continues today. The effects of altered flow regimes from modern dams may also be significant but require further study.

### **Classification relationships**

This site occurs in Ecological Site Group 2 (Open Wetlands) of MLRA 143 (The Northeastern Mountains), in the Northeastern Forage and Forest Region (Land Resource Region R).

The Northeastern Forage and Forest LRR includes all of Maine, New Hampshire, Vermont, Rhode Island, and Connecticut, as well as large portions of Massachusetts, New York, New Jersey, Pennsylvania, and Ohio. Its southern boundary marks the extent of the Wisconsin ice sheet, which engulfed the entire LRR as recently as 10,000 to 15,000 years ago. Erosional and depositional processes associated with glaciation created many of the topographic patterns that distinguish MLRAs within the Northeastern region. Harder granitic and metamorphic bedrock to the north were more resistant to glacial erosion, resulting in the relatively nutrient poor mountains of MLRA 143; whereas nutrient-rich sedimentary bedrock of MLRAs 139, 140, and 146 resulted in relatively flat, fertile landscapes ideal for cultivation. Other areas were depressed below sea-level by the sheer mass of the glacier, resulting in pockets of marine sediments which distinguish MLRAs 142, 144A, 144B, and 145.

Precipitation is sufficient to support productive forestland throughout the Northeastern region. Still, a latitudinal temperature gradient from mesic to frigid soil temperatures results in a general transition from central hardwoods and pine in the southern MLRAs to northern hardwoods and spruce-fir forests farther north (no true boreal forests exist in the region). Elevations are generally low throughout the Northeastern region, with the exception of MLRA 143 which has many high mountain ecosystems with cryic temperature regimes and alpine vegetation above the tree line.

## Ecological site concept

This site occurs in flat, low-lying areas characterized by very poorly-drained, semi-acidic peat soils and bog vegetation. Soil pH is typically between 4.5 and 6.0 throughout, allowing for more overall species diversity than true acid bogs, but also lacking many true acid bog indicator species. It is dominated by sphagnum moss and heath shrubs, and supports other common bog species such as cotton grass in lower quantities. This site may also support low cover of black spruce and larch trees in some areas.

This ecological site is resistant to major disturbances except for small scale hydrologic alterations that may create small patches of drained or ponded peatland. This ecological resistance can be attributed to the ability of these bogs to respond to large fluctuations in water, as well as general resistance to fire, insects, disease, construction, land management, etc.. Further study is needed to identify alternative states for this site.

## Associated sites

F143XY230ME	<b>Acidic Peat Wetland Complex</b> The Semi-acidic Peat Wetland Complex may grade into the Acidic Peat Wetland Complex, usually with the latter being toward the center of the bog and grading outward to be less acidic toward the surrounding forested land.
-------------	---

## Similar sites

F143XY230ME	<b>Acidic Peat Wetland Complex</b> The Acidic Peat Wetland Complex has pH less than 4.5 throughout the profile, compared to pH of greater than 4.5 in at least part of the profile for the Semi-acidic Peat Wetland Complex site. The lower pH result in the most acid bog vegetation indicators, such as pitcher plant and sundew.
F143XY210ME	<b>Marsh Wetland Complex</b> The Marsh Wetland Complex occurs in a similar landscape position, but has more nutrient and oxygen-rich soil water conditions, resulting in the decomposition of organic matter into muck, rather than the peat accumulation characteristic of the Semi-acidic Peat Wetland Complex.

Table 1. Dominant plant species

Tree	(1) <i>Acer saccharinum</i> (2) <i>Abies balsamea</i>
------	--

Shrub	Not specified
Herbaceous	Not specified

## Physiographic features

This site occurs in flat, wet, low-lying areas where large amounts of water collect throughout the area. Water ponds on the surface for significant periods of time, and the lack of soil oxygen and nutrients impeded the decomposition of organic matter over time.

**Table 2. Representative physiographic features**


Landforms	(1) Bog (2) Marsh (3) Swamp
Flooding frequency	None
Ponding duration	Long (7 to 30 days) to very long (more than 30 days)
Ponding frequency	Occasional to frequent
Elevation	10–2,100 ft
Slope	0–2%
Aspect	Aspect is not a significant factor

## Climatic features


As the northernmost MLRA in the region, this site experiences frigid and snowy winters, warm rainy summers, and a relatively short five to six month growing season. Precipitation is considerably constant from month to month; however, areas of higher elevations may receive up to double the annual precipitation of the lower elevations and have a three to four month growing season with extremely cold winters.

**Table 3. Representative climatic features**


Frost-free period (characteristic range)	106-129 days
Freeze-free period (characteristic range)	139-159 days
Precipitation total (characteristic range)	43-49 in
Frost-free period (actual range)	101-134 days
Freeze-free period (actual range)	134-164 days
Precipitation total (actual range)	41-50 in
Frost-free period (average)	118 days
Freeze-free period (average)	149 days
Precipitation total (average)	46 in




**Figure 1. Monthly precipitation range**



**Figure 2. Monthly minimum temperature range**



**Figure 3. Monthly maximum temperature range**



**Figure 4. Monthly average minimum and maximum temperature**




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

### Climate stations used

- (1) INDIAN LAKE 2SW [USC00304102], Indian Lake, NY
- (2) WESLEY [USC00179294], Machias, ME

### Influencing water features

Large amounts of water saturate the soils of this site throughout much of the year, limiting tree growth and favoring sphagnum moss, heath shrubs, and other common bog vegetation.

### Soil features

Soils of this site are very poorly-drained peat. These soils are very deep, usually with much greater than 40 inches of organic deposits over mineral soil. They act as a sponge with exceedingly high water-holding capacity. Soil pH is expected to be between 4.5 and 6.0, though these peat conditions may exist outside this range. The soils of this site are characterized by not only their semi-acidic pH, but also by the lack of dissolved oxygen in the water source, which inhibits organic matter decomposition, resulting in peat accumulation.

Table 4. Representative soil features

Parent material	(1) Herbaceous organic material (2) Woody organic material
Surface texture	(1) Peat
Drainage class	Very poorly drained
Permeability class	Slow
Soil depth	60 in
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%

Available water capacity (14in)	Not specified
Soil reaction (1:1 water) (0in)	Not specified
Subsurface fragment volume <=3" (10in)	Not specified
Subsurface fragment volume >3" (0in)	Not specified

## Ecological dynamics

[Caveat: The vegetation information contained in this section and is only provisional, based on concepts, and future projects support validation through field work. \*] The vegetation groupings described in this section are based on the terrestrial ecological system classification and vegetation associations developed by NatureServe (Comer et al., 2003) and localized associations provided by the New York Natural Heritage Program (Edinger et al., 2014), Maine Natural Areas Program (Gawler and Cutko, 2010), New Hampshire Natural Heritage Program (Sperduto and Nichols, 2011), and Massachusetts Division of Fisheries and Wildlife (Swain, 2020).

The vegetation of this site is dominated by sphagnum moss and heath shrubs. It also supports other common bog species such as cotton grass in lower quantities. This site may sometimes support low cover of black spruce and larch trees, though the reason for tree presence or absence is poorly-understood.

This ecological site is resistant to major disturbances except for small scale hydrologic alterations that may create small patches of drained or ponded peatland (such as near a culvert). This ecological resistance can be attributed to the ability of these bogs to respond to large fluctuations in water, as the peat acts like a sponge, expanding and contracting with the water supply. There is also a general resistance to fire, insects, disease, construction, land management, etc. due to the wet nature and particular species on the site. Further study is needed to identify alternative states for this site.

## State and transition model

# F143XY220ME – Semi-acidic Peat Wetland Complex

## 1. Reference State/Current Potential

### 1.1 Sphagnum and Heath Shrubs dominant

Sphagnum moss cover greater than 95%. Heath shrubs dominate the vascular plants, with cinnamon fern, cotton grass, three-seeded sedge and a diversity of other species possible.

Figure 7. STM

### Inventory data references

Future work is needed, as described in a future project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

### Other references

Comer, P., D. Faber-Langendoen, R. Evans, S. Grawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schultz, K. Snow, and J. Teague. 2003. Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems. NatureServe, Arlington, Virginia

Edinger, G. J., D. J. Evans, S. Gebauer, T. G. Howard, D. M. Hunt, and A. M. Olivero (editors). 2014. Ecological Communities of New York State. Second Edition. A revised and expanded edition of Carol Reschke's Ecological Communities of New York State. New York Natural Heritage Program, New York State Department of Environmental Conservation, Albany, NY.

Gawler, S. and A. Cutko. 2010. Natural Landscapes of Maine: A Guide to Natural Communities and Ecosystems. Maine Natural Areas Program, Maine Department of Conservation, Augusta, Maine.

NatureServe. 2021. NatureServe Explorer: An online encyclopedia of life [web application]. NatureServe, Arlington, Virginia. <https://explorer.natureserve.org/>. (accessed 10 July. 2021).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Official Soil Series Descriptions. Available online. (accessed 11 Aug. 2021).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Climate Research Station Data. Available online. (accessed 23 June. 2021).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Survey Geographic (SSURGO) Database for [MLRA 141, Maine]. Available online. (accessed 14 Oct. 2021).

Sperduto, D.D. and William F. Nichols. 2011. Natural Communities of New Hampshire. 2nd Ed. NH Natural Heritage Bureau, Concord, NH. Pub. UNH Cooperative Extension, Durham, NH.

Swain, P. C. 2020. Classification of the Natural Communities of Massachusetts. Massachusetts Division of Fisheries and Wildlife, Westborough, MA

USNVC [United States National Vegetation Classification]. 2017. United States National Vegetation Classification Database V2.01. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC. Available The U.S. National Vegetation Classification (usnvc.org) (accessed 2 July. 2021).

## Contributors

Christopher Mann  
Jamin Johanson

## Approval

Greg Schmidt, 10/07/2024

## Acknowledgments

Nels Barrett, Nick Butler, and Carl Bickford provided considerable review of this ecological site concept.

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/10/2025
Approved by	Greg Schmidt
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

### 1. Number and extent of rills:

---



2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**

---