

# Ecological site R144AY046RI Subaqueous Freshwater Organic Deposits

Last updated: 10/04/2024 Accessed: 05/13/2025

#### **General information**

**Provisional**. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

#### **MLRA** notes

Major Land Resource Area (MLRA): 144A-New England and Eastern New York Upland, Southern Part

#### MLRA 144A: New England and Eastern New York Upland, Southern Part

The eastern half of the eastern part of this MLRA is in the Seaboard Lowland Section of the New England Province of the Appalachian Highlands. The western half of the eastern part and the southeastern half of the western part are in the New England Upland Section of the same province and division. The northwestern half of the western part is in the Hudson Valley Section of the Valley and Ridge Province of the Appalachian Highlands. This MLRA is a very scenic area of rolling to hilly uplands that are broken by many gently sloping to level valleys that terminate in coastal lowlands. Elevation ranges from sea level to 1,000 feet (0 to 305 meters) in much of the area, but it is 2,000 feet (610 meters) on some hills. Relief is mostly about 6 to 65 feet (2 to 20 meters) in the valleys and about 80 to 330 feet (25 to 100 meters) in the uplands.

This area has been glaciated and consists almost entirely of till hills, drumlins, and bedrock-controlled uplands with a mantle of till. It is dissected by narrow glacio-fluvial valleys. The southernmost boundary of the area marks the farthest southward extent of Wisconsinian glaciation on the eastern seaboard. The river valleys and coastal plains are filled with glacial lake sediments, marine sediments, and glacial outwash. The bedrock in the eastern half of the area consists primarily of igneous and metamorphic rocks of early Paleozoic age. Granite is the most common igneous rock, and gneiss, schist, and slate are the most common metamorphic rocks. In the parts of the MLRA in eastern and southeastern New York, Devonian- to Pennsylvanian-age sandstone, shale, and limestone are dominant. Carbonate rocks, primarily dolomite and limestone, are the dominant kinds of bedrock in the part of this MLRA in northwestern Connecticut.

## **Ecological site concept**

The site consists of very deep organic subaqueous soils permanently submerged beneath approximately 0 to 3 meters of water on submerged organic landscapes of freshwater ponds, lakes, and reservoirs. The soils formed in thick organic deposits. Slope ranges from 0 to 3 percent. Representative soils are Tuckertown and Wickford. Areas of this soil are used for recreational fishing and swimming. Vegetation includes bladderwort (Utricularia sp.), fanwort (*Cabomba caroliniana*), variable milfoil (*Myriophyllum heterophyllum*), white water lily (Nymphaea ordorata), and yellow water lily (Nuphar variegatum).

#### Similar sites

F144AY028MA Wet Outwash

#### Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified

## **Physiographic features**

The site occurs on permanently submerged outwash and plains of freshwater ponds, lakes, and reservoirs.

	.,
Landforms	(1) Lowland > Outwash plain
Runoff class	Negligible
Flooding frequency	None
Ponding frequency	Frequent
Elevation	0–173 m
Slope	0–3%
Water table depth	0 cm
Aspect	Aspect is not a significant factor

#### Table 2. Representative physiographic features

## **Climatic features**

The Koppen-Geiger climate classification of the area in which this MLRA occurs varies between Dfb (Warmsummer humid continental) in the North, and Dfa (Hot-summer humid continental) in the southern portion of the MLRA. Precipitation is usually uniformly distributed throughout the year. Near the coast, however, it is slightly lower in summer. Precipitation is slightly higher in spring and fall in inland areas. Rainfall occurs as high-intensity, convective thunderstorms during the summer. During the winter, most of the precipitation occurs as moderateintensity storms (northeasters) that produce large amounts of rain or snow. The freeze-free period increases in length to the south.

Frost-free period (characteristic range)	124-145 days
Freeze-free period (characteristic range)	152-188 days
Precipitation total (characteristic range)	1,143-1,219 mm
Frost-free period (actual range)	116-145 days
Freeze-free period (actual range)	147-190 days
Precipitation total (actual range)	1,067-1,219 mm
Frost-free period (average)	134 days
Freeze-free period (average)	170 days
Precipitation total (average)	1,168 mm

#### Table 3. Representative climatic features



Figure 1. Monthly precipitation range



Figure 2. Monthly minimum temperature range



Figure 3. Monthly maximum temperature range



Figure 4. Monthly average minimum and maximum temperature



Figure 5. Annual precipitation pattern



Figure 6. Annual average temperature pattern

### **Climate stations used**

- (1) RUTLAND [USC00436995], Rutland, VT
- (2) GLOVERSVILLE [USC00303319], Gloversville, NY
- (3) ALLENTOWN INTL AP [USW00014737], Allentown, PA
- (4) NEW BEDFORD MUNI AP [USW00094726], New Bedford, MA
- (5) DURHAM 2 SSW [USW00054795], Durham, NH
- (6) BELCHERTOWN [USC00190562], Belchertown, MA

## Influencing water features

Subaqueous soils differ from subaerial, or terrestrial, soils by having perennial water on the soil surface. These soils occur in shallow freshwater and marine environments, such as ponds, lakes, and the subtidal areas of estuaries and tidal embayments.

#### Wetland description

National Wetland Classification (Cowardin et al., 1979):

System: Lacustrine

Subsystem: Littoral

Class: Unconsolidated Bottom/Shore, Aquatic Bed, Emergent

Subclass: Mud, Algal, Rooted/Floating Vascular, Organic, Persistent/Non-persistent

Water Regime: Permanently Flooded

## Soil features

The site consists of very deep organic subaqueous Wickford muck soil, permanently submerged beneath

approximately 0 to 3 meters of water on submerged organic landscapes of freshwater ponds, lakes, and reservoirs. The soil formed in thick organic deposits. Soil drainage class is subaqueous.

Table 4.	. Representative	soil features
----------	------------------	---------------

Parent material	(1) Organic material
Surface texture	(1) Muck
Drainage class	Subaqueous
Permeability class	Slow
Depth to restrictive layer	183 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (Depth not specified)	27.94 cm
Soil reaction (1:1 water) (Depth not specified)	3–6.5
Subsurface fragment volume <=3" (Depth not specified)	0%
Subsurface fragment volume >3" (Depth not specified)	0%

## **Ecological dynamics**

Vegetation includes white water lily (Nymphaea ordorata), and yellow water lily (Nuphar variegatum, as well as bladderwort (Utricularia sp.), fanwort (*Cabomba caroliniana*), variable milfoil (*Myriophyllum heterophyllum*).

## State and transition model

## 144AY046–Subaqueous Freshwater Organic Deposits



Transition	Drivers/practices
T1-2	Dredging
R2-1	Fill and aquatic planting

Figure 8. STM\_144AY046\_Subaqueous\_Freshwater\_Organic\_Deposit

## State 1 Minimally-managed reference state

The vegetation of brackish tidal wetlands is variable, ranging from Brackish meadows of *Spartina patens*, *Eleocharis uniglumis*, and *Panicum virgatum* to Brackish "reed" marshes, dominated by *Typha angustifolia* and Typha x glauca and bulrushes Schenoplectus (aka Bulboschoenus spp.).

Community 1.1 Nuphar advena - Nymphaea odorata aquatic vegetation

Community 1.2 Vallisneria americana - potamogeton perfoliatus aquatic vegetation

Community 1.3 Unvegetated

State 2 Semi-natural / dredged state

Community 2.1 Invasive spp. Myriophyllum

Community 2.2 Unvegetated

Transition T1A State 1 to 2

Dredging

# Restoration pathway R2A State 2 to 1

Fill and aquatic planting

## Additional community tables

## Inventory data references

Future work is needed, as described in a future project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

## **Other references**

#### REFERENCES

Bradley, M.P. and Stolt, M.H., 2003. Subaqueous soil-landscape relationships in a Rhode Island estuary. Soil Science Society of America Journal, 67(5)\_1487-1495.

Ditzler, C.A., Ahrens, R.J., Rabenhorst, M.C., Stolt, M., Hipple, K., and Turenne, J. s.d. Classification, Mapping, and Interpretation of Subaqueous Soils. Unpubl. Manuscript.

Stolt, M., Bradley, M., Turenne, J., Payne, M., Scherer, E., Cicchetti, G., Shumchenia, E., Guarinello, M., King, J., Boothroyd, J. and Oakley, B., 2011. Mapping shallow coastal ecosystems: a case study of a Rhode Island Iagoon. Journal of Coastal Research, 27(6A)\_1-15.

### Contributors

Nels Barrett, Ph.D. Michael Margo

## Approval

Greg Schmidt, 10/04/2024

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/13/2025
Approved by	Greg Schmidt
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

#### Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:

<sup>4.</sup> Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values):
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth ( in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):

- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 17. Perennial plant reproductive capability: