

## Ecological site R144AY047RI Subaqueous Haline Glacial Deposits

Last updated: 10/04/2024  
Accessed: 05/11/2025

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 144A–New England and Eastern New York Upland, Southern Part

MLRA 144A: New England and Eastern New York Upland, Southern Part

The eastern half of the eastern part of this MLRA is in the Seaboard Lowland Section of the New England Province of the Appalachian Highlands. The western half of the eastern part and the southeastern half of the western part are in the New England Upland Section of the same province and division. The northwestern half of the western part is in the Hudson Valley Section of the Valley and Ridge Province of the Appalachian Highlands. This MLRA is a very scenic area of rolling to hilly uplands that are broken by many gently sloping to level valleys that terminate in coastal lowlands. Elevation ranges from sea level to 1,000 feet (0 to 305 meters) in much of the area, but it is 2,000 feet (610 meters) on some hills. Relief is mostly about 6 to 65 feet (2 to 20 meters) in the valleys and about 80 to 330 feet (25 to 100 meters) in the uplands.

This area has been glaciated and consists almost entirely of till hills, drumlins, and bedrock-controlled uplands with a mantle of till. It is dissected by narrow glacio-fluvial valleys. The southernmost boundary of the area marks the farthest southward extent of Wisconsinian glaciation on the eastern seaboard. The river valleys and coastal plains are filled with glacial lake sediments, marine sediments, and glacial outwash. The bedrock in the eastern half of the area consists primarily of igneous and metamorphic rocks of early Paleozoic age. Granite is the most common igneous rock, and gneiss, schist, and slate are the most common metamorphic rocks. In the parts of the MLRA in eastern and southeastern New York, Devonian- to Pennsylvanian-age sandstone, shale, and limestone are dominant. Carbonate rocks, primarily dolomite and limestone, are the dominant kinds of bedrock in the part of this MLRA in northwestern Connecticut.

### Ecological site concept

The Subaqueous Haline Glacial Deposits ecological site consists of very deep, subaqueous soils that are permanently submerged beneath 10 through 150 cm of tidal estuarine water in mainland coves and submerged mainland beaches within coastal lagoons and open bays. The soils are formed in sandy marine deposits over sandy and gravelly outwash. Slope ranges from 0 through 5 percent. Representative soil is Anguilla. This soil supports submerged aquatic vegetation and aquatic habitats. Some areas are vegetated with eelgrass (*Zostera marina*), widgeon grass (*Ruppia maritima*) and native macroalgae. Vegetative cover typically ranges from 0 to 35 percent.

This soil supports submerged aquatic vegetation and aquatic habitats. The area is used by recreational fishermen for the harvest of crabs. In addition fishing is commonplace and the species found in the area are smelt, small cod, flounder, scup, menhaden, and white perch.

### Associated sites

R144AY048RI	Subaqueous Haline Low Energy Basins
R144AY049RI	Subaqueous Haline Slopes
R144AY050RI	Subaqueous Haline Flats

## Similar sites

R144AY049RI	Subaqueous Haline Slopes
R144AY050RI	Subaqueous Haline Flats
R144AY048RI	Subaqueous Haline Low Energy Basins

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Zostera marina</i> (2) <i>Ruppia maritima</i>

## Physiographic features

This site occurs on permanently submerged haline embayments, headlands, or coves. These may occur adjacent to glaciated uplands with numerous boulders and stones that are exposed to the air at low tides. Slope ranges from 0 through 5 percent; below sea level.

Table 2. Representative physiographic features

Landforms	(1) Bay > Bay bottom (2) Estuary > Headland (3) Lagoon > Cove
Elevation	0 ft
Slope	0–5%
Water table depth	0 in
Aspect	Aspect is not a significant factor

## Climatic features

The Koppen-Geiger climate classification of the area in which this MLRA occurs varies between Dfb (Warm-summer humid continental) in the North, and Dfa (Hot-summer humid continental) in the southern portion of the MLRA. Precipitation is usually uniformly distributed throughout the year. Near the coast, however, it is slightly lower in summer. Precipitation is slightly higher in spring and fall in inland areas. Rainfall occurs as high-intensity, convective thunderstorms during the summer. During the winter, most of the precipitation occurs as moderate-intensity storms (northeasters) that produce large amounts of rain or snow. The freeze-free period increases in length to the south.

Table 3. Representative climatic features

Frost-free period (characteristic range)	142-174 days
Freeze-free period (characteristic range)	176-214 days
Precipitation total (characteristic range)	46-49 in
Frost-free period (actual range)	140-181 days
Freeze-free period (actual range)	168-216 days
Precipitation total (actual range)	44-51 in
Frost-free period (average)	157 days
Freeze-free period (average)	196 days
Precipitation total (average)	47 in




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range






Figure 4. Monthly average minimum and maximum temperature



**Figure 5. Annual precipitation pattern**



**Figure 6. Annual average temperature pattern**

## Climate stations used

- (1) ROCHESTER SKYHAVEN AP [USW00054791], Rochester, NH
- (2) BEVERLY MUNI AP [USW00054733], Wenham, MA
- (3) PLYMOUTH MUNI AP [USW00054769], Carver, MA
- (4) NEWPORT STATE AP [USW00014787], Middletown, RI
- (5) GROTON NEW LONDON AP [USW00014707], Groton, CT
- (6) BRIDGEPORT SIKORSKY MEM AP [USW00094702], Stratford, CT
- (7) DOBBS FERRY ARDSLEY [USC00302129], Ardsley, NY

## Influencing water features

Subaqueous soils differ from subaerial, or terrestrial, soils by having perennial water on the soil surface in either shallow freshwater or marine environments. These subaqueous soils are formed in the subtidal areas of estuaries and tidal embayments.

## Wetland description

National Wetland Classification (Cowardin et al., 1979):

System: Marine/estuarine

Subsystem: Subtidal

Class: Unconsolidated Bottom, Aquatic Bed, Unconsolidated Shore, Emergent

Subclass: Cobble-Gravel, Sand, Mud, Algal, Rooted/Floating Vascular, Organic, Persistent/Non-persistent

Water Regime: Subtidal

## Soil features

This site consists of very deep, subaqueous soils formed in sandy marine deposits over sandy and gravelly glaciofluvial deposits, submerged terrestrial loess, or till deposits. Representative soils are Napatree and Anguilla. The saturated hydraulic conductivity is high or very high.

**Table 4. Representative soil features**

Parent material	(1) Loess (2) Glaciofluvial deposits (3) Marine deposits (4) Till
Surface texture	(1) Sandy loam (2) Coarse sand (3) Silt loam (4) Loam (5) Very fine sandy loam (6) Fine sandy loam (7) Loamy sand (8) Loamy coarse sand (9) Fine sand (10) Sand (11) Gravelly sand (12) Mucky sand
Family particle size	(1) Sandy (2) Coarse-loamy
Drainage class	Subaqueous
Permeability class	Very slow
Depth to restrictive layer	72 in
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Soil reaction (1:1 water) (Depth not specified)	3.2–7.4
Subsurface fragment volume <=3" (Depth not specified)	0–50%
Subsurface fragment volume >3" (Depth not specified)	0%

## Ecological dynamics

[Caveat: The vegetation information contained in this section and is only provisional, based on concepts, not yet validated with field work.\*]

The vegetation groupings described in this section are based on the terrestrial ecological system classification and vegetation associations developed by NatureServe (Comer 2003). Terrestrial ecological SYSTEMS are specifically defined as a group of plant community-types called ASSOCIATIONS that tend to [co-]occur within landscapes with similar ecological processes, substrates, and/or environmental gradients. Any given system will typically manifest itself in a landscape at intermediate geographic scales of tens-to-thousands of hectares and will persist for 50 or more years. A vegetation association is a plant community that is much more specific to a given soil, geology, landform, climate, hydrology, and disturbance history. It is the basic unit for vegetation classification and recognized by the US National Vegetation Classification (US FDGC 2008). Each association will be named by the diagnostic and often dominant species that occupy the different height strata (tree, sapling, shrub, and herb). Within the NatureServe Explorer database (NatureServe, 2015), ecological systems are numbered by a Community Ecological System Code (CES) and individual vegetation associations are assigned an identification number called a Community Element Global Code (CEGL).


Additional and more localized vegetation information is provided by the State Natural Heritage Programs of Connecticut (Metzler and Barrett 2001), Massachusetts (Swain and Kearsley 2001), New Hampshire (Sperduto and Nichols, 2011), New York (Edinger et al., 2014), and Rhode Island (Enser and Lungren, 2006).

The Subaqueous Haline Submerged Glacial Deposits is characterized by the North Atlantic Rocky Intertidal system (CES201.048), and the North Atlantic Tidal Sand Flat system (CES201.049). The site consists of very deep,

subaqueous soils that are permanently submerged beneath 10 through 150 cm of tidal estuarine water in mainland coves and submerged mainland beaches within coastal lagoons and open bays. The soils are formed in sandy marine deposits over sandy and gravelly outwash. Slope ranges from 0 through 5 percent. Representative soil is Anguilla. Some areas are vegetated with native macroalgae, eelgrass (*Zostera marina*), and widgeon grass (*Ruppia maritima*). Vegetative cover typically ranges from 0 to 35 percent. The site is subject to recreational fishing, and crabbing. The site is subject to dredging activities and aquaculture management.

[\*Caveat] The information presented is representative of very complex vegetation communities. Key indicator plants and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and geography. The reference plant community is not necessarily the management goal. The drafts of species lists are merely representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.

## State and transition model



### State 1 Minimally-managed reference state

The reference state includes: • *Zostera marina* Atlantic Aquatic Vegetation Translated Name: Seawrack Atlantic Aquatic Vegetation Common Name: North Atlantic Seawrack Bed (CEGL004336) Other associations may include: •

*Ruppia maritima* Acadian/Virginian Zone Temperate Aquatic Vegetation Translated Name: Widgeongrass  
Acadian/Virginian Zone Temperate Aquatic Vegetation Common Name: North Atlantic Coast Widgeongrass Bed (CEGL006167) • *Ascophyllum nodosum* - *Fucus vesiculosus* Tidal Algal Nonvascular Vegetation Translated Name: Yellow Tang - Black Tang Tidal Algal Nonvascular Vegetation Common Name: New England Rocky Intertidal Rocky Shore (CEGL006341) • *Laminaria agardhii* Tidal Algal Nonvascular Vegetation Translated Name: Common Southern Kelp Tidal Algal Nonvascular Vegetation Common Name: New England Kelp Bed (CEGL006344)

## Community 1.1

### Eelgrass (*Zostera marina*) Atlantic aquatic vegetation (CEGL006167)

*Zostera marina* Atlantic Aquatic Vegetation Translated Name: Seawrack Atlantic Aquatic Vegetation Common Name: North Atlantic Seawrack Bed (CEGL004336) Eelgrass (*Zostera marina*) is dominant and occurs most often in nearly pure stands. Widgeonweed (*Ruppia maritima*) can occur sporadically in this association, especially as waters become less saline. Additional associated species include various macroalgae, especially *Ulva lactuca*, *Enteromorpha* spp., *Cladophora* spp., and *Polysiphonia* spp. Where water is less saline, *Enteromorpha*, *Chaetomorpha*, *Gracilaria*, *Agardhiella*, and *Ectocarpus* can occur. Elevation/depth of the beds is determined by low tide level at the upper end and light penetration at the lower end, the latter being a function of water depth and turbidity. The beds generally occur in areas with only moderate wave action where salinity fluctuations are minor. Eel-grass beds tend to stabilize and enrich substrate and provide habitat for epiphytes and other marine organisms. (Source: NatureServe 2018 [accessed 2019], USNVC 2017 [accessed 2019]). Cross-referenced plant community concepts (typically by political State): CT: Eelgrass Permanently Flooded Vegetation (Metzler and Barrett, 2006) MA: Seagrass community (Swain and Kearsley, 2001) NH: Eelgrass Bed (Sperduto and Nichols, 2011) NY: Marine Eelgrass Meadow (Edinger et al., 2014) RI: Marine Subtidal Aquatic Bed (Enser and Lundgren, 2006)

## Community 1.2

### Widgeonweed (*Ruppia maritima*) Acadian/Virginian Zone Temperate Aquatic Vegetation (CEGL006167)

This brackish/saline tidal community of the central and northern Atlantic coast is dominated by widgeonweed (*Ruppia maritima*). It occurs in large beds in estuarine bays as well as small patches within saline/brackish tidal creeks. Substrates are sand or muck, and salinity is generally more brackish. Widgeonweed has a wide range of salinity tolerance and overlaps with other species, although generally not in the same locations. Common associates in more brackish/freshwater tidal conditions include horned pondweed (*Zannichellia palustris*), sago false pondweed (*Stuckenia pectinata*), and clasping-leaved pondweed (*Potamogeton perfoliatus*) or eelgrass (*Zostera marina*) as waters get deeper and more saline. There can also be a diverse array of macroalgae. (Source: NatureServe 2018 [accessed 2019], USNVC 2017 [accessed 2019]). Cross-referenced plant community concepts (typically by political State): CT: Widgeonweed Permanently Flooded Vegetation (Metzler and Barrett, 2006) MA: Seagrass community (Swain and Kearsley, 2001) NH: Undisclosed (Sperduto and Nichols, 2011) NY: Brackish Subtidal Aquatic Bed (Edinger et al., 2014) RI: Marine Subtidal Aquatic Bed (Enser and Lundgren, 2006)

## Community 1.3

### Unvegetated

Unvegetated

## State 2

### Semi-natural / dredged state

The Semi-natural State would expect plant communities where ecological processes are primarily operating with some land conditioning in the past or present, e.g., aquaculture, dredge spoils.

## Community 2.1

### Eelgrass (*Zostera marina*) Atlantic aquatic vegetation (CEGL006167)

## Community 2.2

### Widgeonweed (*Ruppia maritima*) Acadian/Virginian Zone Temperate Aquatic Vegetation

**(CEGL006167)**

**Community 2.3  
Unvegetated**

**State 3  
Cultural state**

The Cultural State would expect the ecological site to be very strongly conditioned by management conversion.

**Community 3.1  
Aquaculture**

**Transition T1A  
State 1 to 2**

Dredging

**Transition T1B  
State 1 to 3**

Aquacultural practices

**Restoration pathway R2A  
State 2 to 1**

Fill and seagrass planting

**Transition T2A  
State 2 to 3**

Aquacultural practices

**Restoration pathway R3A  
State 3 to 1**

Fill and seagrass planting

**Transition T3A  
State 3 to 2**

Dredging

**Additional community tables**

**Inventory data references**

Future work is needed, as described in a future project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

**Other references**

REFERENCES



Bradley, M.P. and Stolt, M.H., 2003. Subaqueous soil-landscape relationships in a Rhode Island estuary. Soil Science Society of America Journal, 67(5)\_1487-1495.

Ditzler, C.A., Ahrens, R.J., Rabenhorst, M.C., Stolt, M., Hipple, K., and Turenne, J. s.d. Classification, Mapping, and Interpretation of Subaqueous Soils. Unpubl. Manuscript.

Stolt, M., Bradley, M., Turenne, J., Payne, M., Scherer, E., Cicchetti, G., Shumchenia, E., Guarinello, M., King, J., Boothroyd, J. and Oakley, B., 2011. Mapping shallow coastal ecosystems: a case study of a Rhode Island lagoon. Journal of Coastal Research, 27(6A)\_1-15.

## Contributors

Nels Barrett, Ph.D.  
Michael Margo

## Approval

Greg Schmidt, 10/04/2024

## Acknowledgments

Acknowledgements to Dr. Mark Stolt (URI) for the development of the subaqueous soil descriptions.

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/11/2025
Approved by	Greg Schmidt
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

### 1. Number and extent of rills:

---

### 2. Presence of water flow patterns:

---

### 3. Number and height of erosional pedestals or terracettes:

---

### 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

- 
5. **Number of gullies and erosion associated with gullies:**
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:**
- 
7. **Amount of litter movement (describe size and distance expected to travel):**
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 

17. **Perennial plant reproductive capability:**
-