

Ecological site F144BY402ME

Clay Hills

Last updated: 9/27/2024
Accessed: 05/11/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 144B—New England and Eastern New York Upland, Northern Part

This major land resource area (MLRA) is in Maine (56 percent), New Hampshire (22 percent), Vermont (14 percent), Massachusetts (6 percent), Connecticut (1 percent), and New York (1 percent). It makes up about 22,728 square miles (58,864 square kilometers). The MLRA consists of a relatively young landscape shaped by the Laurentide Ice Sheet, which covered the region from 35,000 to 10,000 years ago. Rolling hills of dense basal till converge on ridges of shallow bedrock that were scoured by glacial ice. River valleys that were flooded by melting glacial water or seawater house large expanses of glacial outwash and stratified drift in inland areas and, to a lesser extent, glaciomarine and glaciolacustrine sediment deposits in coastal areas. Organic bogs, ablation till, and alluvial flood plains make up the remaining portions of the MLRA.

The soils in this region are dominantly Entisols, Spodosols, and Inceptisols. They commonly have a fragipan. The dominant suborders are Ochrepts, Orthods, Aquepts, Fluvents, and Saprists. The soils in the region dominantly have a frigid soil temperature regime with some cryic areas at higher elevation, a udic soil moisture regime, and mixed mineralogy. Most of the land is forested, and 98 percent is privately owned. Significant amounts of forest products are produced including lumber, pulpwood, Christmas trees, and maple syrup. Principal agricultural crops include forage and grains for dairy cattle, potatoes, apples, and blueberries. Wildlife habitat and recreation are important land uses. Stoniness, steep slopes, and poor drainage limit the use of many of the soils.

Classification relationships

NRCS:

Land Resource Region: R—Northeastern Forage and Forest Region

MLRA: 144B—New England and Eastern New York Upland, Northern PartMLRA resources

Major Land Resource Area (MLRA): 144B—New England and Eastern New York Upland, Northern Part

Ecological site concept

This site occurs on the remnants of gently-sloping (0-15%) glacial lake beds, lake plains and terraces. Soils have clay textures and very few rock fragments throughout the profile. These are moderately-well and well drained soils with a seasonally-high water table within 18-36 inches of the soil surface. Tree species are diverse, typically with conifers such as red spruce, larch, and white pine more abundant than hardwoods, which include red maple and grey birch.

Associated sites

F144BY301ME	Loamy Till Swamp The Loamy Till Swamp site occurs lower in the watershed than the Loamy Flat site. The two sites occur together along a soil drainage gradient from somewhat poorly to poorly- and very poorly-drained.
-------------	---

F144BY502ME	Loamy Till Toeslope The Loamy Flat site and Loamy Till Toeslopes site share many of the same soils, but Loamy Flats occur on flatter areas grading into wetlands and produce spruce-fir forests, whereas the Loamy Till Toeslope occurs at the base of slopes and produces semi-rich mixedwood forests.
F144BY304ME	Wet Clay Flat The Loamy Flat site occurs on somewhat-poorly and poorly-drained soil complexes that are somewhat drier and have significantly less understory production than the Wet Loamy Flat site.

Similar sites

F144BY502ME	Loamy Till Toeslope The Loamy Flat site and Loamy Till Toeslopes site share many of the same soils, but Loamy Flats occur on flatter areas grading into wetlands and produce spruce-fir forests, whereas the Loamy Till Toeslope occurs at the base of slopes and produces semi-rich mixedwood forests.
F144BY304ME	Wet Clay Flat The Loamy Flat site occurs on somewhat-poorly and poorly-drained soil complexes that are somewhat drier and have significantly less understory production than the Wet Loamy Flat site.

Table 1. Dominant plant species

Tree	(1) <i>Pinus strobus</i> (2) <i>Tsuga canadensis</i>
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site is found on marine and lake sediments that were deposited during deglaciation when much of coastal Maine was submerged under the Atlantic Ocean. As the ocean subsided, it left behind clay sediments from which these soils developed. In steeper areas, gullying is a natural process forming an eroded, wavy land surface.

Table 2. Representative physiographic features

Landforms	(1) Upland > Marine terrace (2) Plains > Marine terrace (3) Lake plain > Lake plain (4) River valley > Stream terrace (5) Coastal plain (6) Outwash delta (7) Outwash plain (8) Plain
Flooding frequency	None
Ponding duration	Brief (2 to 7 days) to long (7 to 30 days)
Ponding frequency	None to frequent
Elevation	0–1,500 ft
Slope	0–45%
Ponding depth	0–6 in
Water table depth	12–36 in
Aspect	Aspect is not a significant factor

Climatic features

The climate is humid and temperate and is characterized by warm summers and cold winters. Precipitation generally is evenly distributed throughout the year. Near the coast, it is slightly lower in summer. In inland areas, it is slightly higher in spring and fall. Rainfall occurs during high-intensity, convective thunderstorms in summer. In

winter, most of the precipitation occurs as moderate-intensity storms (northeasters) that produce large amounts of rain or snow. Heavy snowfalls commonly occur late in winter. Temperatures and the length of the freeze-free period increase from north to south and closer to the coast.

This major land resource area (MLRA) covers four states and may have substantial climate variability among locations: Maine (56 percent), New Hampshire (22 percent), Vermont (14 percent), Massachusetts (6 percent), Connecticut (1 percent), and New York (1 percent).

Table 3. Representative climatic features

Frost-free period (characteristic range)	117-140 days
Freeze-free period (characteristic range)	144-170 days
Precipitation total (characteristic range)	42-48 in
Frost-free period (actual range)	98-146 days
Freeze-free period (actual range)	133-180 days
Precipitation total (actual range)	40-54 in
Frost-free period (average)	126 days
Freeze-free period (average)	159 days
Precipitation total (average)	46 in




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) BELFAST [USC00170480], Belfast, ME
- (2) ACADIA NP [USC00170100], Bar Harbor, ME
- (3) CORINNA [USC00171628], Corinna, ME

- (4) DOVER-FOXCROFT WWTP [USC00171975], Dover Foxcroft, ME
- (5) FARMINGTON [USC00172765], Farmington, ME
- (6) GARDINER [USC00173046], Gardiner, ME
- (7) JONESBORO [USC00174183], Addison, ME
- (8) LEWISTON [USC00174566], Auburn, ME
- (9) MADISON [USC00174927], Anson, ME
- (10) NEWCASTLE [USC00175675], Newcastle, ME
- (11) ORONO [USC00176430], Old Town, ME
- (12) WATERVILLE TRTMT PLT [USC00179151], Waterville, ME
- (13) WEST ROCKPORT 1 NNW [USC00179593], Rockport, ME
- (14) AUGUSTA STATE AP [USW00014605], Augusta, ME
- (15) BANGOR INTL AP [USW00014606], Bangor, ME
- (16) PORTLAND INTL JETPORT [USW00014764], Portland, ME

Influencing water features

This site is not typically influenced by streams or wetlands.

Soil features

This site is characterized by deep clay soils formed in marine or lacustrine sediments. They are moderately well- to well-drained and have very few rock fragments. Soil pH is typically somewhat acidic, but not as acidic as soils formed in other parent materials.

Table 4. Representative soil features

Parent material	(1) Glaciomarine deposits (2) Glaciolacustrine deposits
Surface texture	(1) Silt loam (2) Fine sandy loam
Drainage class	Somewhat poorly drained to well drained
Permeability class	Very slow
Soil depth	0–60 in
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (3-8in)	Not specified
Soil reaction (1:1 water) (3.6-7.3in)	Not specified
Subsurface fragment volume <=3" (0-2in)	Not specified
Subsurface fragment volume >3" (0in)	Not specified

Ecological dynamics

[Caveat: The vegetation information contained in this section and is only provisional, based on concepts, and future projects support validation through field work. *] The vegetation groupings described in this section are based on the terrestrial ecological system classification and vegetation associations developed by NatureServe (Comer et al., 2003) and localized associations provided by the New York Natural Heritage Program (Edinger et al., 2014), Maine Natural Areas Program (Gawler and Cutko, 2010), New Hampshire Natural Heritage Program (Sperduto and Nichols, 2011), and Massachusetts Division of Fisheries and Wildlife (Swain, 2020).

This site typically supports stands of conifer-dominated mixedwood. Tree species are diverse, typically with conifers such as red spruce, larch, and white pine more abundant than hardwoods, which include red maple and grey birch.

Logging and blowdowns create open patches where herbaceous and mid-seral communities occur following disturbance. Insects and disease may weaken trees on this site as well. The lack of rocks and relatively higher soil nutrient levels of this site are conducive to land-use conversion from forest land to cropland, hayland, or pasture. Pine plantations have been observed on this site.


Relationship to Other Classification Systems

This site includes the following state natural heritage program types:

NONE

State and transition model

F144BY402ME – Clay Hills



State 1

Reference State / Current Potential

Community 1.1

Conifer-dominated Mixedwoods Forest

Softwoods predominate, typically white pine, red spruce and larch, with some hardwoods present.

Community 1.2

Herbaceous Phase

Wild raspberry, ferns, and other herbs colonize the open land

Community 1.3

Doghair Forest

Spruce, white pine, larch and balsam fir saplings growing close together with sparsely occurring red maple and other hardwoods.

Community 1.4

Mixedwoods Forest

50-80 year old hardwoods. Early successional species (white birch, fir, grey birch, aspen) dying out.

Pathway P1.1-1.2

Community 1.1 to 1.2

windthrow, blowdown, fire

Pathway P1.2-1.3

Community 1.2 to 1.3

vegetation development (succession)

Pathway P1.3-1.4

Community 1.3 to 1.4

vegetation development (succession)

Pathway P1.4-1.1

Community 1.4 to 1.1

windthrow, blowdown, fire

Pathway P1.4-1.2

Community 1.4 to 1.2

windthrow, blowdown, fire

State 2

Grassland / Hay land

Community 2.1

Pasture or Hay Land (FSG4)

Cleared and planted fields of mostly perennial herbaceous species.

State 3

Crop Land

Community 3.1

Annual or Perennial Crops

Cleared and cultivated fields, heavily managed with regular soil disturbance.

State 4

White Pine

Community 4.1

Herbs and Shrubs

Wild raspberry, ferns, and other herbs colonize the open land

Community 4.2

White Pine Forest

Single age white pine forest.

Pathway P4.1-4.2 Community 4.1 to 4.2

Vegetation development (succession)

Pathway P4.2-4.1 Community 4.2 to 4.1

harvest, logging

Conservation practices

Forest Stand Improvement
Forest Land Management

Transition T1-2 State 1 to 2

tree removal, pasture or hayfield establishment

Conservation practices

Clearing and Snagging
Land Clearing
Invasive Plant Species Control
Managed Haying/Grazing

Transition T1-3 State 1 to 3

Tree clearing, crop establishment

Conservation practices

Clearing and Snagging
Cover Crop
Land Clearing

Transition T1-4 State 1 to 4

selective harvest

Conservation practices

Forest Stand Improvement
Forest Land Management

Transition R2-1 State 2 to 1

abandonment, vegetation development (succession), planting

Conservation practices

Tree/Shrub Site Preparation
Tree/Shrub Establishment
Upland Wildlife Habitat Management
Restoration and Management of Natural Ecosystems
Native Plant Community Restoration and Management
Invasive Plant Species Control
Managed Haying/Grazing

Transition T2-4
State 2 to 4

tree establishment

Conservation practices

Tree/Shrub Site Preparation
Tree/Shrub Establishment
Invasive Plant Species Control

Restoration pathway R3-1
State 3 to 1

abandonment, vegetation development (succession), tree planting

Conservation practices

Tree/Shrub Establishment
Upland Wildlife Habitat Management
Tree/Shrub Pruning
Restoration and Management of Natural Ecosystems
Native Plant Community Restoration and Management
Invasive Plant Species Control

Transition T3-4
State 3 to 4

tree planting

Conservation practices

Tree/Shrub Site Preparation
Tree/Shrub Establishment
Invasive Plant Species Control

Restoration pathway R4-1
State 4 to 1

abandonment, vegetation development (succession), plantings

Conservation practices

Tree/Shrub Site Preparation
Tree/Shrub Establishment
Upland Wildlife Habitat Management
Restoration and Management of Natural Ecosystems
Native Plant Community Restoration and Management
Invasive Plant Species Control

Restoration pathway T4-2

State 4 to 2

Tree removal, pasture or hay land establishment

Conservation practices

Clearing and Snagging
Land Clearing

Transition T4-3

State 4 to 3

tree removal, cropland establishment

Conservation practices

Clearing and Snagging
Cover Crop
Land Clearing

Additional community tables

Inventory data references

Future work is needed, as described in a future project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

Other references

Comer, P., D. Faber-Langendoen, R. Evans, S. Grawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schultz, K. Snow, and J. Teague. 2003. Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems. NatureServe, Arlington, Virginia

Edinger, G. J., D. J. Evans, S. Gebauer, T. G. Howard, D. M. Hunt, and A. M. Olivero (editors). 2014. Ecological Communities of New York State. Second Edition. A revised and expanded edition of Carol Reschke's Ecological Communities of New York State. New York Natural Heritage Program, New York State Department of Environmental Conservation, Albany, NY.

Gawler, S. and A. Cutko. 2010. Natural Landscapes of Maine: A Guide to Natural Communities and Ecosystems.

Maine Natural Areas Program, Maine Department of Conservation, Augusta, Maine.

NatureServe. 2021. NatureServe Explorer: An online encyclopedia of life [web application]. NatureServe, Arlington, Virginia. <https://explorer.natureserve.org/>. (accessed 10 July. 2021).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. Agricultural Handbook 296

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Official Soil Series Descriptions. Available online. (accessed 11 Aug. 2021).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Climate Research Station Data. Available online. (accessed 23 June. 2021).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Survey Geographic (SSURGO) Database for [MLRA 141, Maine]. Available online. (accessed 14 Oct. 2021).

Sperduto, D.D. and William F. Nichols. 2011. Natural Communities of New Hampshire. 2nd Ed. NH Natural Heritage Bureau, Concord, NH. Pub. UNH Cooperative Extension, Durham, NH.

Swain, P. C. 2020. Classification of the Natural Communities of Massachusetts. Massachusetts Division of Fisheries and Wildlife, Westborough, MA

USNVC [United States National Vegetation Classification]. 2017. United States National Vegetation Classification Database V2.01. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC. Available The U.S. National Vegetation Classification (usnvc.org) (accessed 2 July. 2021).

Contributors

Christopher Mann

Approval

Nels Barrett, 9/27/2024

Acknowledgments

Nels Barrett and Nick Butler provided considerable review of this ecological site concept.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	06/29/2020
Approved by	Nels Barrett
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-