

Ecological site F231XY110AK Boreal Forest Gravelly Slopes Steep

Last updated: 2/13/2024
Accessed: 05/10/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 231X–Interior Alaska Highlands

The Interior Alaska Uplands (MLRA 231X) is in the Interior Region of Alaska and includes the extensive hills, mountains, and valleys between the Tanana River to the south and the Brooks Range to the north. These hills and mountains surround the Yukon Flats Lowlands (MLRA 232X). MLRA 231X makes up about 69,175 square miles. The hills and mountains of the area tend to be moderately steep to steep resulting in high-relief slopes. The mountains are generally rounded at lower elevations and sharp-ridged at higher elevations. Elevation ranges from about 400 feet in the west, along the boundary with the Interior Alaska Lowlands (MLRA 229X), to 6,583 feet at the summit of Mt. Harper, in the southeast. Major tributaries include large sections of the Yukon, Koyukuk, Kanuti, Charley, Coleen, and Chatanika Rivers. This area is traversed by several major roads, including the Taylor Highway in the east and the Steese, Elliott, and Dalton Highways north of Fairbanks. The area is mostly undeveloped wild land that is sparsely populated. The largest community along the road system is Fairbanks with smaller communities like Alatna, Allakaket, Chicken, Eagle, Eagle Village, Hughes, and Rampart occurring along the previously mentioned rivers and highways.

The vast majority of this MLRA was unglaciated during the Pleistocene epoch with the exceptions being the highest mountains and where glaciers extended into the area from the Brooks Range. For the most part, glacial moraines and drift are limited to the upper elevations of the highest mountains. Most of the landscape is mantled with bedrock colluvium originating from the underlying bedrock. Valley bottoms are filled with Holocene fluvial deposits and colluvium from the adjacent mountain slopes. Silty loess, which originated from unvegetated flood plains in and adjacent to this area, covers much of the surface. On hill and mountain slopes proximal to major river valleys (e.g., Tanana and Yukon Rivers), the loess is many feet thick. As elevation and distance from major river valleys increases, loess thickness decreases significantly. Bedrock is commonly exposed on the highest ridges.

This area is in the zone of discontinuous permafrost. Permafrost commonly is close to the surface in areas of the finer textured sediments throughout the MLRA. Isolated masses of ground ice occur in thick deposits of loess on terraces and the lower side slopes of hills. Solifluction lobes, frost boils, and circles and stripes are periglacial features common on mountain slopes in this area. Pingos, thermokarst pits and mounds, ice-wedge polygons, and earth hummocks are periglacial features common on terraces, lower slopes of hills and mountains, and in upland valleys in the area.

The dominant soil orders in this area are Gelisols, Inceptisols, Spodosols, and Entisols. The soils in the area have a subgelic or cryic soil temperature regime, an aquic or udic soil moisture regime, and mixed mineralogy. Gelisols are common on north facing slopes, south facing footslopes, valley bottoms, and stream terraces. Gelisols are typically shallow or moderately deep to permafrost (10 to 40 inches) and are poorly or very poorly drained. Wildfires can disturb the insulating organic material at the surface, lowering the permafrost layer, eliminating perched water tables from Gelisols, and thus changing the soil classification. Inceptisols and Spodosols commonly form on south facing hill and mountain slopes. Entisols are common on flood plains and high elevation mountain slopes. Miscellaneous (non-soil) areas make up about 2 percent of this MLRA. The most common miscellaneous areas are rock outcrop and rubble land. In many valleys placer mine tailings are common.

Short, warm summers and long, cold winters characterize the subarctic continental climate of the area. The mean annual temperature of the area ranges from 22 to 27 degrees F. The mean annual temperature of the southern half of the area is approximately 3 degrees warmer compared to the northern half (PRISM 2018). The warmest months span June through August with mean monthly temperatures ranging from 50 to 56 degrees F. The coldest months span November through February with mean monthly temperatures ranging from -5 to 3 degrees F. When compared to the high-elevation alpine and subalpine life zones, the lower elevation boreal life zone tends to be 2-3 degrees F colder during the coldest months and 1-2 degrees F warmer during the warmest months (PRISM 2018). The freeze-free period at the lower elevations averages about 60 to 100 days, and the temperature usually remains above freezing from June through mid-September.

Precipitation is limited across this area, with the average annual precipitation ranging from 12 to 19 inches. The southern half of the areas receives approximately 2.5 inches more annual precipitation than the northern half (PRISM 2018). The lower elevation boreal life zone receives approximately 2.5 inches less annual precipitation than the high-elevation alpine and subalpine life zones (PRISM 2018). Approximately 3/5th of the annual precipitation occurs during the months of June through September with thunderstorms being common. The average annual snowfall ranges from about 45 to 100 inches. The ground is consistently covered with snow from November through March.

Most of this area is forested below an elevation of about 2500 feet. Dominant tree species on slopes are white spruce and black spruce. Black spruce stands are most common on north-facing slopes, stream terraces, and other sites with poor drainage and permafrost. White spruce stands are most common on warm slopes with dry soils. At lower elevations, lightning-caused wildfires are common, often burning many thousands of acres during a single fire. Following wildfires, forbs, grasses, willow, ericaceous shrubs, paper birch, and quaking aspen communities are common until they are eventually replaced by stands of spruce. Tall willow and alder scrub is extensive on low flood plains. White spruce and balsam poplar are common on high flood plains.

With increasing elevation, the forests and woodlands give way to subalpine communities dominated by krummholz spruce, shrub birch, willow, and ericaceous shrubs. At even higher elevations, alpine communities prevail which are characterized by diverse forbs, dwarf ericaceous shrubs, and eightpetal mountain-avens. Many of these high elevation communities have a considerable amount of lichen cover and bare ground.

LRU notes

This area supports three life zones defined by the physiological limits of plant communities along an elevational gradient: boreal, subalpine, and alpine. The boreal life zone is the elevational band where forest communities dominate. Not all areas in the boreal life zone are forest communities, however, particularly in places with too wet or dry soil to support tree growth (e.g., bogs or river bluffs). Above the boreal band of elevation, subalpine and alpine vegetation dominate. The subalpine zone is typically a narrow transitional band between the boreal and the alpine life zones, and is characterized by sparse, stunted trees. In the subalpine, certain types of birch and willow shrub species grow at ≥ 1 m in height (commonly *Betula glandulosa* and *Salix pulchra*). In the alpine, trees no longer occur, and all shrubs are dwarf or lay prostrate on the ground. In this area, the boreal life zone occurs below 2500 feet elevation on average. The transition between boreal and alpine vegetation can occur within a range of elevations, and is highly dependent on slope, aspect, and shading from adjacent mountains.

Within each life zone, there are plant assemblages that are typically associated with cold slopes and warm slopes. Cold slopes and warm slopes are created by the combination of the steepness of the slope, the aspect, and shading from surrounding ridges and mountains. Warm slope positions typically occur on southeast to west facing slopes that are moderate to very steep ($>10\%$ slope) and are not shaded by the surrounding landscape. Cold slopes typically occur on northwest to east facing slopes, occur in shaded slope positions, or occur in low-lying areas that are cold air sinks. Examples of shaded positions include head slopes, low relief backslopes of hills, and the base of hills and mountains shaded by adjacent mountain peaks. Warm boreal slope soils have a cryic soil temperature regime and lack permafrost. In this area, white spruce forests are an indicator of warm boreal slopes. Cold boreal slope soils typically have a gelic soil temperature regime and commonly have permafrost. In this area, black spruce forests and woodlands are an indicator of cold boreal slopes. The boreal life zone can occur at higher elevations on warm slopes, and lower elevations on cold slopes.

Classification relationships

Landfire BPS – 7416061 – Western North American Boreal Dry Aspen-Steppe Bluff – Lower Elevations (Landfire 2009)

Ecological site concept

This boreal site occurs on warm hillslopes and escarpments that are very steep. Associated soils lack permafrost, do not have a seasonal water table, and are considered somewhat excessively drained. The soil profile has a thin layer of organic material over a layer of windblown silt or silty colluvium of variable thickness over gravelly residuum or colluvium. Soils with residuum commonly contact bedrock but at depths most commonly greater than 20 inches.

Multiple plant communities occur within the reference state and the vegetation in each community differs in large part due to fire. When the reference state vegetation burns, the post-fire plant community is dominantly forbs, grasses, and weedy mosses. With time and lack of another fire event, the post-fire vegetation goes through multiple stages of succession. For this site, the reference plant community is the most stable with the longest time since the vegetation was burned. This community is typically characterized as open needleleaf forest (Viereck et al. 1992) with white spruce as the dominant tree. For this ecological site to progress from the earliest stages of post-fire succession to the oldest stages of succession, data suggest that 150 years or more must elapse without another fire event (Foot 1982; Chapin et al. 2006; Landfire 2009).

The reference plant community understory commonly has twinflower, lingonberry, common juniper, kinnikinnick, bluejoint, purple reedgrass, false toadflax, arctic lupine, tall bluebells, sidebells wintergreen, splendid feathermoss, rhytidium moss, and Schreber's big red stem moss. White spruce tree cover is primarily in the tall tree stratum (greater than 40 feet in height). The understory vegetative strata that characterize this community phase are bryophytes and dwarf shrubs (less than 8 inches in height).

Associated sites

R231XY109AK	Boreal Scrub Gravelly Slopes Dry Occurs on the same escarpments but on steeper and more erosive slopes.
F231XY182AK	Boreal Forest Gravelly Slopes Occurs adjacent to this site but on less steep slopes.

Similar sites

R231XY109AK	Boreal Scrub Gravelly Slopes Dry This site occurs on steeper and more erosive slopes that do not support stands of trees.
F231XY182AK	Boreal Forest Gravelly Slopes Both sites occur on the same warm boreal slopes. The steeper slopes associated with site 110 have comparatively drier soils that result in different kinds and amounts of understory vegetation.
F231XY181AK	Boreal Forest Gravelly Slopes Steep Cold Site 181 occurs on cold boreal slopes that are very steep. These colder slopes have less productive white spruce stands and different kinds and amounts of understory vegetation.

Table 1. Dominant plant species

Tree	(1) <i>Picea glauca</i>
Shrub	(1) <i>Juniperus communis</i> (2) <i>Arctostaphylos uva-ursi</i>
Herbaceous	(1) <i>Hylocomium splendens</i> (2) <i>Rhytidium rugosum</i>

Physiographic features

This boreal site occurs on very steep and warm slopes of hills and escarpments. While backslopes are the most

common hillslope position, this site occasionally occurs on shoulders. On rare occasion, this site is associated with climbing dunes on hills. The boreal life zone typically occurs below 2500 feet but this site at times occurs on warmer hillslopes at 2900 feet or more elevation. These are very steep slopes commonly ranging between 45 and 85 percent or more and are southeast to northwest facing. This site does not flood or pond and does not have a water table in the soil profile. This site generates moderate amounts of runoff to adjacent, downslope ecological sites.

Table 2. Representative physiographic features

Hillslope profile	(1) Backslope (2) Shoulder
Landforms	(1) Hill (2) Escarpment (3) Climbing dune
Runoff class	Medium
Flooding frequency	None
Ponding frequency	None
Elevation	875–2,500 ft
Slope	45–85%
Water table depth	60 in
Aspect	W, NW, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	350–2,975 ft
Slope	20–90%
Water table depth	Not specified

Climatic features

Short, warm summers and long, cold winters characterize the subarctic continental climate associated with this boreal site. The mean annual temperature of the site ranges from 22 to 27 degrees F. The warmest months span June through August with mean normal maximum monthly temperatures ranging from 60 to 66 degrees F. The coldest months span November through February with mean normal minimum temperatures ranging from -3 to -12 degrees F. The freeze-free period for the site ranges from 80 to 120 days, and the temperature usually remains above freezing from late May through mid-September.

The area receives minimal annual precipitation with the summer months being the wettest. Average annual precipitation across the area typically ranges between 12 to 18 inches. Approximately 3/5th of the annual precipitation occurs during the months of June through September with thunderstorms common. The average annual snowfall ranges from about 45 to 100 inches. The ground is consistently covered with snow from November through March.

Table 4. Representative climatic features

Frost-free period (characteristic range)	16-78 days
Freeze-free period (characteristic range)	76-114 days
Precipitation total (characteristic range)	12-18 in
Frost-free period (actual range)	4-87 days
Freeze-free period (actual range)	48-120 days

Precipitation total (actual range)	9-20 in
Frost-free period (average)	53 days
Freeze-free period (average)	90 days
Precipitation total (average)	15 in




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) EAGLE AP [USW00026422], Tok, AK
- (2) CHICKEN [USC00501684], Tok, AK
- (3) MILE 42 STEESE [USC00505880], Fairbanks, AK
- (4) BETTLES AP [USW00026533], Bettles Field, AK
- (5) CIRCLE HOT SPRINGS [USC00501987], Central, AK
- (6) FT KNOX MINE [USC00503160], Fairbanks, AK
- (7) GILMORE CREEK [USC00503275], Fairbanks, AK
- (8) FOX 2SE [USC00503181], Fairbanks, AK
- (9) ESTER DOME [USC00502868], Fairbanks, AK
- (10) ESTER 5NE [USC00502871], Fairbanks, AK
- (11) COLLEGE 5 NW [USC00502112], Fairbanks, AK
- (12) COLLEGE OBSY [USC00502107], Fairbanks, AK
- (13) KEYSTONE RIDGE [USC00504621], Fairbanks, AK

Influencing water features

Due to its landscape position, this site is neither associated with or influenced by streams or wetlands. Precipitation

and throughflow are the main sources of water for this ecological site.

Wetland description

n/a

Soil features

Soils commonly formed in windblown silts over residuum or gravelly colluvium and do not have permafrost. Rock fragments do not occur on the soil surface. These are mineral soils commonly capped with 1 to 3 inches of organic material. The mineral soil below the organic material is a silt loam formed from wind-blown loess, which lacks rock fragments and has high water holding capacity. The thickness of the loess layer is highly variable ranging from 4 to 25 inches or more. Below the silty loess, the soil parent material is most commonly residuum or colluvium with rock fragments ranging between 40 and 80 percent of the soil profile by volume. These are typically very deep soils. At times, soils with residuum contact bedrock at shallow to moderate depths (19 to 33 inches). In rare instances, there is an abrupt change between the silty loess and gravelly colluvium resulting in restrictions at very shallow depth (4 to 7 inches). The pH of the soil profile often ranges from moderately acidic to neutral but on escarpment can be moderately alkaline. The soils are dry, lack a seasonal water table, and are generally somewhat excessively drained.



Figure 7. A typical soil profile associated with this site.

Table 5. Representative soil features

Parent material	(1) Loess (2) Colluvium (3) Residuum
Surface texture	(1) Silt loam (2) Stony silt loam
Family particle size	(1) Coarse-loamy (2) Loamy-skeletal (3) Coarse-silty
Drainage class	Somewhat excessively drained
Permeability class	Moderately rapid
Depth to restrictive layer	Not specified
Soil depth	60 in
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	2.3–5 in

Calcium carbonate equivalent (10-40in)	0–1%
Clay content (0-20in)	5–10%
Electrical conductivity (10-40in)	0–2 mmhos/cm
Sodium adsorption ratio (10-40in)	0
Soil reaction (1:1 water) (10-40in)	5.6–7.3
Subsurface fragment volume ≤3" (0-60in)	30–55%
Subsurface fragment volume >3" (0-60in)	10–25%

Table 6. Representative soil features (actual values)

Drainage class	Well drained to somewhat excessively drained
Permeability class	Moderately rapid to rapid
Depth to restrictive layer	19 in
Soil depth	19 in
Surface fragment cover ≤3"	0–1%
Surface fragment cover >3"	0–19%
Available water capacity (0-40in)	1.4–8.7 in
Calcium carbonate equivalent (10-40in)	0–5%
Clay content (0-20in)	Not specified
Electrical conductivity (10-40in)	0–5 mmhos/cm
Sodium adsorption ratio (10-40in)	Not specified
Soil reaction (1:1 water) (10-40in)	3.9–8.1
Subsurface fragment volume ≤3" (0-60in)	0–55%
Subsurface fragment volume >3" (0-60in)	0–27%

Ecological dynamics

Fire

In the Interior Alaska Uplands area, fire is a common and natural event that has a significant control on the vegetation dynamics across the landscape. A typical fire event in the lands associated with this ecological site will reset plant succession and alter dynamic soil properties (e.g., thickness of the organic material). For this ecological site to progress from the earliest stages of post-fire succession dominated by grasses and forbs to the oldest stages of succession dominated by white spruce forests, data suggest that 150 years or more must elapse without another fire event (Foot 1982; Chapin et al. 2006; Landfire 2009).

Within this area, fire is considered a natural and common event that typically is unmanaged. Fire suppression is

limited, and generally occurs adjacent to Fairbanks and the various villages spread throughout the area or on allotments with known structures, all of which have a relatively limited acre footprint. Most fires are caused by lightning strikes. From 2000 to 2020, 596 known fire events occurred in the Interior Alaska Uplands area and the burn perimeter of the fires totaled about 13.8 million acres (AICC 2022). Fire-related disturbances are highly patchy and can leave undisturbed areas within the burn perimeter. During this time frame, 80% of the fire events were smaller than 20,000 acres but 18 fire events were greater than 200,000 acres in size (AICC 2022). These burn perimeters cover approximately 30% of the Interior Alaska Uplands area over a period of 20 years.

The fire regime within Interior Alaska follows two basic scenarios—low-severity burns and high-severity burns. It should be noted, however, that the fire regime in Interior Alaska is generally thought to be much more complex (Johnstone et al. 2008). Burn severity refers to the proportion of the vegetative canopy and organic material consumed in a fire event (Chapin et al. 2006). Fires in cool and moist habitat tend to result in low-severity burns, while fires in warm and dry habitat tend to result in high-severity burns. Because the soils have a thin organic cap and are well drained, the typical fire scenario for this ecological site is considered to result in a high-severity burn.

Large portions of the organic mat are consumed during a high-severity fire event, commonly exposing pockets of mineral soil. The loss of this organic mat, which insulates the mineral soil, and the decrease in site albedo tends to cause overall soil temperatures to increase (Hinzman et al. 2006). These alterations to soil temperature may result in increased depths of seasonal frost in the soil profile. High-severity fire events also destroy a majority of the vascular and nonvascular biomass above ground.


Field data from this site suggest that each of the forested community phases will burn and that fire events will cause a transition to the pioneering stage of fire succession. This stage (community phase 1.5) is a mix of species that either regenerate in place (e.g., subterranean root crowns for willow and rhizomes for graminoids) and/or from wind-dispersed seed or spores that colonize exposed mineral soil (e.g., quaking aspen [*Populus tremuloides*] and *Ceratodon* moss [*Ceratodon purpureus*]). The pioneering stage of fire succession is primarily composed of tree seedlings, forbs, grasses, and weedy bryophytes. This stage of succession is thought to persist for up to 10 years post-fire. Willow (*Salix* spp.) and quick growing deciduous tree seedlings continue to colonize and grow in stature on recently burned sites until they become dominant in the overstory, which marks the transition to the early stage of fire succession (community phase 1.4). This early stage of fire succession is thought to persist 10 to 30 years post-fire. In the absence of fire, tree species continue to become more dominant in the stand and eventually develop into forests.

The later stages of succession have an overstory that is dominantly deciduous trees (community phase 1.3), a mix of broadleaf and needleleaf trees (community phase 1.2), or needleleaf trees (community phase 1.1). The recruitment of trees species during the pioneering and early stages of post-fire succession largely controls the composition of the stand of trees in the later stages of post-fire succession (Johnstone et al. 2010a). During these later stages of succession, the slower growing white spruce seedlings mature and eventually replace the shade-intolerant broadleaf tree species. The typical fire return interval for white spruce stands in Interior Alaska is 150 years (Landfire 2009; Abrahamson 2014).


Lands associated with this site may be burning more frequently than in the past, which may result in alternative pathways of succession. The historic fire return interval for white spruce stands in Interior Alaska occurs approximately once 150 years (Landfire 2009; Abrahamson 2014). Due to global climate change, stands of spruce in certain portions of the Alaskan boreal forest are burning more frequently than these historic averages (Kelly et al. 2013). Increases to burn frequency favors forested stands dominated by quick growing deciduous trees (community 1.3). A major reason being that increased fire frequency decreases the presence and abundance of mature, cone-bearing trees. Less mature trees result in less spruce seedlings post-fire and an overall decreased abundance of spruce in the developing forest canopy. Increased burn frequency in the boreal forest may result in alternative pathways of post-fire succession with stands of deciduous trees persisting for longer than normal durations of time (Johnstone et al. 2010b).

State and transition model


Ecosystem states



State 1 submodel, plant communities



Communities 1, 5 and 2 (additional pathways)



- 1.1a - A high-severity fire sweeps through and incinerates much of the above ground vegetation.
- 1.2b - Time without fire.
- 1.2a - A high-severity fire sweeps through and incinerates much of the above ground vegetation.
- 1.3b - Time without fire.
- 1.3a - A high-severity fire sweeps through and incinerates much of the above ground vegetation.
- 1.4b - Time without fire.
- 1.4a - A high-severity fire sweeps through and incinerates much of the above ground vegetation.
- 1.5a - Time without fire.

State 1
Reference State



Figure 8. Warm boreal slopes that are very steep in the Interior Alaska Uplands.

The reference plant community is open needleleaf forest (Viereck et al. 1992) with the dominant tree being white spruce. There are five community phases within the reference state related to fire.

Dominant plant species

- white spruce (*Picea glauca*), tree
- common juniper (*Juniperus communis*), shrub
- kinnikinnick (*Arctostaphylos uva-ursi*), shrub
- splendid feather moss (*Hylocomium splendens*), other herbaceous
- rhytidium moss (*Rhytidium rugosum*), other herbaceous

Community 1.1

White Spruce / common juniper - kinnikinnick / splendid feather moss - rhytidium moss



Figure 9. Typical plant community associated with community phase 1.1.

The reference plant community is characterized as an open needleleaf forest (25 to 60 percent cover) composed primarily of mature white spruce. White spruce tree cover is primarily in the tall tree stratum (greater than 40 feet in height). Gaps occur in the tree canopy, but they are limited in size and extent and are likely the result of occasional windthrow. Live deciduous trees, primarily quaking aspen, occasionally occur in the tree canopy, but most have been replaced by white spruce. The soil surface is primarily covered with bryophytes. Common and abundant understory species for this community include twinflower, lingonberry, common juniper, kinnikinnick, bluejoint, purple reedgrass, false toadflax, arctic lupine, tall bluebells, sidebells wintergreen, splendid feathermoss, rhytidium moss, and Schreber's big red stem moss. The understory vegetative strata that characterize this community phase are bryophytes and dwarf shrubs (less than 8 inches in height).

Dominant plant species

- white spruce (*Picea glauca*), tree
- twinflower (*Linnaea borealis*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- common juniper (*Juniperus communis*), shrub
- kinnikinnick (*Arctostaphylos uva-ursi*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- purple reedgrass (*Calamagrostis purpurascens*), grass
- splendid feather moss (*Hylocomium splendens*), other herbaceous
- rhytidium moss (*Rhytidium rugosum*), other herbaceous
- Schreber's big red stem moss (*Pleurozium schreberi*), other herbaceous
- false toadflax (*Geocaulon lividum*), other herbaceous
- arctic lupine (*Lupinus arcticus*), other herbaceous
- tall bluebells (*Mertensia paniculata*), other herbaceous
- sidebells wintergreen (*Orthilia secunda*), other herbaceous

Community 1.2

white spruce - quaking aspen / prickly rose - kinnikinnick / false toadflax - tall bluebells



Figure 10. Typical plant community associated with community phase 1.2.

Community 1.2 is in the late stage of fire-induced secondary succession for this ecological site. It is characterized as open mixed forest (Viereck et al. 1992) with mature aspen and a mixture of immature and mature white spruce as the dominant trees. Tree cover is generally split between immature medium-sized trees (15 to 40 feet in height) and mature tall trees (greater than 40 feet in height). The soil surface is primarily covered with herbaceous litter. Common and abundant understory species for this community include prickly rose, russet buffaloberry, lingonberry, twinflower, squashberry, common juniper, kinnikinnick, bluejoint, false toadflax, tall bluebells, sidebells wintergreen, fireweed, northern bedstraw, and splendid feathermoss. The understory vegetative strata that characterize this community phase are low shrubs (between 8 and 36 inches tall), dwarf shrubs (less than 8 inches), and medium forbs (between 4 and 24 inches tall).

Dominant plant species

- white spruce (*Picea glauca*), tree
- quaking aspen (*Populus tremuloides*), tree
- prickly rose (*Rosa acicularis*), shrub
- russet buffaloberry (*Shepherdia canadensis*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- twinflower (*Linnaea borealis*), shrub
- squashberry (*Viburnum edule*), shrub
- common juniper (*Juniperus communis*), shrub
- kinnikinnick (*Arctostaphylos uva-ursi*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- false toadflax (*Geocaulon lividum*), other herbaceous
- fireweed (*Chamerion angustifolium*), other herbaceous
- splendid feather moss (*Hylocomium splendens*), other herbaceous
- tall bluebells (*Mertensia paniculata*), other herbaceous
- sidebells wintergreen (*Orthilia secunda*), other herbaceous
- northern bedstraw (*Galium boreale*), other herbaceous

Community 1.3

quaking aspen / prickly rose - kinnikinnick / fireweed - false toadflax



Figure 11. Typical plant community associated with community phase 1.3.

Community 1.3 is in the middle stage of fire-induced secondary succession for this ecological site. It is characterized as open deciduous forest (Viereck et al. 1992) with mature stands of quaking aspen. Immature white spruce are a common subdominant tree in the canopy. Tree cover primarily occurs as medium-sized trees (15 to 40 feet in height). The soil surface is primarily covered with herbaceous litter. Common and abundant understory species for this community include prickly rose, russet buffaloberry, kinnikinnick, lingonberry, twinflower, squashberry, bluejoint, fireweed, northern bedstraw, and tall bluebells. The understory vegetative strata that characterize this community phase are low shrubs (between 8 and 36 inches tall), dwarf shrubs (less than 8 inches), and medium forbs (between 4 and 24 inches tall).

Dominant plant species

- quaking aspen (*Populus tremuloides*), tree
- prickly rose (*Rosa acicularis*), shrub
- russet buffaloberry (*Shepherdia canadensis*), shrub
- kinnikinnick (*Arctostaphylos uva-ursi*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- squashberry (*Viburnum edule*), shrub
- twinflower (*Linnaea borealis*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- fireweed (*Chamerion angustifolium*), other herbaceous
- tall bluebells (*Mertensia paniculata*), other herbaceous
- northern bedstraw (*Galium boreale*), other herbaceous

Community 1.4

quaking aspen / prickly rose - russet buffaloberry / bluejoint



Figure 12. Typical plant community associated with community phase 1.4.

Community phase 1.4 is in the early stage of fire-induced secondary succession for this ecological site. It is characterized as open tall scrubland. The overstory canopy is primarily composed of broadleaf tree species commonly being quaking aspen and tall scrub commonly being willow. Tree cover primarily is in the regenerative tree stratum (less than 15 feet in height). White spruce seedlings are common in the understory, but they are not abundant. The soil surface is primarily covered with herbaceous litter and woody debris. Common and abundant understory species for this community include prickly rose, russet buffaloberry, kinnikinnick, lingonberry, twinflower, squashberry, bluejoint, purple reedgrass, fireweed, northern bedstraw, tall bluebells, and weedy bryophytes. The understory vegetative strata that characterize this community phase are low shrubs (between 8 and 36 inches tall), mosses, and medium forbs (between 4 and 24 inches tall).

Dominant plant species

- quaking aspen (*Populus tremuloides*), tree
- prickly rose (*Rosa acicularis*), shrub
- russet buffaloberry (*Shepherdia canadensis*), shrub
- kinnikinnick (*Arctostaphylos uva-ursi*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- Bebb willow (*Salix bebbiana*), shrub
- purple reedgrass (*Calamagrostis purpurascens*), grass
- bluejoint (*Calamagrostis canadensis*), grass
- fireweed (*Chamerion angustifolium*), other herbaceous
- pohlia moss (*Pohlia nutans*), other herbaceous
- juniper polytrichum moss (*Polytrichum juniperinum*), other herbaceous
- common yarrow (*Achillea millefolium*), other herbaceous
- Rocky Mountain goldenrod (*Solidago multiradiata*), other herbaceous
- tall bluebells (*Mertensia paniculata*), other herbaceous
- Alaska wild rhubarb (*Polygonum alpinum*), other herbaceous
- three toothed saxifrage (*Saxifraga tricuspidata*), other herbaceous

- dwarf scouringrush (*Equisetum scirpoides*), other herbaceous

Community 1.5

bluejoint / fireweed



Figure 13. Typical plant community associated with community phase 1.5.

Community phase 1.5 is in the pioneering stage of fire-induced secondary succession for this ecological site. It is characterized as a mesic forb herbaceous community. Tree cover primarily is in the regenerative tree stratum (less than 15 feet in height). Deciduous tree seedlings, primarily quaking aspen, are common throughout the community. Although small areas of exposed bare soil are common, the soil surface is primarily covered with a mixture of weedy bryophyte species, woody debris, and herbaceous litter. Commonly observed species include an assortment of willow, kinnikinnick, prickly rose, purple reedgrass, bluejoint, Altai fescue, northern bedstraw, fireweed, alpine sweetvetch, rocky mountain goldenrod, and common yarrow. The vegetative strata that characterize this community phase are tree regeneration (less than 15 feet in height), medium forbs (between 4 inches and 2 feet in height), and medium graminoids (between 4 inches and 2 feet in height).

Dominant plant species

- willow (*Salix*), shrub
- kinnikinnick (*Arctostaphylos uva-ursi*), shrub
- prickly rose (*Rosa acicularis*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- Altai fescue (*Festuca altaica*), grass
- purple reedgrass (*Calamagrostis purpurascens*), grass
- pohlia moss (*Pohlia nutans*), other herbaceous
- ceratodon moss (*Ceratodon purpureus*), other herbaceous
- juniper polytrichum moss (*Polytrichum juniperinum*), other herbaceous
- northern bedstraw (*Galium boreale*), other herbaceous
- alpine sweetvetch (*Hedysarum alpinum*), other herbaceous
- Rocky Mountain goldenrod (*Solidago multiradiata*), other herbaceous
- common yarrow (*Achillea millefolium*), other herbaceous
- American thorn wax (*Bupleurum americanum*), other herbaceous
- Jakutsk snowparsley (*Cnidium cniidiifolium*), other herbaceous
- eastern pasqueflower (*Pulsatilla patens*), other herbaceous

Pathway 1.1a

Community 1.1 to 1.5



White Spruce / common juniper - kinnikinnick / splendid feather moss - rhytidium moss



bluejoint / fireweed

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated dry soils, this site commonly experiences high-severity fires. A significant proportion of organic matter is consumed, leaving exposed mineral soil. Vegetation usually resprouts from surviving individuals or is recruited from nearby areas via seed or seedbank.

Pathway 1.2b Community 1.2 to 1.1



white spruce - quaking aspen / prickly rose - kinnikinnick / false toadflax - tall bluebells



White Spruce / common juniper - kinnikinnick / splendid feather moss - rhytidium moss

Time without fire results in the continued growth and increased abundance of white spruce, which overtop and remove the shade intolerant deciduous tree species from the forest canopy.

Pathway 1.2a Community 1.2 to 1.5



white spruce - quaking aspen / prickly rose - kinnikinnick / false toadflax - tall bluebells



bluejoint / fireweed

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated dry soils, this site commonly experiences high-severity fires. A significant proportion of organic matter is consumed, leaving exposed mineral soil. Vegetation usually resprouts from surviving individuals or is recruited from nearby areas via seed or seedbank.

Pathway 1.3b Community 1.3 to 1.2



quaking aspen / prickly rose - kinnikinnick / fireweed - false toadflax




white spruce - quaking aspen / prickly rose - kinnikinnick / false toadflax - tall bluebells

Time without fire results in the continued growth and increased abundance of white spruce, which overtop and remove the shade intolerant deciduous tree species from the forest canopy.


Pathway 1.3a

Community 1.3 to 1.5




A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated dry soils, this site commonly experiences high-severity fires. A significant proportion of organic matter is consumed, leaving exposed mineral soil. Vegetation usually resprouts from surviving individuals or is recruited from nearby areas via seed or seedbank.

Pathway 1.4b
Community 1.4 to 1.3




Time without fire results in the continued development of a forest canopy dominated by quaking aspen.

Pathway 1.4a
Community 1.4 to 1.5



A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated dry soils, this site commonly experiences high-severity fires. A significant proportion of organic matter is consumed, leaving exposed mineral soil. Vegetation usually resprouts from surviving individuals or is recruited from nearby areas via seed or seedbank.

Pathway 1.5a
Community 1.5 to 1.4



Time without fire results in the herbaceous community being overtopped by willow and deciduous tree seedlings.

Additional community tables

Table 7. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)	Diameter (In)	Basal Area (Square Ft/Acre)
Tree							
white spruce	PIGL	<i>Picea glauca</i>	Native	31–55	55–65	6.5–13.5	–

Table 8. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
bluejoint	CACA4	<i>Calamagrostis canadensis</i>	Native	0.3–2	0–3
purple reedgrass	CAPU	<i>Calamagrostis purpurascens</i>	Native	0.3–2	0–2
Forb/Herb					
false toadflax	GELI2	<i>Geocaulon lividum</i>	Native	0.3–2	0–5
tall bluebells	MEPA	<i>Mertensia paniculata</i>	Native	0.3–2	0–2
arctic lupine	LUAR2	<i>Lupinus arcticus</i>	Native	0.3–2	0–1
sidebells wintergreen	ORSE	<i>Orthilia secunda</i>	Native	0.3–2	0–1
bluntleaf sandwort	MOLA6	<i>Moehringia lateriflora</i>	Native	0.3–2	0–0.1
northern bedstraw	GABO2	<i>Galium boreale</i>	Native	0.3–2	0–0.1
alpine sweetvetch	HEAL	<i>Hedysarum alpinum</i>	Native	0.3–2	0–0.1
Shrub/Subshrub					
lingonberry	VAVI	<i>Vaccinium vitis-idaea</i>	Native	0–0.6	0–15
twinflower	LIBO3	<i>Linnaea borealis</i>	Native	0.1–0.6	0.1–12
common juniper	JUCO6	<i>Juniperus communis</i>	Native	0.6–3	0–5
prickly rose	ROAC	<i>Rosa acicularis</i>	Native	0.6–3	0–5
kinnikinnick	ARUV	<i>Arctostaphylos uva-ursi</i>	Native	0.1–0.6	0–0.1
Nonvascular					
Schreber's big red stem moss	PLSC70	<i>Pleurozium schreberi</i>	Native	0–0.1	0–80
splendid feather moss	HYSP70	<i>Hylocomium splendens</i>	Native	0.1–0.2	1–75
rhytidium moss	RHRU70	<i>Rhytidium rugosum</i>	Native	0.1–0.2	0–25
felt lichen	PEAP60	<i>Peltigera aphthosa</i>	Native	0.1–0.2	0–3

Animal community

n/a

Hydrological functions

n/a

Recreational uses

n/a

Wood products

n/a

Other products

n/a

Other information

n/a

Inventory data references

Tier 2 sampling plots used to develop the reference state. Plot numbers as recorded in NASIS with associated community phase.

Community 1.1

11BB00501, 12AK2902008, 12SN02902, 13BA00204, 2016AK290638

Community 1.2

08CS00604, 08CS00902, 08TC00201, 09NP00304, 10TC02102

Community 1.3

08CS01003, 10TC03801, 10TC04503, 11MC01901, 14EG00202

Community 1.4

10NP03302, 10TC02401

Community 1.5

08TC01304, 09NP00302, 09NP01301

References

Abrahamson, I.L. 2014. Fire Regimes of Alaskan White Spruce Communities.

Chapin, F.S., L.A. Viereck, P.C. Adams, K.V. Cleve, C.L. Fastie, R.A. Ott, D. Mann, and J.F. Johnstone. 2006. Successional processes in the Alaskan boreal forest. Page 100 in *Alaska's changing boreal forest*. Oxford University Press.

Foote, M.J. 1983. Classification, description, and dynamics of plant communities after fire in the taiga of interior Alaska. US Department of Agriculture, Forest Service, Pacific Northwest Forest and

Hinzman, L.D., L.A. Viereck, P.C. Adams, V.E. Romanovsky, and K. Yoshikawa. 2006. Climate and permafrost dynamics of the Alaskan boreal forest. *Alaska's changing boreal forest* 39–61.

Johnstone, J.F., F.S. Chapin, T.N. Hollingsworth, M.C. Mack, V. Romanovsky, and M. Turetsky. 2010. Fire, climate change, and forest resilience in interior Alaska. *Canadian Journal of Forest Research* 40:1302–1312.

Johnstone, J.F. 2008. A key for predicting postfire successional trajectories in black spruce stands of interior Alaska. US Department of Agriculture, Forest Service, Pacific Northwest Research Station.

Kelly, R., M.L. Chipman, P.E. Higuera, I. Stefanova, L.B. Brubaker, and F.S. Hu. 2013. Recent burning of boreal forests exceeds fire regime limits of the past 10,000 years. *Proceedings of the National Academy of Sciences* 110:13055–13060.

Schoeneberger, P.J. and D.A. Wysocki. 2012. Geomorphic Description System. Natural Resources Conservation Service, 4.2 edition. National Soil Survey Center, Lincoln, NE.

United States Department of Agriculture, . 2022. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin.

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

Other references

Alaska Interagency Coordination Center (AICC). 2022. <http://fire.ak.blm.gov/>

LANDFIRE. 2009. Western North American Boreal White Spruce-Hardwood Forest. In: LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior. Washington, DC.

PRISM Climate Group. 2018. Alaska – average monthly and annual precipitation and minimum, maximum, and mean temperature for the period 1981-2010. Oregon State University, Corvallis, Oregon.
<https://prism.oregonstate.edu/projects/alaska.php>. (Accessed 4 September 2019).

United States Department of Agriculture-Natural Resources Conservation Service. 2016. U.S. General Soil Map (STATSGO2). Web Soil Survey. Available online at <http://websoilsurvey.nrcs.usda.gov>. Accessed (Accessed 3 March 2021).

Contributors

Blaine Spellman
Jamin Johanson
Stephanie Shoemaker
Philip Barber

Approval

Kirt Walstad, 2/13/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/10/2025
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

-
2. **Presence of water flow patterns:**
-
3. **Number and height of erosional pedestals or terracettes:**
-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
-
5. **Number of gullies and erosion associated with gullies:**
-
6. **Extent of wind scoured, blowouts and/or depositional areas:**
-
7. **Amount of litter movement (describe size and distance expected to travel):**
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or**

decadence):

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-