

Ecological site R239XY051AK

Arctic Shrub Loamy Frozen Tidal Marsh

Last updated: 2/18/2025
Accessed: 05/11/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 239X–Northern Bering Sea Islands

The Northern Bering Sea Islands (MLRA 239X) occurs in Western Alaska and includes Saint Lawrence (1,792 square miles), Nunivak (1,632 square miles), and Saint Matthew (137 square miles) Islands and several smaller adjacent islands all of which are surrounded by the Bering Sea. This MLRA makes up 3,705 square miles. The terrain primarily consists of nearly level to rolling plains and highlands with mostly gentle slopes. Coastal lowlands dotted with numerous small- and medium-size lakes make up a significant part of St. Lawrence Island. Steep, low-relief volcanic cones, vents, and lava flows are common throughout Nunivak Island and less common on St. Lawrence and St. Matthew Islands. Narrow, discontinuous sea cliffs, sand dunes, and sand sheets are along many stretches of the coast. Elevation ranges from sea level along the coast to 2,207 feet at the summit of Atuk Mountain, on St. Lawrence Island. The area is mostly undeveloped wild land that is sparsely populated. Residents use this remote area primarily for subsistence hunting, fishing, and gathering. Reindeer and/or muskox herding provides meat and other products to residents on Nunivak Island and St. Lawrence Islands. The largest communities on the islands are Diomedes, Gambell, Mekoryuk, and Savoonga.

Geology and Soils

Across the islands, most of the landscape is mantled with late Tertiary and Quaternary alluvial, marine, and eolian surficial deposits. While a small portion of the northwest coast of St. Lawrence Island was glaciated (Patton et al. 2011), the vast majority of the MLRA was unglaciated during the Pleistocene Epoch. St. Lawrence Island is the most geologically complex of the islands in this area. The St. Lawrence Island coastal plain is dotted with numerous small- and medium-size lakes with a mosaic primarily composed of surficial deposits and volcanic and sedimentary rock, including coal beds and limestone. The highlands on this island are primarily composed of Cretaceous granitic bedrock except for Atuk Mountain which is composed of young volcanic bedrock from the Quaternary to late Tertiary. Nunivak and St. Matthew Islands are made up almost exclusively of early and late Tertiary and Quaternary volcanic rocks.

These islands are in the zone of discontinuous permafrost. Frozen soils are common across the vast extents of rolling plains and gentle sloping highlands. In these areas, the layer of permafrost is generally thin or moderately thick and occurs primarily in fine textured deposits. Permafrost generally does not occur on flood plains, in coarse textured sediments on the slopes of volcanic cones and other highlands, along the coast, or near lakes and other bodies of water. Common periglacial features include solifluction lobes, frost boils, and palsens (Swanson et al. 1986, USDA 2022).

The majority of soils are acidic, and the dominant soil order is Gelisols. Except for some non-acidic uplands on St. Lawrence Island, the vast majority of soil substrate across the MLRA is acidic (pH less than 5.5) (CAVM Team 2023). The Gelisols are shallow or moderately deep to permafrost (10 to 40 inches) and are typically very poorly to poorly drained. Common Gelisol suborders are Histels, Orthels, and Turbels. The Histels have thick accumulations of surface organic material and primarily occur in very wet coastal plain depressions and low-gradient drainageways. The Orthels and Turbels have comparably thinner surface organic material and primarily occur on

the coastal lowlands and other areas with gentle slopes. The MLRA also has small areas of Andisols, Entisols, Inceptisols, and Mollisols. Andisols and Inceptisols primarily occur on volcanic cones and other slopes with coarse textured, acidic soils. Mollisols occur on areas with limestone on St. Lawrence Island (USDA 2022). Entisols primarily occur on flood plains and estuaries. Miscellaneous (non-soil) areas make up about 10 percent of the area and are primarily water, lava flows, rubble composed of volcanic rock, and beach sediments.

Climate

The presence of sea ice in the Bering Sea strongly influences the climate across the islands in this area. Sea ice in the Bering Sea historically forms in early December, increases in thickness until late April, and breaks apart in June (Zuesler 1941). When sea ice is absent, the Bering Sea and North Pacific Ocean moderate diurnal and monthly temperatures resulting in a maritime climate. As sea ice forms around the islands, temperatures decrease significantly with the area shifting to a continental climate.

Vegetation

Tidal flats and estuaries support sedge dominant communities, while drier beach dune communities support American dunegrass and seacoast angelica communities (Swanson et al. 1986). The coastal lowlands and nearly level to rolling plains have a mosaic of sedge and moss dominant wetlands and various tundra. The tundra often has dwarf shrubs like crowberry; tussock forming and non-tussock forming sedges; and a variety of forbs, lichen, and mosses. Very wet drainages and the shores of lakes support wet sedge meadows. Drier soils on flood plains commonly support low to tall willow scrub with dense grasses and forbs in the understory. Shallow soils with coarse textured rocks common on volcanic cones, mountain slopes, and ridges commonly support alpine dwarf scrub dominated by ericaceous shrubs, *Dryas*, and dwarf willows. These communities commonly have a considerable amount of lichen and bare ground. Bedrock exposures and barrens with lichens and scattered shrubs and herbs in pockets of fine earth dominate the highest elevations, ridges, and other windblown sites.

Introduced ungulates

Introduced herds of reindeer and muskox provide a rich history of land use across the Northern Bering Sea Islands MLRA. Of the many islands in this MLRA, Nunivak was the only island historically grazed by ungulates. Inhabited by caribou until the late 1800's, the caribou on Nunivak Island were extirpated with the introduction of rifles (Griffin 2001).

Reindeer were introduced to St. Lawrence Island as early as 1901 (Jackson 1902), Nunivak Island in 1920, and St. Matthew Island in 1944 (Swanson and Barker 1991). Muskox were introduced to Nunivak Island in 1930 (ADFG 2024). Nunivak Island currently has managed herds of reindeer and muskox, St. Lawrence Island currently has managed herds of reindeer, and St. Matthew Island currently has no herds of reindeer. Some small islands in this MLRA are believed to have no history of natural or introduced ungulate herds (e.g. Pinnacle Islands, Hall Island, and Punuk Islands).

LRU notes

There are two distinct bioclimate in this MLRA resulting in slight differences in vegetation. St. Lawrence Island is more than 200 miles North of Nunivak and St. Matthew Islands. As a result, St. Lawrence Island is significantly colder. Mean annual air temperatures on Nunivak and St. Matthew Islands typically range from 30 to 34 degrees Fahrenheit and are between 24 to 28 degrees Fahrenheit on St. Lawrence Islands (PRISM 2018). More southerly islands in this area fall into the Circumpolar Arctic Vegetation Mapping (CAVM) subzone E and more northerly islands fall into CAVM subzone D (CAVM 2022). Moist and dry tundra common to the near level to rolling plains across the islands are thought to support plant communities with similar species but have different plant community structures. Subzone E supports low shrub communities and subzone D erect dwarf shrub communities (CAVM 2022). At this time, these differences in community structure are recognized but unique ecological sites for each CAVM bioclimate subzone were not developed.

This area supports two life zones defined by the physiological limits of plant communities along an elevational gradient: arctic and alpine. In this MLRA, the arctic life zone occurs below 500 feet elevation on average (Swanson et al. 1986) and is the elevational band where lowland vegetation dominates. For this MLRA, certain vascular plant species are common in the lowlands and much less common in the alpine (i.e. *Salix pulchra*, *Salix fuscescens*, *Betula nana*, *Ledum palustre* ssp. *decumbens*, and *Calamagrostis canadensis*). Above the arctic band of elevation,

alpine vegetation dominates. For this MLRA, certain vascular plant and lichen species are common in the alpine and much less common in the lowlands (i.e. *Dryas octopetala* ssp. *octopetala*, *Diapensia lapponica* var. *obovata*, *Anthoxanthum monticola* ssp. *alpinum*, *Oxytropis nigrescens*, *Alectoria ochroleuca*, and *Flavocetraria nivalis*). The lowlands also have much higher potential for lichen biomass yields compared to the alpine (Swanson et al. 1986). The transition between arctic and alpine vegetation can occur within a range of elevations, and is highly dependent on latitude, slope, aspect, and shading from adjacent mountains.

Classification relationships

Landfire BPS – 7617010 - Alaska Arctic Coastal Sedge-Dwarf-Shrubland (Landfire 2009)

Halophytic Sedge Wet Meadow (Viereck et al. 1992)

Marsh (Tidal) (Swanson et al. 1986)

Ecological site concept

This arctic ecological site occurs on portions of the coastal plain that are frequently inundated with tidal water and have wet, loamy soils with permafrost. Associated soils flood frequently for brief durations, pond frequently for long durations, have a high-water table throughout the growing season, and are considered very poorly drained. A typical soil profile has 7 to 11 inches of peat over loamy marine deposits.

The presence of introduced ungulate herds on Nunivak, St Lawrence, and St. Matthews Islands, in some places for over a century, plays an integral role in shaping vegetation across this MLRA. Islands in this MLRA without a history of introduced reindeer and muskox herds are associated with reference state vegetation, while islands with introduced herds are associated with grazing state vegetation.

One plant community has been documented within the grazing state for this ecological site. This community is characterized as halophytic sedge wet meadow (Viereck et al. 1992) with the dominant plants being oval-leaf willow, crowberry, water sedge, and lesser saltmarsh sedge. Other commonly observed species include red fescue, arctic daisy, purple marshlocks, pacific silverweed, saltmarsh starwort, Schreber's big red stem moss, and *Sphagnum* moss. The vegetative strata that characterize this community are dwarf shrubs (less than 8 inches height), medium graminoids (between 4 and 24 inches in height) and moss.

Associated sites

R239XY053AK	Arctic Sedge Loamy Tidal Marsh Occurs on estuaries and portions of the coastal plain that are lower and closer to the ocean. When compared to ecological site 51, this ecological site experiences more frequent and longer duration tidal flooding and soils have much higher sodium concentrations.
-------------	---

Similar sites

R239XY053AK	Arctic Sedge Loamy Tidal Marsh While ecological sites 51 and 53 are characterized as halophytic sedge wet meadow (Viereck et al. 1992) there are differences in the kinds and amounts of dominant vegetation. Site 53 has different sedges and lacks shrubby vegetation.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Salix ovalifolia</i> (2) <i>Empetrum nigrum</i>
Herbaceous	(1) <i>Carex aquatilis</i> (2) <i>Carex glareosa</i>

Physiographic features

This ecological site occurs on high coastal plains within a shore complex. On Nunivak Island, additional associated landforms were identified as margins of lagoons and backshores (Swanson et al. 1986). Slope is nearly level and

occurs on all aspects. Elevation ranges between 10 and 30 feet. Tidal flooding occurs frequently for brief durations. Ponding occurs frequently for long durations of time with ponding depth ranging between 0 and 12 inches. During the growing season, a water table commonly occurs at the soil surface. This site generates negligible runoff to adjacent, downslope ecological sites.

Table 2. Representative physiographic features

Landforms	(1) Lowland > Coastal plain (2) Shore complex > Coastal plain (3) Shore complex > Lagoon (4) Shore complex > Backshore
Runoff class	Negligible
Flooding duration	Brief (2 to 7 days)
Flooding frequency	Frequent
Ponding duration	Long (7 to 30 days)
Ponding frequency	Frequent
Elevation	10–30 ft
Slope	0–2%
Ponding depth	0–12 in
Water table depth	0 in
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

Sea ice strongly influences the climate of the islands in MLRA 239X. For the Northern Bering Sea Islands, sea ice starts forming in December and often persists through early June. In the absence of sea ice, the Bering Sea and North Pacific Ocean moderate diurnal and monthly temperatures resulting in a maritime climate. Summer temperatures (June through August) are relatively stable with mean maximum monthly temperatures ranging between 50 to 55 degrees Fahrenheit. As sea ice forms around the islands, temperatures decrease significantly with the area shifting to a continental climate. The coldest months (January through March) have mean monthly minimum temperatures ranging from 4 to 6 degrees Fahrenheit. The extent, thickness, and duration of the Bering Sea ice appears to be in flux resulting in southerly storms that can bring significantly warmer winter monthly temperatures (Stabeno et al. 2018, Gramling 2019).

The Northern Bering Sea Islands have summers that are short and cool and winters that are long and cold. Strong winds are common throughout the year. Mean annual air temperatures typically range from 26 to 32 degrees Fahrenheit with Saint Lawrence Island (mean annual air temperatures between 24 to 28 degrees Fahrenheit) being significantly colder compared to Nunivak and Saint Michael Islands (mean annual air temperatures between 30 to 34 degrees Fahrenheit) (PRISM 2018). The warmest months are June, July, and August. During these summer months, the typical freeze free period for the area ranges from 94 to 111 days. The coldest months are January, February, and March.

This area is semi-arid with mean annual precipitation typically ranging between 14 and 17 inches. The warmest months have overcast skies with frequent fog and precipitation while the coldest months have clear skies. The two wettest months are August and September where the islands typically receive a quarter of the annual precipitation. The rest of the months receive similar amounts of precipitation. Saint Michael Island receives greater mean annual precipitation (between 17 and 21 inches) compared to Nunivak and Saint Lawrence Islands (between 13 to 17 inches) (PRISM 2018). The average annual snowfall ranges from about 50 to 80 inches (USDA 2022) with the highest snowfall occurring during the months spanning November through March (USDA 1986).

Table 3. Representative climatic features

Frost-free period (characteristic range)	51-75 days
Freeze-free period (characteristic range)	94-111 days

Precipitation total (characteristic range)	14-17 in
Frost-free period (actual range)	50-85 days
Freeze-free period (actual range)	93-117 days
Precipitation total (actual range)	13-21 in
Frost-free period (average)	64 days
Freeze-free period (average)	103 days
Precipitation total (average)	15 in




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) NOME MUNI AP [USW00026617], Nome, AK
- (2) WALES [USW00026618], Wales, AK
- (3) BETHEL AP [USW00026615], Bethel, AK

Influencing water features

This site is classified as an Estuarine Fringe wetland under the Hydrogeomorphic (HGM) classification system (USDA-NRCS 2008). In the associated high coastal plains, periodic tidal flooding, precipitation, and ground water discharge are the main sources of water (USDA-NRCS 2008).

Depth to the water table may decrease following summer storm events or spring snowmelt and increase during extended dry periods.

Wetland description

n/a

Soil features

Soils formed in a thick layer of loamy marine deposits. Rock fragments do not occur on the soil surface. These are mineral soils capped with 7 to 11 inches of saturated organic material. The mineral soil below the organic material is a silt loam, which lacks rock fragments and has high water holding capacity. While soils are considered very deep, permafrost occurs at shallow to moderate depths (12 to 24 inches). The pH of the soil profile is very strongly acidic to moderately acidic. The soils are wet for long portions of the growing season and are considered very poorly drained.

Table 4. Representative soil features

Parent material	(1) Marine deposits
Surface texture	(1) Silt loam
Family particle size	(1) Loamy
Drainage class	Very poorly drained
Permeability class	Moderately rapid
Depth to restrictive layer	12–24 in
Soil depth	60 in
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	8.1–10.8 in
Calcium carbonate equivalent (10-40in)	0%
Clay content (0-20in)	1–5%
Electrical conductivity (10-40in)	0 mmhos/cm
Sodium adsorption ratio (10-40in)	0–3
Soil reaction (1:1 water) (10-40in)	4.8–6
Subsurface fragment volume <=3" (0-60in)	0%
Subsurface fragment volume >3" (0-60in)	0%

Ecological dynamics

The Northern Bering Sea Islands MLRA (MLRA 239X) occurs in the arctic where the harsh climate limits the composition and structure of plant communities. This area has cool and short summers and long and cold winters. Limited warmth during the short summer months, inhibits trees from occurring, and the expansive tundra is composed of a mosaic of low growing shrubs, sedges, moss, and lichen. The cold temperatures limit the vertical and horizontal structure of shrubs and other functional groups of the tundra (CAVM 2022). For instance, shrubs do not typically exceed 80 cm in height across these islands (Swanson et al. 1986; CAVM 2022). The exception is micro-climates where snow can accumulate, which protects vegetation from harsh winds and insulates soils. These micro-climates allow for shrubs to grow much taller and occur on landforms such as flood plains.

Saltwater Influences

Within the shore complex, there are multiple landforms such as tidal flats, foredunes, estuaries, and backshores each with unique tidal influences. Tidal flats, foredunes, and estuaries are all proximal to the ocean and receive frequent salt spray and/or routine tidal flooding. Backshores are comparatively distal landforms that receive tidal

flooding only during exceptionally severe storms or unusually high tides. Based on the associated soils and vegetation, this ecological site occurs on distal landforms within the shore complex.


Ungulate History and Use

In this MLRA, the lack of predators paired with quality forage can lead to dramatic population growth of reindeer which in turn can lead to significant die-offs. Eighty-one reindeer were introduced to Nunivak Island in 1920. Due to a lack of predators and an abundance of high-quality forage, the reindeer population climbed to peaks of greater than 30 thousand in 1944 and 23 thousand in 1965 (Swanson and Barker 1991). After each peak in population, the reindeer herds experienced dramatic population die offs that resulted in less than 5 thousand animals (Swanson and Barker 1991). These die offs are largely attributed to lichen range depletion. Lichen forage makes up 47 percent of the March diet for reindeer herds on Nunivak Island (Swanson et al. 1986) so the depletion of lichen range can directly lead to stress and mortality of reindeer populations.

The presence of introduced ungulate herds on Nunivak, St Lawrence, and St. Matthews Islands, in some places for over a century, plays an integral role in shaping vegetation across this MLRA. Some small islands in this MLRA are believed to have no history of natural or introduced ungulate herds (e.g. Pinnacle Islands, Hall Island, and Punuk Islands). On islands with introduced herds, grazing by reindeer and/or muskox has impacted the potential natural vegetation. For instance, continuous grazing of slow growing fruticose lichen can lead to changes in lichen species composition (Swanson and Barker 1991) and can lead to increases in shrub and bryophyte cover (Kautz et al. 1992). Because of the mixed history in grazing in this MLRA, the STM for this ecological site has two states. Islands in this MLRA without a history of ungulate herds are associated with reference state vegetation, while islands with introduced ungulate herds are associated with grazing state vegetation.

State and transition model


Ecosystem states




T1 - Human introduction of reindeer and/or muskox to islands

R1 - Long periods of time after extirpation of human introduced ungulates

State 1 submodel, plant communities



State 2 submodel, plant communities



State 1 Reference State

The historic and current use of introduced ungulates in this MLRA may have altered the potential natural vegetation

on these islands. Islands in this MLRA without a history of introduced grazing have reference state vegetation, while islands with introduced herds of reindeer and/or muskox (Nunivak, St. Lawrence, and St. Matthews Islands) have grazing state vegetation. Currently no data has been collected in areas of this MLRA in reference condition. Future targeted data collection efforts can address whether range in excellent condition within the grazing state is similar to reference state vegetation and these results could dramatically alter this provisional state and transition model.

Dominant plant species

- black crowberry (*Empetrum nigrum*), shrub
- oval-leaf willow (*Salix ovalifolia*), shrub
- Alaska bog willow (*Salix fuscescens*), shrub
- sedge (*Carex*), grass
- circumpolar reedgrass (*Calamagrostis deschampsoides*), grass
- Schreber's big red stem moss (*Pleurozium schreberi*), other herbaceous
- sphagnum (*Sphagnum*), other herbaceous

Community 1.1

crowberry - oval-leaf willow / sedges - circumpolar reedgrass

Community 1.1 is the potential natural vegetation for this state. It is characterized as halophytic sedge wet meadow (Vioreck et al. 1992). Common and abundant understory species include crowberry, oval-leaf willow, Alaska bog willow, various sedges, and circumpolar reedgrass, and Sphagnum moss.

Dominant plant species

- black crowberry (*Empetrum nigrum*), shrub
- oval-leaf willow (*Salix ovalifolia*), shrub
- Alaska bog willow (*Salix fuscescens*), shrub
- sedge (*Carex*), grass
- circumpolar reedgrass (*Calamagrostis deschampsoides*), grass
- sphagnum (*Sphagnum*), other herbaceous

State 2

Grazing State



Figure 7. A shore complex adjacent to Duchikthluk Bay on the southern end of Nunivak Island.

One plant community has been documented within the grazing state. The data for this state is based on a historical range survey conducted on Nunivak Island (Swanson et al. 1986) Future work will be required to determine if the vegetation on Nunivak Island represent the vegetation across the grazed islands of this MLRA.

Community 2.1

oval-leaf willow - crowberry / water sedge - lesser saltmarsh sedge / Schreber's big red stem moss - Sphagnum moss



Figure 8. A tidal marsh associated with this ecological site on Nunivak Island.

One plant community has been documented within the grazing state for this ecological site. This community is characterized as halophytic sedge wet meadow (Vioreck et al. 1992) with the dominant plants being oval-leaf willow, crowberry, water sedge, lesser saltmarsh sedge, Schreber's big red stem moss, and Sphagnum moss. Other commonly observed species include red fescue, arctic daisy, purple marshlocks, pacific silverweed, saltmarsh starwort, Schreber's big red stem moss, and Sphagnum moss. The vegetative strata that characterize this community are dwarf shrubs (less than 8 inches height), medium graminoids (between 4 and 24 inches in height) and moss.

Forest understory. Live lichen and moss annual production cannot be measured accurately due to a lack of information on growth rates and/or slow annual growth rates. Lichen and moss biomass data below refers to total biomass, while vascular plants biomass refers to annual production.

Dominant plant species

- oval-leaf willow (*Salix ovalifolia*), shrub
- black crowberry (*Empetrum nigrum*), shrub
- arctic willow (*Salix arctica*), shrub
- netleaf willow (*Salix reticulata*), shrub
- water sedge (*Carex aquatilis*), grass
- lesser saltmarsh sedge (*Carex glareosa*), grass
- red fescue (*Festuca rubra*), grass
- arctic daisy (*Chrysanthemum arcticum*), other herbaceous
- purple marshlocks (*Comarum palustre*), other herbaceous
- Pacific silverweed (*Argentina egedii* ssp. *groenlandica*), other herbaceous
- saltmarsh starwort (*Stellaria humifusa*), other herbaceous
- Schreber's big red stem moss (*Pleurozium schreberi*), other herbaceous
- sphagnum (*Sphagnum*), other herbaceous

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	350	490	630
Moss	200	300	400
Shrub/Vine	100	140	180
Forb	50	70	90
Lichen	0	0	0
Total	700	1000	1300

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	5-15%
Grass/grasslike foliar cover	50-75%
Forb foliar cover	10-25%
Non-vascular plants	5-15%
Biological crusts	0%
Litter	50-75%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 7. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	5-15%	—	—
>0.5 <= 1	—	—	—	10-25%
>1 <= 2	—	—	50-75%	—
>2 <= 4.5	—	—	—	—
>4.5 <= 13	—	—	—	—
>13 <= 40	—	—	—	—
>40 <= 80	—	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

Transition T1

State 1 to 2

Human introduction of reindeer and/or muskox to islands

Restoration pathway R1

State 2 to 1

Long periods of time after extirpation of human introduced ungulates

Additional community tables

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Shrub/Vine					
1	Shrub Total Annual Production			100–180	
	oval-leaf willow	SAOV	<i>Salix ovalifolia</i>	40–110	–
	black crowberry	EMNI	<i>Empetrum nigrum</i>	10–55	–
	arctic willow	SAAR27	<i>Salix arctica</i>	10–55	–
	netleaf willow	SARE2	<i>Salix reticulata</i>	5–20	–
Grass/Grasslike					
2	Graminoid Total Annual Production			350–630	
	water sedge	CAAQ	<i>Carex aquatilis</i>	100–315	–
	lesser saltmarsh sedge	CAGL4	<i>Carex glareosa</i>	100–270	–
	red fescue	FERU2	<i>Festuca rubra</i>	25–90	–
	tall cottongrass	ERAN6	<i>Eriophorum angustifolium</i>	0–45	–
	rush	JUNCU	<i>Juncus</i>	5–30	–
Forb					
3	Forb Total Annual Production			50–90	
	Pacific silverweed	AREGG	<i>Argentina egedii</i> ssp. <i>groenlandica</i>	10–55	–
	purple marshlocks	COPA28	<i>Comarum palustre</i>	10–55	–
	arctic daisy	CHAR13	<i>Chrysanthemum arcticum</i>	5–35	–
	saltmarsh starwort	STHU	<i>Stellaria humifusa</i>	5–30	–
	ledge stonecrop	RHIN11	<i>Rhodiola integrifolia</i>	0–10	–
Moss					
4	Total Bryophyte Biomass			200–400	
	Schreber's big red stem moss	PLSC70	<i>Pleurozium schreberi</i>	80–240	–
	sphagnum	SPHAG2	<i>Sphagnum</i>	80–240	–
Lichen					
5	Total Lichen Biomass			0	

Animal community

The principal use of this site by wildlife is feeding for a variety of shorebirds and waterfowl. Brant, emperor geese, turnstones, and sandhill cranes commonly use this site for nesting and feeding, while a large variety of other bird species may forage and rest in these areas. Muskoxen also use this site in addition to reindeer, especially during the summer to escape from oestrid flies; the coastal breezes can provide escape from these insects.

Hydrological functions

n/a

Recreational uses

The varied seascapes and landscapes present scenic views of high aesthetic value. Opportunities for photography are excellent.

Wood products

No wood products available from this site.

Other products

Reindeer Grazing

This site is best suited for spring and summer range. Growing grasses and sedges produce high levels of digestible protein. Browse also provides forage during the fall and early winter. Some sedges may provide high quality forage during the winter.

Other information

These interpretive narratives were all developed in a report for range sites on Nunivak Island (Swanson et al.1986).

Inventory data references

Plant species and production information are largely based on a historic range survey on Nunivak Island (Swanson et al. 1986).

References

Alaska Department of Fish and Game Staff. 2024 (Date accessed). Muskox (*Ovibos moschatus*).
<https://www.adfg.alaska.gov/index.cfm?adfg=muskox.main>.

CAVM Team. 2023. Raster Circumpolar Arctic Vegetation Map. Scale 1:7,000,000. Conservation of Arctic Flora and Fauna, Akureyri.

Gramling, C. 2019. What happens when the Bering Sea's ice disappears?.

Griffin, D. 2001. Nunivak Island, Alaska: A history of contact and trade.

Jackson, S. 1902. Eleventh Annual Report on Introduction of Domestic Reindeer into Alaska, with Map and Illustrations, 1901. Senate of the United States, 54th Congress, 1st Session. Document No. 98..

Kautz RD, Swanson JD, Barker M, and Morgart J. 1992. Nunivak Island Trend Study, 1989-1990 Nunivak Island, AK. USDA NRCS.

Landfire. 2009. Biophysical Setting. LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior, Washington, DC..

Patton Jr, W., F. Wilson, and T. Taylor. 2011. Geologic Map of Saint Lawrence Island.

Smith, R.D., A.P. Ammann, C.C. Bartoldus, and M.M. Brinson. 1995. An approach for assessing wetland functions using hydrogeomorphic classification, reference wetlands, and functional indices.

Stabeno, P.J., C.A. Ladd, C. Mordy, and R.M. McCabe. 2018. How the Absence of Sea Ice Altered the Physical Oceanography of the Northern Bering Sea. AGU Fall Meeting Abstracts 2018:OS53B–04.

Swanson, J.D., D. Lehner, J. Zimmerman, and D. Pauling. 1986. Range survey of Nunivak Island, Alaska.

Swanson, J. and M. Barker. 1992. Assessment of Alaska reindeer populations and range conditions. Rangifer 12:33–43.

United States Department of Agriculture, . 2022. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin.

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

Wald, E.J. 2009. Nunivak Island Reindeer and Muskoxen Survey, 2009. US Fish & Wildlife Service, Yukon Delta National Wildlife Refuge.

Zeusler, F. 2009. Ice in the Bering Sea and Arctic Ocean.

Other references

PRISM Climate Group. 2018. Alaska – average monthly and annual precipitation and minimum, maximum, and mean temperature for the period 1981-2010. Oregon State University, Corvallis, Oregon. <https://prism.oregonstate.edu/projects/alaska.php>. (Accessed 4 September 2019).

Scenarios network for Alaska and arctic planning (SNAP). Historical Monthly Temperature – 1km, 1901-2009. <http://ckan.snap.uaf.edu/dataset/>. (Accessed 5 May 2021).

SNAP. Historical monthly and derived precipitation products downscaled from CRU TS data via the delta methods – 2km, 1901-2009. <http://ckan.snap.uaf.edu/dataset/>. (Accessed 5 May 2021).

Contributors

Blaine Spellman

Approval

Marji Patz, 2/18/2025

Acknowledgments

Phillip Barber, Stephanie Schmit, Michael Singer, Jamin Johanson and Marji Patz are acknowledged for their feedback on the ecological sites of this MLRA and suggestion on soil component correlation. Karin Sonnen is acknowledged for her excellent technical review to make this ecological site description report a much better product.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
--------------------------	--

Contact for lead author	
Date	05/11/2025
Approved by	Marji Patz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-