

# **Ecological site group 029XESG05**

## **Finer Uplands**

Last updated: 04/18/2023  
Accessed: 05/10/2025

---

### **Key Characteristics**

- Uplands
- <75% bedrock outcrop
- Surface SAR <8 and Subsurface EC <8 and Surface EC <4
- EC <1.5 surface and <2 subsurface
- slope <35% or <40% surface rock
- Depth >55cm
- Rock <30% surface and <30% subsurface
- Clay <30% surface and <35% subsurface
- sand <75% and texture is Sandy Loam or finer in surface & subsurface
- Clay >20% and texture is finer than Sandy Loam in surface

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### **Physiography**

This ESG is located on alluvial fans, inset fans, and mountains.

### **Vegetation dynamics**

The modal ecological site for this ESG is R029XY048NV Outwash Plain.

### **Major Land Resource Area**

MLRA 029X  
Southern Nevada Basin and Range

### **Subclasses**

- R029XG020CA–Moist Floodplain
- R029XY006NV–LOAMY 8-10 P.Z.
- R029XY017NV–LOAMY 5-8 P.Z.
- R029XY048NV–OUTWASH PLAIN
- R029XY114NV–LOAMY FAN 8-12 P.Z.
- R029XY310UT–Upland Loam (Utah Serviceberry)

### **Correlated Map Unit Components**

23237362, 23237253, 22663252, 23152931, 23157400, 23158048, 22816731, 22822266, 22822222, 22822455, 22822462, 22601393, 22728266, 22728267

### **Stage**

Provisional

## **Contributors**

Curtis Talbot  
Travis Nauman

## **State and transition model**

## **Citations**