Major Land Resource Area 009X Palouse and Nez Perce Prairies

Accessed: 05/12/2025

Ecological site keys

MLRA 9 - Southern Section - LRU A - Basalt plateaus and high basins

- I. Sites found in open drainageways or swales receiving supplemental subsurface flow
 - A. Site predominately found within a 14-17" precipitation zone
 - 1 Soils typically silt loam over silty clay loam ... R009XY005OR Cold Swale 13-17 PZ
 - 2 Soils typically silty clay loam over clay ... R009XY006OR Cold Clayey Swale 13-17 PZ
 - B. Site predominately found within a 17-24" precipitation zone
 - 1 Soils shallow (10 20"), somewhat poorly drained ... R009XY007OR Cold Swale 17-24 PZ
 - 2 Soils moderately deep (20 40"), poorly drained ... R009XY008OR Cold Wet Swale 17-24 PZ

II. Not as above

- A. Site predominately found within a 13 17" precipitation zone
 - 1 Sites flat to moderately sloping, not influenced by aspect
 - i. Soils very shallow (0 10") ... R009XY027OR Cold Very Shallow 13+ PZ
 - ii. Soils shallow (10 20") ... R009XY022OR Cold Shallow 13+ PZ
 - iii. Soils moderately deep (20 40") ... R009XY017OR Cold Loamy 13-17 PZ
 - iv. Soils deep to very deep (40" +) .. R009XY205OR Cold Deep Loamy 13-17 PZ (missing concept)
 - 2 Sites gently to steeply sloping, influenced by aspect
 - i. Sites occur on south slope aspects
 - a. Soils shallow (10 20") ... R009XY036OR Cold Shallow South 13+ PZ
 - b. Soils moderately deep to very deep (20" +)
 - 1) Slopes 12 30% ... R009XY034OR Cold South 13-17 PZ
 - 2) Slopes 30 90% .. R009XY203OR Cold Steep South 13-17 PZ (missing concept)
 - ii. Sites occur on north slope aspects
 - a. Soils shallow (10 20") ... R009XY048OR Shallow North 14+ PZ
 - b. Soils moderately deep to very deep (20" +)
 - 1) Clayey family particle size class, shrubs dominated by Symphorocarpus and Rosa spp. ... R009XY041OR Deep North 14-17 PZ
 - 2) Loamy family particle size class
 - a) Stony silt loam surface texture, shrubs dominated by Physocarpus malvaceus ... R009XY060OR Shrubby North 15+ PZ
 - b) Silt loam surface texture, shrubs dominated by Crataegus spp. ... R009XY046OR Shrubby Moist North 15+ PZ
- B. Site predominately found within a 17 24" precipitation zone
 - 2 Sites flat to moderately sloping, not influenced by aspect
 - i. Soils very shallow (0 10") ... R009XY027OR Cold Very Shallow 13+ PZ
 - ii. Soils shallow (10 20") ... R009XY022OR Cold Shallow 13+ PZ
 - iii. Soils moderately deep (20 40") ... R009XY018OR Cold Loamy 17-24 PZ

- iv. Soils deep to very deep (40" +) .. R009XY206OR Cold Deep Loamy 17-24 PZ (missing concept)
- 1 Sites gently to steeply sloping, influenced by aspect
 - i. Sites occur on south slope aspects
 - a. Soils shallow (10 20") ... R009XY036OR Cold Shallow South 13+ PZ
 - b. Soils moderately deep to very deep (20" +)
 - 1) Slopes 30 60% ... R009XY035OR Cold South 17-24 PZ
 - 2) Slopes 60 90% .. R009XY204OR Cold Steep South 17-24 PZ (missing concept)
 - ii. Sites occur on north slope aspects
 - a. Soils very shallow to shallow (0 20")
 - 1) Soils very shallow (0-10") .. R009XY202OR Very Shallow North 14+ PZ (missing concept)
 - 2) Soils shallow (10 20") ... R009XY048OR Shallow North 14+ PZ
 - b. Soils moderately deep to very deep (20" +)
 - 1) Clayey family particle size class, shrubs dominated by Symphorocarpus and Rosa spp. ... $R009XY045OR-North\ 17-24\ PZ$
 - 2) Loamy family particle size class
 - a) Stony silt loam surface texture, shrubs dominated by Physocarpus malvaceus ... R009XY060OR Shrubby North 15+ PZ
 - b) Silt loam surface texture, shrubs dominated by Crataegus spp. ... R009XY046OR Shrubby Moist North 15+ PZ
- C. Site predominately found within a 30-40" precipitation zone. R009XY070OR. ... R009XY070OR High Ridge 30+ PZ